

COLD AND COUGH- diphenhydramine hcl, phenylephrine hcl solution

Meijer Distribution Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Meijer Distribution, Inc. Cold & Cough Drug Facts

Active ingredients (in each 5 mL)

Diphenhydramine HCl 6.25 mg

Phenylephrine HCl 2.5 mg

Purposes

Antihistamine/cough suppressant

Nasal decongestant

Uses

- temporarily relieves:
- sneezing
- itchy nose or throat
- runny nose
- itchy, watery eyes due to hay fever
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation as may occur with a cold

Warnings

Do not use

- in a child under 4 years of age
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- with any other product containing diphenhydramine, even one used on skin
- to make a child sleepy

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)

- chronic cough that lasts or as occurs with asthma
- a breathing problem such as chronic bronchitis

Ask a doctor or pharmacist before use if the child is
taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- may cause marked drowsiness
- sedatives and tranquilizers may increase drowsiness
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or occur with fever
- cough persists for more than 7 days, comes back, or occurs with fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- may be given every 4 hours. Do not give more than 6 doses in 24 hours unless directed by a doctor.
- use enclosed dosing cup only. Keep for use with this product only. Do not use any other dosing device.

Age	Dose
children under 4 years of age	do not use
children 4 to under 6 years of age	do not use unless directed by a doctor
children 6 to under 12 years of age	10 mL

Other information

- **each 5 mL contains:** sodium 3 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

acesulfame potassium, anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, maltitol solution, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Children's Triaminic® Night Time Cold & Cough active ingredients

For Ages 6-12 Years

CHILDREN'S

nighttime cold & cough

Diphenhydramine HCl

Antihistamine | Cough suppressant

Phenylephrine HCl | Nasal decongestant

Multi-Symptom Relief

Grape Flavor

Cough | Runny, Stuffy Nose

Itchy Throat

Sugar Free | Alcohol Free

4 FL OZ (118 mL)



COLD AND COUGH

diphenhydramine hcl, phenylephrine hcl solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-913
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Drug Facts

Active ingredients (in each 5 mL)
 Diphenhydramine HCl 6.25 mg.....Antihistamine/cough suppressant
 Phenylephrine HCl 2.5 mg.....Nasal decongestant

Purposes

Uses
 temporarily relieves:
 ■ sneezing ■ itchy nose or throat
 ■ runny nose ■ itchy, watery eyes due to hay fever
 ■ nasal and sinus congestion
 ■ cough due to minor throat and bronchial irritation as may occur with a cold

Warnings
Do not use
 ■ in a child under 4 years of age
 ■ in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
 ■ with any other product containing diphenhydramine, even one used on skin
 ■ to make a child sleepy

Ask a doctor before use if the child has
 ■ heart disease ■ high blood pressure
 ■ thyroid disease ■ diabetes ■ glaucoma
 ■ cough that occurs with too much phlegm (mucus)
 ■ chronic cough that lasts or as occurs with asthma
 ■ a breathing problem such as chronic bronchitis

Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers

When using this product
 ■ do not exceed recommended dosage
 ■ may cause marked drowsiness
 ■ sedatives and tranquilizers may increase drowsiness
 ■ excitability may occur, especially in children

Stop use and ask a doctor if
 ■ nervousness, dizziness or sleeplessness occur
 ■ symptoms do not improve within 7 days or occur with fever
 ■ cough persists for more than 7 days, comes back, or occurs with fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Directions
 ■ may be given every 4 hours. Do not give more than 6 doses in 24 hours unless directed by a doctor.
 ■ use enclosed dosing cup only. Keep for use with this product only. Do not use any other dosing device.

Age	Dose
children under 4 years of age	do not use
children 4 to under 6 years of age	do not use unless directed by a doctor
children 6 to under 12 years of age	10 mL

Other information
 ■ each 5 mL contains: sodium 3 mg
 ■ store at 20-25°C (68-77°F)

Inactive ingredients: acesulfame potassium, anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, maltitol solution, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by Novartis Consumer Health S.A., owner of the registered trademark Triaminic®.

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

DIST. BY MEIJER DISTRIBUTION, INC.
 GRAND RAPIDS, MI 49544
 www.meijer.com



PID 416721

91326 6E C5

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	6.25 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	

Product Characteristics

Color	PURPLE	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-913-26	1 in 1 CARTON	10/31/2007	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/31/2007	

Labeler - Meijer Distribution Inc (006959555)

Revised: 1/2019

Meijer Distribution Inc