

**CURANOL- acetaminophen suspension**  
**Wittman Pharma, Inc.**

-----  
**CuraNOL**

**Directions:**

- Dosage may be repeated every 4 hours, or as directed by doctor
- Do not exceed recommended dose.
- Find dose in chart below ■ If possible, use weight to dose, otherwise use age

**Dosage**

Weight (lbs)	Under 24	24-35	36-47	48-59	60-71	72-95	Over 95
Age (years)	Under 2	2 to 3	4 to5	6 to 8	9 to 10	11	12 to Adult
Dosage Teaspoonful (tsp)	consult a doctor	1 tsp (5 mL)	1 1/2 tsp (7.5 mL)	2 tsp (10mL)	2 1/2 tsp (12.5 mL)	3 tsp (15 mL)	4 tsp (20mL)

**Warnings:** Liver Warning: This product contains acetaminophen.

Severe liver damage may occur if.

- adults take more than 6 doses in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen (prescription or nonprescription)
- adult has 3 or more alcoholic drinks while using this product.

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■rash

If skin reaction occurs, stop use and seek medical help right away

Do not use with any other product containing acetaminophen (prescription or nonprescription), this will provide more than the recommended dose (overdose) of acetaminophen and could cause serious health concerns. If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**WHEN USING THIS PRODUCT, DO NOT EXCEED RECOMMENDED DOSE**

Ask a doctor before use if you have a liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask doctor if: ■ pain gets worse or lasts more than 10 days in adults  
■ pain gets worse or lasts more than 5 days in children under 12 years

fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition  
If pregnant or breast-feeding, ask a health professional before use.

**Inactive ingredients:** Citric Acid, Flavor, Methylparaben, Monoammonium

Glycyrrhizinate, Potassium Citrate, Propylene Glycol, Propylparaben, Purified Water, Sorbitol,

Sucralose

**Uses:** Temporarily relieves minor aches and pains due to: ■ headache ■ backache ■ muscular aches ■ minor pain of arthritis ■ the common cold ■ toothache ■ premenstrual and menstrual cramps

Temporarily reduces fever.

**Keep out of reach of children.**

**Purpose:**

Pain reliever/Fever reducer

**Active Ingredient** (in each 5 mL tsp)

Acetaminophen, 160mg

<b>Drug Facts</b>	
<b>Active Ingredient</b> (in each 5 mL tsp)	<b>Purpose</b>
Acetaminophen, 160 mg	Pain reliever/Fever reducer
<b>Uses:</b> Temporarily relieves minor aches and pains due to: ■ headache ■ backache ■ muscular aches ■ minor pain of arthritis ■ the common cold ■ toothache ■ premenstrual and menstrual cramps. Temporarily reduces fever.	
<b>Warnings: Liver Warning:</b> This product contains acetaminophen. Severe liver damage may occur if: ■ adults take more than 6 doses in 24 hours, which is the maximum daily amount ■ child takes more than 5 doses in 24 hours, which is the maximum daily amount ■ taken with other drugs containing acetaminophen (prescription or nonprescription) ■ adult has 3 or more alcoholic drinks daily, while using this product.	
<b>Allergy Alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.	
<b>Do not use</b> with any other product containing acetaminophen (prescription or nonprescription), this will provide more than the recommended dose (overdose) of acetaminophen and could cause serious health concerns. If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	
<b>WHEN USING THIS PRODUCT, DO NOT EXCEED RECOMMENDED DOSE</b> <b>Ask a doctor before use if you have a liver disease.</b> Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.	
<b>Stop use and ask doctor if:</b> ■ pain gets worse or lasts more than 10 days in adults ■ pain gets worse or lasts more than 5 days in children under 12 years, and before fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.	

MADE IN USA NDC 83335-104-16



# CuraNOL

Pain Reliever / Fever Reducer

**ACETAMINOPHEN**  
**160 mg / 5 mL**

Compare to the active ingredient of Tylenol® Liquid\*


✓ SUGAR FREE    ✓ ALCOHOL FREE  
 ✓ DYE FREE      ✓ GLUTEN FREE

Grape Flavor



**Wittman Pharma**

Net Content: 16 fl oz (473 mL)

<b>Drug Facts (continued)</b>							
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults/children even if you do not notice any signs or symptoms.							
<b>Directions:</b> ■ Dosage may be repeated every 4 hours, or as directed by doctor. ■ Do not exceed recommended dose. ■ Find dose in chart below. ■ If possible, use weight to dose, otherwise use age.							
Weight (lbs)	Under 24	24-35	36-47	48-59	60-71	72-95	Over 95
Age (years)	Under 2	2 to 3	4 to 5	6 to 8	9 to 10	11	12 to adult
Dosage Teaspoonful (tsp)	consult a doctor	1 tsp (5 mL)	1½ tsp (7.5 mL)	2 tsp (10 mL)	2½ tsp (12.5 mL)	3 tsp (15 mL)	4 tsp (20 mL)
<b>Other information:</b> ■ Store at 15°-30°C (59°-86°F). Do not refrigerate. ■ Protect from light. Avoid excessive heat or humidity. ■ Tamper evident, do not use if safety seal under cap is broken or missing.							
<b>Inactive ingredients:</b> Citric Acid, Flavor, Methylparaben, Monoammonium Glycyrrhizinate, Potassium Citrate, Propylene Glycol, Propylparaben, Purified Water, Sorbitol, Sucralose							
<b>Questions?</b> Call 1-352-549-9917 M-F 9am - 5pm EST Manufactured for: Wittman Pharma, Inc. Brooksville, FL 34604							
*This product is not manufactured or distributed by McNeil Consumer Healthcare distributor of Tylenol® Liquid. Rev. 3/2024							
 3 83335 10416 3							

## CURANOL

acetaminophen suspension

### Product Information

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:83335-104

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
METHYLPARABEN (UNII: A2I8C7HI9T)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SORBITOL (UNII: 506T60A25R)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

### Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83335-104-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/01/2023	
2	NDC:83335-104-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/01/2023	
3	NDC:83335-104-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/01/2023	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/01/2023	

Labeler - Wittman Pharma, Inc. (830980947)

Establishment

Name	Address	ID/FEI	Business Operations
Wittman Pharma, Inc.		830980947	label(83335-104) , manufacture(83335-104) , analysis(83335-104)

Revised: 4/2024

Wittman Pharma, Inc.