

CURELIEF- diphenhydramine hcl solution
Wittman Pharma, Inc.

CuRELIEF

Directions: Take every 4-6 hours, or as directed by doctor. * Do NOT exceed the recommended dose

Age	Dose
Adults and Children 12 years of age and over	take 10 to 20 mL (2 to 4 tsp), not to exceed 300 mg in 24 hours
Children 6 to under 12 years of age	take 5 to 10 mL (1 to 2 tsp) not to exceed 150 mg in 24 hours
Children under 6 years of age	Consult a doctor

Warnings:

Do not use: ■ with any other product containing diphenhydramine, even one used on skin ■ to make a child sleepy

Ask a doctor before use if you have: ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to the enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product: ■ excitability might occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness effect ■ be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Inactive Ingredients: Citric Acid, Flavor, Methylparaben, Monoammonium

Glycyrrhizinate, Potassium Citrate, Propylene Glycol, Propylparaben, Purified Water, Sorbitol, Sucralose

Uses: Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat ■ Temporarily relieves these symptoms due to the common cold: ■ runny nose ■ sneezing

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Purpose: Antihistamine

Active Ingredient (in each 5 mL tsp)

Diphenhydramine HCL, 12.5 mg

Drug Facts

Drug Facts	
Active Ingredient (in each 5 mL tsp)	Purpose
Diphenhydramine HCL, 12.5 mg	Antihistamine
Uses: Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat ■ Temporarily relieves these symptoms due to the common cold: ■ runny nose ■ sneezing	
Warnings: Do not use: ■ with any other product containing diphenhydramine, even one used on skin ■ to make a child sleepy Ask a doctor before use if you have: ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to the enlarged prostate gland	
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. When using this product: ■ excitability might occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness effect ■ be careful when driving a motor vehicle or operating machinery	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children.	
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	

MADE IN USA NDC 83335-103-16



CuRELIEF

ANTIHISTAMINE

Diphenhydramine HCL
12.5 mg / 5 mL

Compare to the active ingredient of
Benadryl® Allergy Liquid*


✓ SUGAR FREE
✓ ALCOHOL FREE

✓ DYE FREE
✓ GLUTEN FREE

Cherry Flavor


**Wittman
Pharma**

Net Content: 16 fl oz (473 mL)

Drug Facts (continued)	
Directions: ■ Take every 4 to 6 hours, or as directed by doctor. ■ Do not exceed recommended dose.	
Age	Dose
Adults and children 12 years of age and over:	take 10 to 20 mL (2 to 4 tsp) not to exceed 300 mg in 24 hrs
Children 6 to under 12 years of age:	take 5 to 10 mL (1 to 2 tsp) not to exceed 150 mg in 24 hrs
Children under 6 years of age:	consult a doctor
Other information: ■ Store at 15°-30°C (59°-86°F). ■ Do not refrigerate. ■ Avoid excessive heat or humidity. ■ Protect from light. ■ Tamper evident, do not use if safety seal under cap is broken or missing.	
Inactive Ingredients: Citric Acid, Flavor, Methylparaben, Monoammonium Glycyrhizinate, Potassium Citrate, Propylene Glycol, Propylparaben, Purified Water, Sorbitol, Sucralose	
Questions? Call 1-352-549-9917 M-F 9am - 5pm EST. Manufactured for: Wittman Pharma, Inc. Brooksville, FL 34604	
*Benadryl® is a registered trademark of Johnson & Johnson. Rev. 3/2024	
 3 83335 10316 6	

CURELIEF

diphenhydramine hcl solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:83335-103

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SORBITOL (UNII: 506T60A25R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83335-103-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/2024	
2	NDC:83335-103-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/01/2024	

Labeler - Wittman Pharma, Inc. (830980947)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Revised: 4/2024

Wittman Pharma, Inc.