

SEVERE COLD AND FLU NIGHTTIME- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hci liquid
P & L Development, LLC

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal Decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- runny nose and sneezing
- cough to help you sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed (see overdose warning)**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or last more than 3 days
- redness or swelling is present
- cough comes back, or occurs with rash or headache that lasts.

These could be a signs of a serious conditions.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **take only as directed - see Overdose warning**
- mL = milliliter
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- do not exceed 4 doses per 24 hours
- adults and children 12 years and over: 30 mL every 4 hours
- children under 12 years of age: do not use
- **When using Day Time or Night Time products, carefully read each label to ensure correct dosing.**

Other information

- **each 30 mL contains;** sodium 79 mg
- store between 15-30°C (59-86°F)
- do not refrigerate

Inactive ingredients

anhydrous citric acid, FD&C blue1, FD&C red 40, Flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

Compare to active ingredients in Vicks® NyQuil® Severe Cold & Flu*

maximum strength

severe night time cold & flu

cold & flu

Acetaminophen

dextromethorphan HBr

doxylamine succinate

phenylephrine HCl

relieves:

- aches, fever & sore throat
- cough
- runny nose & sneezing
- nasal & sinus Congestion

alcohol free

berry flavor

FL OZ (mL)

*This product is not manufactured or distributed by Procter & Gamble, distributor or Vicks® NyQuil® Severe Cold & Flu.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL IS BROKEN OR MISSING.

Manufactured by:

PL Developments

11865 S. Alameda St

Lynwood, CA 90262

Product Label

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL IS BROKEN OR MISSING.

PLD-A341A LB003625

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Severe Cold & Flu.

Manufactured by:
PL Developments
11865 S. Alameda St
Lynwood, CA 90262



Compare to active ingredients in
Vicks® NyQuil® Severe
Cold & Flu*
NDC 49580-0416-8

maximum strength
**severe
night time
cold & flu**
Acetaminophen
dextromethorphan HBr
doxylamine succinate
phenylephrine HCl

- relieves:
- aches, fever & sore throat
 - cough
 - runny nose & sneezing
 - nasal & sinus congestion



alcohol free
8 fl oz (237 mL)

Drug Facts	
Active ingredients (in each 30 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine
Phenylephrine HCl 10 mg.....	Nasal decongestant
Uses • temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion and pressure • cough due to minor throat and bronchial irritation • minor aches and pains • headache • fever • sore throat • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage • runny nose and sneezing • cough to help you sleep	
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: • more than 4,000 mg in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	

Drug Facts (continued under label) PEEL HERE →

Drug Facts (continued)
Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema • a sodium- restricted diet
Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin
When using this product • do not take more than directed • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness
Stop use and ask a doctor if • nervousness, dizziness, or sleeplessness occur • pain, nasal congestion, or cough gets worse or lasts more than 7 days • new symptoms occur • fever gets worse or lasts more than 3 days • redness or swelling is present • cough comes back, or occurs with rash or headache that lasts.

Drug Facts (continued)
These could be signs of a serious condition.
If pregnant or breast-feeding , ask a health professional before use.
Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.
Directions • take only as directed - see Overdose warning • mL = milliliter • use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device. • do not exceed 4 doses per 24 hours • adults and children 12 years and over: 30 mL every 4 hours • children under 12 years of age: do not use • when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing
Other information • each 30 mL contains: sodium 79 mg • store between 15-30°C (59-86°F) • do not refrigerate
Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum
Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Readyincase NightTime Severe Cold & Flu Berry Liquid

SEVERE COLD AND FLU NIGHTTIME			
acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hci liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49580-0416
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
			650 mg

ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49580-0416-8	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/31/2015	

Labeler - P & L Development, LLC (101896231)