

**COLD AND FLU NIGHTTIME- acetaminophen, dextromethorphan hbr,
doxylamine succinate solution**
Lidl US LLC

Lidl 44-014

Active ingredients (in each 30 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 30 mg
Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - runny nose and sneezing
 - sore throat
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- liver disease
- glaucoma
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

- do not exceed 4 doses per 24 hours
- mL = milliliter
- only use the dose cup provided
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years: do not use

Other information

- **each 30 mL contains:** sodium 22 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, D&C yellow #10, FD&C green #3, FD&C yellow #6, flavors, glycerin, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate dihydrate, sodium saccharin, sucralose

Questions or comments?

1-800-426-9391

Principal display panel

*Compare to the active ingredients in Vicks[®] NyQuil[®] Cold & Flu

NDC 71141-180-32

RELIEVES Headache, Fever, Sore Throat, Minor Aches & Pains, Sneezing, Runny Nose, Cough

LIQUID

Nighttime
cold & flu
RELIEF

Acetaminophen

Pain Reliever/Fever Reducer

Dextromethorphan HBr
Cough Suppressant

Doxylamine Succinate
Antihistamine

For ages 12 years

and over
Nighttime relief

12 FL OZ (355 mL)

by **LIDL®**

**TAMPER EVIDENT: DO NOT USE IF PRINTED
NECK WRAP IS BROKEN OR MISSING**

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil® Cold & Flu.

50844 ORG032301402

DISTRIBUTED BY
LIDL US, LLC
ARLINGTON, VA 22202

1-844-344-5071
www.lidl.com

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts (continued)

- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil® Cold & Flu. 50844 ORG032301402

Discard Seal, Empty & Recycle Cap. www.recycling.com

PLASTIC BOTTLE

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

5117817A

**No Print / No Varnish Area
Lot # and Exp. Info**

PEEL HERE FOR MORE DRUG FACTS

*Compare to the active ingredients in Vicks® NyQuil® Cold & Flu

NDC 71141-180-32

RELIEVES Headache, Fever, Sore Throat, Minor Aches & Pains, Sneezing, Runny Nose, Cough

Liquid

Nighttime cold & flu RELIEF

Acetaminophen Pain Reliever/Fever Reducer
Dextromethorphan HBr Cough Suppressant
Doxylamine Succinate Antihistamine

For ages 12 years and over
Nighttime relief

12 FL OZ (355 mL)

by **LIDL**®

Drug Facts TAMPER EVIDENT: DO NOT USE IF PRINTED NECK WRAP IS BROKEN OR MISSING

Active ingredients (in each 30 mL)

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
- runny nose and sneezing
- sore throat
- headache
- minor aches and pains
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- liver disease
- glaucoma

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- do not exceed 4 doses per 24 hours
- mL = milliliter
- only use the dose cup provided
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years: do not use

Other information

- each 30 mL contains: sodium 22 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients anhydrous citric acid, D&C yellow #10, FD&C green #3, FD&C yellow #6, flavors, glycerin, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate dihydrate, sodium saccharin, sucralose

Questions or comments? 1-800-426-9391

LiDL 44-014

COLD AND FLU NIGHTTIME

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:71141-180

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	green	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71141-180-32	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/07/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/07/2023	

Labeler - Lidl US LLC (079389709)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(71141-180) , pack(71141-180)

Revised: 4/2024

Lidl US LLC