# LAXATIVE- bisacodyl tablet, delayed release TOPCO ASSOCIATES LLC

-----

#### TopCare 44-676A

#### Active ingredient (in each tablet)

Bisacodyl USP, 5 mg

## Purpose

Stimulant laxative

#### Uses

- for relief of occasional constipation (irregularity)
- this product generally produces bowel movement in 6 to 12 hours

#### Warnings

#### Do not use

if you cannot swallow without chewing.

# Ask a doctor before use if you have

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

# When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness, and cramps

# Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative.
   These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

# If pregnant or breast-feeding,

ask a health professional before use.

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

#### do not take more than directed

take with a glass of water

adults and children 12 years	take 1 to 3 tablets in a single
	daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
children o to under 12 years	dose
	ask a doctor

#### Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- see end flap for expiration date and lot number

## Inactive ingredients

carmine, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

#### Questions or comments?

1-888-423-0139

#### **Principal Display Panel**

+TopCare®

health

NDC 76162-676-01

#### Gentle

Laxative

BISACODYL USP 5 mg • STIMULANT LAXATIVE

Gentle, Dependable, Relief from Constipation

#### **30**

ENTERIC COATED TABLETS

actual size

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844 REV0923B67601

DISTRIBUTED BY
TOPCO ASSOCIATES LLC
ITASCA, IL 60143
©TOPCO LNKA1223
QUESTIONS? 1-888-423-0139
topcare@topco.com
www.topcarebrand.com
QUALITY GUARANTEED

medical help or contact a Poison Control Center right away. Keep out of reach of children. In case of overdose, get

If pregnant or breast-feeding, ask a health professional

- you need to use a laxative for more than 1 week condition.
- after use of a laxative. These could be signs of a serious ■ you have rectal bleeding or fail to have a bowel movement Stop use and ask a doctor if
- it may cause stomach discomfort, faintness, and cramps
- do not use within 1 hour after taking an antacid or milk do not chew or crush tablet(s)
  - When using this product
  - sto mach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 Aska doctor before use if you have

Do not use if you cannot swallow without chewing. Warnings

**Drug Facts** (continued)

magnesium stearate, methacrylic acid, micro crystalline aluminum lake, hypromellose, lactose anhydrous, carmine, colloidal silicon dioxide, com starch, FD&C red #40 thactive ingredients

> ■ see end flap for expiration date and lot number 15°-30°C (59°-86°F) ■ avoid excessive humidity ■ store at 25°C (77°F); excursions permit ted between

OPENED OR BLISTER IS TORN OR BROKEN ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS Other information

children under 6 years	sak a doctor
children 6 to under 12 years	take 1 tablet in a single daily dose
adults and children 12 years and over	take 1 to 3 tablets in a single daily dose

take with a glass of water Directions 

a do not take more than directed

Drug Facts (continued)

B-1910-676A-01-GI REV0923B67601

œ

┚ П

Ш

Bisacodyl USP, 5 mg ......Stimulant laxative,

(in each tablet) Active ingredient

Drug Facts

COMPLETE PRODUCT INFORMATION KEEP OUTER PACKAGE FOR

Purpose

12 hours ■ this product generally produces bowel movement in 6 to ■ for relief of occasional constipation (irregularity) səsn

**Drug Facts** (confinued)

NDC 76162-676-01

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING No Print / No Varnish Lot no. & Exp. date

# +TopCai

# Gentle Laxative

BISACODYL USP 5 mg • STIMULANT LAXATIVE

Gentle, Dependable, Relief from Constipation





actual size

# ENTERIC COATED

information or call 1-888-423-0139 Scan here for more



www.lopcarebrand.com topcare@topco.com QUESTIONS? 1-888-423-0139 ©TOPCO LNKA1223 ITASCA, IL 60143 TOPCO ASSOCIATES LLC **DISTRIBUTED BY** REV0923867601 **##809** 

Questions or comments? 1-888-423-0139 stearic acid, talc, titanium dioxide, triacetin, triethyl citrate simethicone, sodium bicarbonate, sodium lauryl sulfate,

cellulose, polydextrose, polyethylene glycol, shellac wax,

Drug Facts (continued)

### LAXATIVE

bisacodyl tablet, delayed release

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:76162-676

**Route of Administration** ORAL

#### **Active Ingredient/Active Moiety**

Ingredient Name Basis of Strength Strength

BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y) BISACODYL 5 mg

# **Inactive Ingredients**

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B710)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

#### **Product Characteristics**

TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	В
Contains			

# **Packaging**

Markating Start Markating Eng

#	Item Code	Package Description	Marketing Start  Date	Marketing End Date
1	NDC:76162- 676-01	2 in 1 BOX	04/20/2024	
1		30 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information				
Marketing Application Number or Monograph Marketing Start Marketing End Category Citation Date Date				
OTC Monograph Drug	505G(a)(3)	04/20/2024		

# Labeler - TOPCO ASSOCIATES LLC (006935977)

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
LNK International, Inc.		038154464	pack(76162-676)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(76162-676) , pack(76162-676)

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
LNK International, Inc.		832867894	manufacture(76162-676)

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
LNK International, Inc.		967626305	pack(76162-676)

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
LNK International, Inc.		117025878	manufacture(76162-676)

Revised: 4/2024 TOPCO ASSOCIATES LLC