

NAPROXEN SODIUM 220MG- naproxen sodium 220mg tablet, coated
Command Brands, LLC

Naproxen Sodium 220mg

Active ingredient

Naproxen sodium 220mg

(naproxen 200mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- **drink a full glass of water with each dose**
- **Adults and children 12 years and older:**
 - take 1 caplet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 caplets within the first hour
 - do not exceed 2 caplets in any 8 to 12 hour period
 - do not exceed 3 caplets in a 24 hour period
- **Children under 12 years:** ask a doctor

Other information

- **each caplet contains:** sodium 20 mg
- store at 20-25°C (68-77° F)
- avoid high humidity and excessive heat above 40° C (104° F)
- **do not use if tamper evident seal under bottle is broken or missing**

Inactive ingredients

FD&C blue #2 aluminum lake, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide

May contain: corn starch, croscarmellose sodium, magnesium stearate, purified water, silicon dioxide, sodium starch glycolate, stearic acid

Questions or Comments?

Call 1-888-952-0050

Monday through Friday 9AM-5PM EST.

Principal Display Panel

Redoxon®

Naproxen Sodium
Naproxeno Sódico

All Day Pain Relief
Todo el Día Sin Dolor

24
Caplets
Cápsulas



Redoxon®

Naproxen Sodium
Naproxeno Sódico

220 mg caplets Naproxen Sodium USP (NSAID)
220 mg cápsulas Naproxeno Sódico USP (NSAID)

Distributed by: Command Brands LLC
Houston, TX 77043



Todo el Día Sin Dolor

Longer and Stronger Analgesic Pain Relief
Más Prolongado y Fuerte Alivio Analgésico Contra el Dolor

Analgesic / Fever Reducer / Reduces Inflammation
Analgésico / Baja la Fiebre / Desinflama

2 Caplets for all day long
2 Cápsulas para todo el día

24
Caplets
Cápsulas

Redoxon®

220 mg caplets Naproxen Sodium USP (NSAID)
220 mg cápsulas Naproxeno Sódico USP (NSAID)

Analgesic / Fever Reducer / Reduces Inflammation
Analgésico / Baja la Fiebre / Desinflama



MADE IN INDIA
HECHO EN LA INDIA



Lot / lote:
Date / Fecha:

Drug Facts

Active Ingredient

(In each caplet)
Naproxen sodium 220mg
Naproxen 200mg (NSAID)*
nonsteroidal anti-inflammatory drug

Purposes

Pain relief/
fever reducer/
muscular aches
and pains
due to:
■ minor pain of arthritis
■ menstrual cramps
■ backache
■ the common cold
■ temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives
■ facial swelling
■ asthma (wheezing)
■ shock
■ skin reddening
■ rash
■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed.

Heart attack and stroke warning:


NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have asthma
Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking any other drug
When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke ■ chest pain ■ trouble breathing
■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ drink a full glass of water with each dose ■ adults and children 12 years and older:
■ take 1 caplet every 8 to 12 hours while symptoms last ■ for the first dose you may take 2 caplets within the first hour ■ do not exceed 2 caplets in any 8 to 12 hour period ■ do not exceed 3 caplets in a 24 hour period
Children under 12 years: ■ ask a doctor
Other information ■ each caplet contains sodium 20 mg ■ store at 20° to 25°C (68° to 77° F) ■ avoid high humidity and excessive heat above 40°C (104°F) ■ do not use if tamper evident seal under bottle is broken or missing
Inactive ingredients F.D.C. Blue #2 aluminum lake, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide
May contain: corn starch, croscarmellose sodium, magnesium stearate, purified water, silicon dioxide, sodium starch glycolate, stearic acid
Questions or Comments?
Call 1-888-952-0050
Monday through Friday 9AM-5PM EST



Distributed by:
Command Brands LLC
Houston, TX 77043
LB1514
R0518



Naproxen Sodium 220mg

NAPROXEN SODIUM 220MG

naproxen sodium 220mg tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69 197-394
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	BLUE	Score	no score
Shape	OVAL	Size	10mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69 197-394-24	1 in 1 CARTON	12/23/2014	

1	24 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090545	12/23/2014	

Labeler - Command Brands, LLC (060684113)

Revised: 12/2018

Command Brands, LLC