

AUREOMYCIN- chlortetracycline hydrochloride granule
Phibro Animal Health

Aureomycin® 100 Granular

Aureomycin® 100 Granular

Chlortetracycline
Type A Medicated Article

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Net Wt 50 LB (22.68 kg)

Active drug ingredient

Chlortetracycline calcium complex equivalent to 100 g chlortetracycline hydrochloride per lb.

Ingredients

Dried *Streptomyces aureofaciens* Fermentation Product and Calcium Sulfate.

For use in the manufacture of medicated animal feeds.
For use in dry feed only. Not for use in liquid medicated feeds.

Use directions

Mix sufficient Aureomycin 100 Granular Medicated Article to supply desired concentration of chlortetracycline per ton with part of the feed ingredients to make a preblend. Add the remainder of the ingredients and mix thoroughly. For specific use levels, see **Indications for use**.

Mixing directions

Level desired grams per ton	Amount of medicated article per ton*
50	1/2 lb
100	1 lb
200	2 lb
400	4 lb
500	5 lb

† It is recommended that 1 pound of Aureomycin 100 Granular Type A Medicated Article be diluted with 3 pounds of one of the feed ingredients to form a 4 pound working premix. Use 2 pounds of the working premix to make a preblend (see **Use directions**) for a Type C feed containing 50 g Aureomycin chlortetracycline / ton of feed.

Indications for use

Indications for use	Chlortetracycline mg per lb body wt per day
<p>Cattle</p> <p>Beef Cattle (over 700 lb): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline.</p>	0.5
<p>Beef and Non-Lactating Dairy Cattle: As an aid in control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline when delivered in a free-choice feed. Free-choice feed must be manufactured under a medicated feed mill license utilizing an FDA approved formulation.</p>	0.5-2.0
<p>Calves, Beef and Non-Lactating Dairy Cattle: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to chlortetracycline. Feed for not more than 5 days. The appropriate amount of Aureomycin-containing feed supplement may be mixed in the cattle's daily ration or administered as a top-dress. If the Aureomycin-containing feed supplement is administered as a top-dress, it must be spread uniformly on top of the ration and sufficient space must be provided so that all cattle can eat at the same time.</p>	10
<p>Swine</p> <p>Control of porcine proliferative enteropathies (ileitis) caused by <i>Lawsonia intracellularis</i> susceptible to chlortetracycline. Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Note: this drug level is equivalent to approximately 400 grams per ton, depending on feed consumption and body weight.) Feed for not more than 14 days.</p>	10
<p>Turkeys</p> <p>Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis; coronaviral enteritis) susceptible to chlortetracycline. Feed continuously for 7 to 14 days.</p>	25

Indications for use	mg per head per day
Cattle	
Growing Cattle (over 400 lb): For the reduction of the incidence of liver abscesses.	70
Beef Cattle and Dairy Replacement Heifers: Control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella</i> spp. susceptible to chlortetracycline.	350
Beef Cattle (under 700 lb): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline.	350
Sheep	
Breeding Sheep: Reduction in the incidence of (vibriotic) abortions caused by <i>Campylobacter fetus</i> infection susceptible to chlortetracycline.	80
Indications for use	In complete feed Chlortetracycline g per ton
Swine	
Reduction in the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E <i>Streptococci</i> susceptible to chlortetracycline.	50-100
Breeding Swine: Control of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline. Feed continuously for not more than 14 days.	400
Ducks	
Control and treatment of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. Feed in complete ration to provide from 8 to 28 mg per pound of body weight per day depending upon age and severity of disease. Feed for not more than 21 days.	200-400
Chickens	
Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days.	100-200

Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days.	200-400
Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline. Feed for 5 days.	500
Turkeys Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days.	200
Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days.	400
Turkey Poults not over 4 weeks of age: Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline.	400
Indications for use Psittacine birds Warning: Psittacosis, avian chlamydiosis, or ornithosis is a reportable communicable disease, transmissible between wild and domestic birds, other animals and man. Contact appropriate public health and regulatory officials. Caution: Aspergilliosis may occur following prolonged treatment.	mg per g feed
Treatment of psittacine birds (parrots, macaws, cockatoos) suspected or known to be infected with psittacosis caused by <i>Chlamydia psittaci</i> sensitive to chlortetracycline. Feed continuously for 45 days. Each bird should consume an amount of medicated feed equal to one-fifth of its body weight daily. During treatment, parrots, macaws, and cockatoos should be kept individually or in pairs in clean cages.	10

WITHDRAWAL PERIODS AND RESIDUE WARNINGS

No withdrawal period is required when used according to labeling. This drug is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not feed to ducks or turkeys producing eggs for human consumption.

Storage

Store below 25°C (77°F), excursions permitted to 37°C (99°F)

**Restricted Drug (California) - Use only as directed. Not for use in humans.
Keep out of reach of children.**

Approved by FDA under NADA # 048-761

Questions/Comments? For technical assistance or to obtain a Safety Data Sheet(s), contact Phibro Animal Health Corporation at 1-888-403-0074 or ww.pahc.com. To report side effects, contact Phibro Animal Health Corporation at 1-888-475-7355. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Phibro, Phibro logo, Aureomycin and the A logo are trademarks owned by or licensed to Phibro Animal Health Corporation or its affiliates.

Distributed by:

Phibro Animal Health Corporation

Teaneck, NJ 07666

176-9003-02

TAKE TIME



**OBSERVE LABEL
DIRECTIONS**

Take Time Statement

PRINCIPAL DISPLAY PANEL - 50 LB Bag



Z Reference No.	TB Master No.	Description	Color
9328611	100622	BAG AUREOMYCIN 100 GRANULAR	Black
714-08143	341	24" x 10" x 5"	Black
341	341	Additional Information	Black
102024	31425		Black

Aureomycin 100 Label

PRINCIPAL DISPLAY PANEL - 2000 LB Supersack

Aureomycin® 100 Granular

Chlortetracycline Type A Medicated Article

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Active Drug Ingredient		Chlortetracycline calcium complex equivalent to 100 g chlortetracycline hydrochloride per lb.													
Ingredients		Dried <i>Streptomyces aureofaciens</i> Fermentation Product and Calcium Sulfate.													
For use in the manufacture of medicated animal feeds. For use in dry feed only. Not for use in liquid medicated feeds.															
Use directions Mix sufficient Aureomycin 100 Granular Medicated Article to supply desired concentration of chlortetracycline per ton with part of the feed ingredients to make a preblend. Add the remainder of the ingredients and mix thoroughly. For specific use levels, see Indications for Use.															
Mixing directions		<table border="1"> <thead> <tr> <th>Level desired grams per ton</th> <th>Amount of medicated article per ton†</th> </tr> </thead> <tbody> <tr> <td>50</td> <td>1/2 lb</td> </tr> <tr> <td>100</td> <td>1 lb</td> </tr> <tr> <td>200</td> <td>2 lb</td> </tr> <tr> <td>400</td> <td>4 lb</td> </tr> <tr> <td>500</td> <td>5 lb</td> </tr> </tbody> </table>	Level desired grams per ton	Amount of medicated article per ton†	50	1/2 lb	100	1 lb	200	2 lb	400	4 lb	500	5 lb	† It is recommended that 1 pound of Aureomycin 100 Granular Type A Medicated Article be diluted with 3 pounds of one of the feed ingredients to form a 4 pound working premix. Use 2 pounds of the working premix to make a preblend (see Use directions) for a Type C feed containing 50 g Aureomycin chlortetracycline / ton of feed.
Level desired grams per ton	Amount of medicated article per ton†														
50	1/2 lb														
100	1 lb														
200	2 lb														
400	4 lb														
500	5 lb														
Indications for Use	Chlortetracycline mg per lb body wt per day	Indications for Use	In complete feed Chlortetracycline g per ton												
Cattle Beef Cattle (over 700 lb): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline. Beef and Non-Lactating Dairy Cattle: As an aid in the control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline when delivered in a free-choice feed. Free-choice feed must be manufactured under a medicated feed mill license utilizing an FDA approved formulation. Calves, Beef and Non-Lactating Dairy Cattle: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to chlortetracycline. Feed for not more than 5 days. The appropriate amount of Aureomycin-containing feed supplement may be mixed in the cattle's daily ration or administered as a top-dress. If the Aureomycin-containing feed supplement is administered as a top-dress, it must be spread uniformly on top of the ration and sufficient space must be provided so that all cattle can eat at the same time.	0.5	Swine Reduction in the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E <i>Streptococci</i> susceptible to chlortetracycline. Breeding Swine: Control of leptospirosis (reducing the incidence of abortion and shedding of leptospires) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline. Feed continuously for not more than 14 days.	50-100												
	0.5-2.0		Ducks Control and treatment of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. Feed in complete ration to provide from 8 to 28 mg per pound of body weight per day depending upon age and severity of disease. Feed for not more than 21 days.	400											
	10	Chickens Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days. Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days. Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline. Feed for 5 days.	200-400												
Swine Control of porcine proliferative enteropathies (ileitis) caused by <i>Lawsonia intracellularis</i> susceptible to chlortetracycline. Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Note: this drug level is equivalent to approximately 400 grams per ton, depending on feed consumption and body weight.) Feed for not more than 14 days.	10	Turkeys Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days. Control of hexamibiasis caused by <i>Haxamita melaegris</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days. Turkey Poults not over 4 weeks of age: Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline.	200-400 500												
	25	Turkeys Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis; oronaviral enteritis) susceptible to chlortetracycline. Feed continuously for 7 to 14 days.	200 400 400												
Indications for Use	mg per head per day	Indications for Use	mg per g feed												
Cattle Growing cattle (over 400 lbs): For the reduction of the incidence of liver abscesses. Beef Cattle and Dairy Replacement Heifers: Control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella</i> spp. susceptible to chlortetracycline. Beef Cattle (under 700 lb): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline.	70 350 350	Psittacine birds Warning: Psittacosis, avian chlamydiosis, or ornithosis is a reportable communicable disease, transmissible between wild and domestic birds, other animals and man. Contact appropriate public health and regulatory officials. Caution: Aspergillosis may occur following prolonged treatment. Treatment of psittacine birds (parrots, macaws, cockatoos) suspected or known to be infected with psittacosis caused by <i>Chlamydia psittaci</i> sensitive to chlortetracycline. Feed continuously for 45 days. Each bird should consume an amount of medicated feed equal to one-fifth of its body weight daily. During treatment, parrots, macaws, and cockatoos should be kept individually or in pairs in clean cages.	10												
	Sheep Breeding Sheep: Reduction in the incidence of (vibriotic) abortions caused by <i>Campylobacter fetus</i> infection susceptible to chlortetracycline.			80											
	WITHDRAWAL PERIODS AND RESIDUE WARNINGS														
No withdrawal period is required when used according to labeling. This drug is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not feed to ducks or turkeys producing eggs for human consumption.															
Restricted Drug (California) - Use only as directed. Not for use in humans. Keep out of reach of children.		Store below 25°C (77°F), excursions permitted to 37°C (99°F) Approved by FDA under NADA # 048-761													
Questions/Comments? For technical assistance or to obtain a Safety Data Sheet(s), contact Phibro Animal Health Corporation at 1-866-403-0074 or www.pahc.com. To report side effects, contact Phibro Animal Health															

corporation at 1-800-475-1305. For additional information about reporting side effects for animal drugs, contact FDA at 1-800-FDA-1088 or <http://www.fda.gov/reportanimal>.
Phibro, Phibro logo and Aureomycin are trademarks owned by or licensed to Phibro Animal Health Corporation or its affiliates.

Distributed by:
Phibro Animal Health Corporation
Teaneck, NJ 07686



Net Wt 2000 LB (907 kg)

Lot No.: XXXXXXXXXX

Expiration Date: ????????????

176-8053-02

Z Reference No.	FG Material No.	Description	Colors
510906/00	10006836	LABEL AUREOMYCIN 100G (USA API)	BLACK Dieline
Specification / DWG	Dimensions	Country	
Z21-30213500R02	6.0" x 10.875"	US	
MPD No.	Date	Additional Information	
12542CR2	3/11/25	Label Matrix File. Wind # 1 .50" quiet zone around all 4 sides of Label	

Aureomycin 100 Label - 2000 lbs

AUREOMYCIN

chlortetracycline hydrochloride granule

Product Information

Product Type	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:66104-1003
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORTETRACYCLINE HYDROCHLORIDE (UNII: O1GX33ON8R) (CHLORTETRACYCLINE - UNII:WCK1KIQ23Q)	CHLORTETRACYCLINE HYDROCHLORIDE	100 g in 0.45 kg

Product Characteristics

Color	gray (gray/brown)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:66104-1003-1	22.68 kg in 1 BAG		
2	NDC:66104-1003-2	907 kg in 1 SUPERSACK		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
NADA	NADA048761		01/01/2009	

Labeler - Phibro Animal Health (006989008)

Revised: 3/2026

Phibro Animal Health