

**TERRAMYCIN 200 GRANULAR- oxytetracycline powder**  
**Phibro Animal Health**

-----  
**Terramycin® 200 Granular**  
**(oxytetracycline)**  
**TYPE A MEDICATED ARTICLE**

**Active Drug Ingredient:**

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to  
oxytetracycline hydrochloride . . . . . 200 g/lb

**CAUTION:**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

**CAUTION:** For use in manufacturing medicated animal feeds only.

**CAUTION:** Certain components of animal feeds, including medicated premixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety.

Precautions such as the following should be considered: dust masks or respirators and protective clothing should be worn; dust-arresting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions—seek prompt medical treatment if such reactions are suspected.

**STORE AT OR BELOW 25°C (77°F), EXCURSIONS PERMITTED UP TO 40°C (104°F)**

**FOR USE IN DRY FEEDS ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS.**

**Mixing and Use Directions**

Thoroughly mix the amount of this Type A medicated article according to the directions below with at least an equal amount by weight of feed formula ingredients prior to blending into a complete feed.

<b>Indications for Use</b>	<b>Oxytetracycline Amount</b>	<b>lb of Terramycin 200 Granular /ton</b>
<b>CHICKENS</b>		

Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> ; control of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	100-200 g/ton Feed continuously for 7-14 days	0.5-1
Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to oxytetracycline	400 g/ton Feed continuously for 7-14 days	2
Reduction of mortality due to air sacculitis (air sac infection) caused by <i>Escherichia coli</i> susceptible to oxytetracycline	500 g/ton Feed continuously for 5 days	2.5
<b>WARNING: At 500 g/ton level, withdraw 24 hours before slaughter. Zero-day withdrawal period for lower use levels. Do not use in low-calcium feed containing less than 0.55 dietary calcium. Use in such may result in violative residues. Do not administer to chickens producing eggs for human consumption.</b>		

## TURKEYS

Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to oxytetracycline	100 g/ton Feed continuously for 7-14 days	0.5
Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to oxytetracycline	200 g/ton Feed continuously for 7-14 days	1
Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline	25 mg/lb of body weight daily Feed continuously for 7-14 days	4.15 <sup>1</sup>

**WARNING: At 200 g/ton use level or higher, withdraw 5 days before slaughter. Zero-day withdrawal period for lower use levels. Do not administer to turkeys producing eggs for human consumption.**

## SWINE

Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and	10 mg/lb of body weight daily Feed continuously for 7-14	2.5 <sup>2</sup>
--	---	------------------

<i>Salmonella choleraesuis</i> susceptible to oxytetracycline and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	days	
For breeding swine for control and treatment of Leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for not more than 14 days	2.5 <sup>2</sup>

**CALVES, BEEF CATTLE, AND NONLACTATING DAIRY CATTLE**

For growing cattle (over 400 lb) for reduction of liver condemnation due to liver abscesses	75 mg/head/day Feed continuously	0.375 <sup>3</sup>
Prevention and treatment of the early stages of shipping fever complex. (Feed 3-5 days before and after arrival in feedlots.)	0.5-2.0 g/head/day	2.5-10 <sup>3</sup>
Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for 7-14 days	25 <sup>4</sup>

**SHEEP**

Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for 7-14 days	6 <sup>5</sup>
--	--	----------------

**WARNING: 5-day withdrawal before slaughter.**

- <sup>1</sup>If bird weighs 10 lb, consuming 0.6 lb of complete feed per day  
<sup>2</sup>Pig consuming 4 lb of complete feed per 100 lb body weight per day  
<sup>3</sup>Include in feed supplement based on consumption of 2 lb of supplement per head per day  
<sup>4</sup>If animal weighs 500 lb, consuming 2 lb of supplement per head per day  
<sup>5</sup>If lamb weighs 60 lb, consuming 1 lb of supplement per head per day

**NOT FOR HUMAN USE**

**RESTRICTED DRUG (CALIFORNIA) - USE ONLY AS DIRECTED  
SEE BACK PANEL FOR COMPLETE MIXING DIRECTIONS,  
USE DIRECTIONS, AND WARNINGS**

**Terramycin is a registered trademark of Pfizer, Inc.,  
licensed to Phibro Animal Health, for Oxytetracycline  
PHIBRO and PHIBRO logo design are trademarks owned by  
Phibro Animal Health Corporation or its affiliates  
Phibro Animal Health Corporation, Teaneck, NJ 07666  
Net Weight 50 lb (22.7 kg)**


Approved by FDA under NADA # 095-143

8846000

102-9142-020-05

**Terramycin 200 Granular**

Terramycin 200 Granular



# TERRAMYCIN 200 GRANULAR

oxytetracycline powder

## Product Information

<b>Product Type</b>	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:66104-9701
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OXYTETRACYCLINE HYDROCHLORIDE</b> (UNII: 4U7K4N52ZM) (OXYTETRACYCLINE ANHYDROUS - UNII:SLF0D9077S)	OXYTETRACYCLINE HYDROCHLORIDE	200 g in 0.45 kg

## Inactive Ingredients

Ingredient Name	Strength
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>RICE BRAN</b> (UNII: R60QEP13IC)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66104-9701-1	22.7 kg in 1 BAG		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA095143	01/01/2012	

**Labeler** - Phibro Animal Health (006989008)**Registrant** - Phibro Animal Health (006989008)

Revised: 11/2023

Phibro Animal Health