

CHILDRENS DELSYM COUGH PLUS COLD NIGHT TIME- acetaminophen, diphenhydramine hydrochloride, and phenylephrine hydrochloride solution
RB Health (US) LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Children's Delsym®
Cough Plus Cold Night Time

Drug Facts

Active ingredients (in each 10 mL)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Diphenhydramine HCl 12.5 mg	Antihistamine/cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sneezing
- temporarily reduces fever
- controls cough to help your child get to sleep

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- with any other product containing diphenhydramine, even one used on the skin
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if the child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not use more than directed (see Overdose warning)**
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.**Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see Overdose warning)**
- measure only with dosing cup provided

- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

Other information

- each 10 mL contains: **sodium 6 mg**
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate¹, xanthan gum

¹ may contain this ingredient

Questions?

1-866-682-4639

You may also report side effects to this phone number.

Dist. by: RB Health (US), Parsippany, NJ 07054-0224

Made in England

PRINCIPAL DISPLAY PANEL - 118 mL Bottle Carton

NDC 63824-212-64

Ages 6 Years+

**Children's
Delsym®**

COUGH + Cold

Night Time

Acetaminophen (Pain Reliever/Fever Reducer)

Diphenhydramine HCl (Antihistamine/Cough Suppressant)

Phenylephrine HCl (Nasal Decongestant)

- ☐ **Relieves Cough**
- ☐ **Relieves
Stuffy Nose**
- ☐ **Relieves
Sore Throat**
- ☐ **Reduces Fever**

Berry Flavor

4 fl oz (118 mL)

Tamper evident Do not use if neckband on bottle cap is broken or missing.

Children's Delsym[®]

COUGH+[®] Cold

Night)Time

NDC 63824-212-64 Ages 6 Years+

Children's Delsym[®]

COUGH+[®] Cold

Night)Time

Acetaminophen (Pain Reliever/Fever Reducer)
Diphenhydramine HCl (Antihistamine/Cough Suppressant)
Phenylephrine HCl (Nasal Decongestant)

- ✓ Relieves Cough
- ✓ Relieves Stuffy Nose
- ✓ Relieves Sore Throat
- ✓ Reduces Fever



Also look for our Children's Delsym 12 Hour Cough Relief In Orange and Grape Flavors




Please visit our website www.delsym.com



DOSING CUP INCLUDED



Dist. by: RB Health (US), Parsippany, NJ 07054-0224
Made in England ©2019 RB Health 031 819 3096409



Berry Flavor
4 fl oz (118 mL)



LOT CODE:

EXP. DATE:



3096409

Drug Facts

**Active ingredients
(in each 10 mL)**

Purposes

Acetaminophen 325 mg	Pain reliever/fever reducer
Diphenhydramine HCl 12.5 mg	Antihistamine/ cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sneezing
- temporarily reduces fever
- controls cough to help your child get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Drug Facts (continued)

When using this product

- do not use more than directed (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical



CHILDRENS DELSYM COUGH PLUS COLD NIGHT TIME

acetaminophen, diphenhydramine hydrochloride, and phenylephrine hydrochloride solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63824-212
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	325 mg in 10 mL
Diphenhydramine Hydrochloride (UNII: TC2D6JAD40) (Diphenhydramine - UNII:8GTS82S83M)	Diphenhydramine Hydrochloride	12.5 mg in 10 mL
Phenylephrine Hydrochloride (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine Hydrochloride	5 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
anhydrous citric acid (UNII: XF417D3PSL)	
edetate disodium (UNII: 7FLD91C86K)	
FD&C Blue No. 1 (UNII: H3R47K3TBD)	
FD&C Red No. 40 (UNII: WZB9127XOA)	
glycerin (UNII: PDC6A3C0OX)	
propylene glycol (UNII: 6DC9Q167V3)	
propyl gallate (UNII: 8D4SNN7V92)	
water (UNII: 059QF0K00R)	
sodium benzoate (UNII: OJ245FE5EU)	
sorbitol (UNII: 506T60A25R)	
sucralose (UNII: 96K6UQ3ZD4)	
trisodium citrate dihydrate (UNII: B22547B95K)	
xanthan gum (UNII: TTV12P4NEE)	

Product Characteristics

Color	BLUE	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63824-212-66	180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)	03/18/2013	
2	NDC:63824-212-64	1 in 1 CARTON	06/09/2017	
2		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	03/18/2013	

