

**ASPERCREME WITH LIDOCAINE DRY- lidocaine spray**  
**Chattem, Inc.**

-----  
**Aspercreme with Lidocaine Dry Spray**

***ASPERCREME with LIDOCAINE SPRAY***

***Drug Facts***

***Active ingredient***

Lidocaine 4%

***Purpose***

Topical anesthetic

***Use***

temporarily relieves minor pain

***Warnings***

**For external use only**

**Do not use**

- on large areas of the body or on cut, irritated or swollen skin
- on puncture wounds
- for more than one week without consulting a doctor

**When using this product**

- use only as directed. Read and follow all directions and warnings on this label.
- do not allow contact with the eyes and mucous membranes
- avoid spraying on face
- avoid inhalation of spray
- do not bandage or apply local heat (such as heating pads) or a medicated patch to area of use
- do not use at the same time as other topical analgesics

**Stop use and ask a doctor if**

- condition worsens

- redness is present
- irritation develops
- symptoms persist for more than 7 days or clear up and occur again within a few days

### **Extremely Flammable**

- do not use near heat or flame or while smoking
- avoid long term storage above 104°F (40°C)
- do not puncture or incinerate. Contents under pressure.
- do not store at temperatures above 120°F (49°C)

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

### ***Directions***

#### **adults and children over 12 years:**

- spray affected area every 6 to 8 hours, not to exceed 3 applications in a 24 hour period
- product will dry quickly on its own, and does not need to be rubbed in

**children 12 years or younger:** ask a doctor

### ***Inactive ingredients:***

alcohol denat. (30%), caprylic/capric triglyceride, isobutane, propylene glycol

### **PRINCIPAL DISPLAY PANEL**

MAX STRENGTH  
Aspercreme  
LIDOCAINE  
PAIN RELIEF SPRAY  
NET WT 4 OZ (113 g)

**MAX STRENGTH**  
**Aspercreme**<sup>®</sup>  
WITH 4% **LIDOCAINE**  
PAIN RELIEF SPRAY

**NUMBS AWAY PAIN**

- FAST ACTING
- SPRAYS AT ANY ANGLE
- QUICK-DRY FORMULA

**FRAGRANCE-FREE**  
\*Among OTC topical analgesics.  
**NET WT 4 OZ (113 g)**

**#1 LIDOCAINE BRAND\***

**DRY SPRAY**

Child-resistant packaging. Replace cap after each use.

<b>Drug Facts</b>	
<b>Active ingredient</b> Lidocaine 4%	<b>Purpose</b> Topical anesthetic
<b>Use</b> temporarily relieves minor pain	
<b>Warnings</b> For external use only	
<b>Do not use</b> <ul style="list-style-type: none"> <li>■ on large areas of the body or on cut, irritated or swollen skin</li> <li>■ on puncture wounds</li> <li>■ for more than one week without consulting a doctor</li> </ul>	
<b>When using this product</b> <ul style="list-style-type: none"> <li>■ use only as directed. Read and follow all directions and warnings on this label.</li> <li>■ do not allow contact with the eyes and mucous membranes</li> <li>■ avoid spraying on face</li> <li>■ avoid inhalation of spray</li> <li>■ do not bandage or apply local heat (such as heating pads) or a medicated patch to area of use</li> <li>■ do not use at the same time as other topical analgesics</li> </ul>	
<b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>■ condition worsens</li> <li>■ redness is present</li> <li>■ irritation develops</li> <li>■ symptoms persist for more than 7 days or clear up and occur again within a few days</li> </ul>	
<b>Extremely Flammable</b> <ul style="list-style-type: none"> <li>■ do not use near heat or flame or while smoking</li> <li>■ avoid long term storage above 104°F (40°C)</li> <li>■ do not puncture or incinerate. Contents under pressure.</li> <li>■ do not store at temperatures above 120°F (49°C)</li> </ul>	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> adults and children over 12 years: <ul style="list-style-type: none"> <li>■ spray affected area every 6 to 8 hours, not to exceed 3 applications in a 24 hour period</li> <li>■ product will dry quickly on its own, and does not need to be rubbed in</li> </ul> children 12 years or younger: ask a doctor	
<b>Inactive ingredients</b> alcohol denat. (30%), caprylic/capric triglyceride, isobutane, propylene glycol	

Dist. by Chatter, Inc., a Sanofi Company  
P.O. Box 2219, Chattanooga, TN 37409-0219  
U.S.A. ©2020  
DOT 2Q M5706 www.aspercreme.com 357-040  
0030398-01

0 41167 05835 0

## ASPERCREME WITH LIDOCAINE DRY

lidocaine spray

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41167-0583
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
ISOBUTANE (UNII: BXR49TP611)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

### Packaging

Marketing Start      Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41167-0583-5	113 mL in 1 CAN; Type 0: Not a Combination Product	01/02/2019	
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M017	01/02/2019		

**Labeler** - Chattem, Inc. (003336013)

Revised: 1/2026

Chattem, Inc.