

ANESTENKA- benzocaine liquid
Menper Distributors, Inc

Anestenka

active Ingredients(%)

Benzocaine 10%

Purpose

Oral Pain Reliever

Uses

For temporary relief of pain due to toothaches and sore gums.

methemoglobinemia warning:

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert:

do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetic. If skin reaction occurs, stop use and seek medical help right away.

Sore throat warning:

if sore throat is severe, persists for more than two days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.

Do not use:

- more than directed
- for teething

- for more than 7 days unless told to do so by a dentist or doctor
- on child under 12 years of age unless told to do so by a dentist or doctor

When using this product:

- Avoid contact with eyes
- do not exceed recommended dosage
- fever and nasal congestion are not symptoms of toothache or sore gums and may indicate the presence of infection. If these symptoms do not go away, advise your dentist or doctor.

Stop use and ask dentist or doctor if :

- Sore mouth symptoms do not get better in 7 days
- swelling, rash or fever develops
- irritation, pain, or redness do not go away
- Redness or swelling is presented

If pregnant or breast-feeding

ask a health professional before use.

Keep out of reach of children,

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

(For adults and children 12 years or older)

- wash hands
- Remove cap and cut off tip dispenser
- use your finger or cotton applicator (swab) to apply a small (pea-size) amount of Anestanka to the affected area
- Apply Anestanka to the affected area up to 4 times daily or as directed by the dentist or doctor
- For children 12 years of age: Consult a dentist or doctor

Other information:

Store at room temperature: 15-30° C (59-86° F)

Tamper Evident Feature:

Do not use if tip is cut prior to opening.

Do not use for toothache as more than a temporary expedient until a dentist can be contacted.

Inactive Ingredients:

methylparaben, propylene glycol, propylparaben

Questions or comments?

call 1-800-560-5223 weekdays 9am-4pm EST/www.menperdistributors.com



Drug Facts	
Active ingredient(s): Benzocaine 10%	Purpose: Oral Pain Reliever
Uses For the temporary relief of pain due to toothaches and sore gums.	
Warnings: Keep the product box for all Warnings and Directions. Warnings: Anesthetics warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used the product before. Stop use and seek immediate medical attention if you or a child in your care experience: ■ pale, gray, or blue colored skin (especially ■ lips/tongue) ■ headache ■ dizziness ■ weakness or loss of consciousness ■ rapid or irregular heartbeat ■ loss of consciousness ■ loss of energy.	
Other warnings: Do not use if pregnant or if you have a history of allergy to local anesthetics such as procaine, butambaine, tetracaine, or other "caine" anesthetics. If a skin reaction occurs, stop use and seek medical help right away.	
See throat warning: If sore throat is severe, persists for more than 7 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore throat symptoms do not improve in 7 days or if it worsens, persists or worsens, see your dentist or doctor promptly.	
Do not use: ■ more than directed ■ for toothache ■ For more than 7 days unless told to do so by a dentist or doctor. ■ In children under 12 years of age unless told to do so by a dentist or doctor.	
When using this product: ■ avoid contact with eyes ■ Do not exceed recommended dosage ■ Fever and nasal congestion are not symptoms of toothaches or sore gums and may indicate the presence of infection. If these symptoms do not go away, consult your dentist or doctor.	
Stop use and ask a dentist or doctor if: ■ Sore mouth symptoms do not get better in 7 days ■ Swelling, rash or other develops ■ irritation, pain or redness do not go away ■ Redness or swelling is present	
If pregnant or breastfeeding, use a health professional before use.	
Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions: For Adults and children 12 years or older: ■ Wash hands ■ Remove cap and cut off tip dispenser ■ Use your finger or rubber applicator (provided) to apply a small (peas-sized) amount of Anesthenka to the affected area. ■ Apply Anesthenka to the affected area up to 4 times daily or as directed by the dentist or doctor. ■ For children under 12 years of age: Consult a dentist or doctor.	
Other information: ■ Store at room temperature (15-30° C/59-86°F) ■ Keep out of children's reach. Do not use for a toothache as more than a temporary expedient until a dentist can be contacted. Do not use for a toothache as more than a temporary expedient until a dentist can be contacted.	
Inactive ingredients: methylparaben, propylene glycol, xylitol, sorbitol.	

ANESTENKA[®]
Benzocaine 10%

Oral Pain Reliever
due to toothaches
and sore gums



ANESTENKA[®]
Benzocaine 10%

Provides Fast Pain Relief

Oral Pain Reliever



ANESTENKA[®]
Benzocaine 10%

Provides Fast Pain Relief



ANESTENKA[®] 

Made in U.S.A.

Net contents: 12 tubes, each 0.07 Fl Oz (2 mL)

Due to toothaches



ANESTENKA

benzocaine liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53145-301
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength

BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)		BENZOCAINE	0.2 mg in 2 mL	
Inactive Ingredients				
Ingredient Name			Strength	
METHYLPARABEN (UNII: A2I8C7HI9T)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53145-301-12	12 in 1 CARTON	01/01/2014	
1		1 in 1 CARTON		
1		2 mL in 1 TUBE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M022	01/01/2014		

Labeler - Menper Distributors, Inc (101947166)

Establishment

Name	Address	ID/FEI	Business Operations
RNV LLC		118917568	manufacture(53145-301)

Revised: 7/2025

Menper Distributors, Inc