TREATONIC TATTOO NUMBING CREAM- tattoo numbing cream cream Shenzhen Situya Trading Co., Ltd.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredient

LIDOCAINE 5%

Purpose

Local Anesthetic

Use

For temporary relief local discomfort, itching, pain, soreness, or burning in the perianal area

associated with anorectal disorders and anorectal inflammation.

Do not use

Pregnant or breastfeeding, if there is any allergic reaction to this product. In case of accidental overdose, get medical help, or contact the poison control center immediately.

When Using

Only take the recommended daily dosage unless otherwise instructed by a doctor. Some individuals may experience allergic reactions to the ingredients contained in this product.

Never put it into the rectum by using the finger or any medical devices.

Stop Use

if discomfort persists for 7 days.

Ask Doctor

if discomfort persists for 7 days.

Keep Oot Of Reach Of Children

and avoid contact with eyes.

In case of accidental ingestion, seek medical attention immediately

Directions

For adults: Cleanse, Rinse, and dry the affected area with warm water. Apply a thick numbing

cream onto the treated area. Apply up to 6 times a day.

For children under 12 years of age: Consult a doctor for guidance.

Other information

Store at room temperature and out of direct sunlight or heat.

Inactive ingredients

Carbomer 934, Emu Oil, Arnica, Lecithin(Soybean), Propylene Glycol, Tocopheryl Acetate, Benzyl Alcohol, Water.

Questions

service@treatonic.com

Warnings

For external use only.

PRINCIPAL DISPLAY PANEL

Recommended State for Proparing States Tationing.

- Cleaser, Risk, and only in the last or any will warm state.

- Apply a trick number grown once in a state date, executing the clean covers the lattice.

- Apply a trick number grown once in a state date, executing the clean covers the lattice.

- When the trick has a splingly will place for any of marine for 60 minutes.

- Bronch to place cleage and expe away any excess orders. Wast for an additional to

- Blags the procedure.

- Blags the procedure.

- Concrete

- Concret

- Concrete

- Concrete

- Concrete

- Concrete

- Concrete

-



Name of the property of the control of the control

178*26



60*60*47mm

150*22mm





Recommended Steps for Preserving Before Transcript.

*Chances, Eners, and by the retained in sent wherein water.

*Apply a historium/bing cream not on the bested since, resulting the cream covers the tabloo area and all in remediate sourcomplete,

*Viriley the bested area before ship flastic ways and remarks for 6 minutes.

*Viriley the bested area before ship flastic ways and remarks for 6 minutes.

*Viriley the bested area before ship flastic ways and remarks for 6 minutes.

*Viriley the production.

*Regis The procedure.

Directions:
For adults: # Cleanse, Rinse, and dry the affected area with warm water. # Apply a th numbing cream onto the treated area. # Apply up to 8 times a day.



Notice:

• Please be aware that the duration of the numbing effect may vary depend on flectors such as temperature and briddeal akin types.

• Additionally, it is generally observed that longer wrapping times lend to produce before results in terms of numbries.

Customer Care:

For any questions, please contact us for a quick and satisfactory solution :

Distributed By: 100 KARMA INC







TREATONIC TATTOO NUMBING CREAM

tattoo numbing cream cream

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:82739-015

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength
	LIDOCAINE (LINII: 98PI200987) (LIDOCAINE - LINII:98PI200987)	LIDOCAINE	5 g in 100 ml

Inactive Ingredients		
Ingredient Name	Strength	
CARBOMER 934 (UNII: Z135WT9208)	0.4 g in 100 mL	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	6 g in 100 mL
.ALPHATOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	1 g in 100 mL
WATER (UNII: 059QF0KO0R)	86.8 g in 100 mL
BENZYL ALCOHOL (UNII: LKG8494WBH)	0.5 g in 100 mL
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	0.3 g in 100 mL

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82739-015- 01	50 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/04/2023	
2	NDC:82739-015- 02	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/04/2023	
3	NDC:82739-015- 03	60 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/04/2023	
4	NDC:82739-015- 04	100 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/04/2023	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	M016	07/04/2023	

Labeler - Shenzhen Situya Trading Co., Ltd. (706154255)

Establishment				
Name	Address	ID/FEI	Business Operations	
Shenzhen Situya Trading Co., Ltd.		706154255	manufacture(82739-015) , label(82739-015)	

Revised: 7/2023 Shenzhen Situya Trading Co., Ltd.