

IBUPROFEN- ibuprofen tablet, coated
Family Dollar (FAMILY WELLNESS)

Drug Facts

Active ingredient (in each tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and older:
 - take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, hypromellose, iron oxide red, lactose monohydrate, povidone (K-30), sodium starch glycolate, stearic acid, titanium dioxide, triacetin

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient of Advil® Tablets†

Ibuprofen

Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

Tablets

DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil® Tablets.

DISTRIBUTED BY:

MIDWOOD BRANDS LLC

500 VOLVO PARKWAY

CHESAPEAKE, VA 23320

Product Label

Lot No.:
Exp. Date:

P1D-K207A FC002223 Product of India



Drug Facts (continued)
Directions
 ■ do not take more than directed
 ■ the smallest effective dose should be used
 ■ adults and children 12 years and older:
 ■ take 1 tablet every 4 to 6 hours while symptoms persist
 ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used
 ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor
 ■ children under 12 years: ask a doctor
Other information
 ■ store between 20°-25°C (68°-77°F)
Inactive ingredients colloidal silicon dioxide, corn starch, hypromellose, iron oxide red, lactose monohydrate, povidone (K-30), sodium starch glycolate, stearic acid, titanium dioxide, triacetin
Questions or comments?
 Call 1-877-753-8955 Monday-Friday 9AM-5PM EST
DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Drug Facts (continued)
Ask a doctor before use if
 ■ stomach bleeding warning applies to you
 ■ you have problems or serious side effects from taking pain relievers or fever reducers
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 ■ you have asthma ■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are
 ■ under a doctor's care for any serious condition
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
 ■ taking any other drug
When using this product
 ■ take with food or milk if stomach upset occurs
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
Stop use and ask a doctor if
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint
 ■ have bloody or black stools ■ vomit blood
 ■ have stomach pain that does not get better
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

50844 ORG091029712
 MATHEWS, NC 28105 USA
 10401 MONROE RD
 FAMILY DOLLAR SERVICES, INC.
 DISTRIBUTED BY:
 VISIT FAMILYDOLLAR.COM
 For a refund (with receipt) or exchange,
 product within 30 days to any Family Dollar store
 Not 100% satisfied? Return package and unused
 product within 30 days to any Family Dollar store
 trademark Advil® tablets.
 Pfizer Consumer Healthcare, owner of the registered
 trademark Advil® tablets.
 This product is not manufactured or distributed by
 away. (1-800-222-1222)

Drug Facts
Active ingredient (in each tablet) Purpose
 ibuprofen USP Pain reliever/
 200 mg (NSAID)* Fever reducer
 *nonsteroidal anti-inflammatory drug
Uses
 ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches
 ■ minor pain of arthritis ■ toothache
 ■ backache ■ the common cold
 ■ menstrual cramps
 ■ temporarily reduces fever
Warnings
 Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
 Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing)
 ■ shock ■ skin redness ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed
Do not use
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
 ■ right before or after heart surgery

NDC 55319-249-01

FAMILY wellness™

COMPARE TO THE ACTIVE INGREDIENT OF ADVIL® TABLETS!

Ibuprofen
Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)

100 COATED TABLETS

100% SATISFACTION OR YOUR MONEY BACK GUARANTEED

ACTUAL SIZE



FAMILY WELLNESS Ibuprofen 200 mg Tablets

IBUPROFEN ibuprofen tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-574
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	brown	Score	no score
Shape	ROUND (Round shaped)	Size	10mm
Flavor		Imprint Code	IBU200
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-574-50	1 in 1 BOX	08/31/2023	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55319-574-10	1 in 1 BOX	08/31/2023	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079129	08/31/2023	

Labeler - Family Dollar (FAMILY WELLNESS) (024472631)