OBSTETRIX DHA COMBO PACK- docosahexaenoic acid, beta carotene, ascorbic acid, cholecalciferol, dl-alpha tocopheryl acetate, thiamine hydrochloride, riboflavin, niacinamide, pyridoxine hydrochloride, l-5methyltetrahydrofolate calcium, cyanocobalamin, carbonyl iron, magnesium oxide, zinc oxide, sodium selenate Seyer Pharmatec, Inc.

\_\_\_\_\_

Obstetrix<sup>™</sup> DHA Combo Pack

#### *Obstetrix*<sup>™</sup> *DHA* Softgels

Supplement Facts Serving Size: 1 Softgel / Servings Per Container: 30		
	Amount	% Daily
	Per Value Serving	
DHA (docosahexaenoic acid)	350 mg	*
* % Daily Value not established.		

#### **OTHER INGREDIENTS:**

Gelatin, Glycerine, Purified Water, Sodium Alginate, Stearic Acid. This product contains purified fish oil.

#### **DESCRIPTION:**

**OBSTETRIX**<sup>™</sup> **DHA** is a multivitamin/multi-mineral dietary supplement with Omega-3 Fatty Acids indicated for improving the nutritional status of women throughout pregnancy and in the postnatal period for both lactating and non-lactating mothers.<sup>†</sup>

It is recommended that for all women folate levels should be high for at least one month prior to possible conception and continued at that level for the first three months of pregnancy.<sup>†</sup>

<sup>†</sup>These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

## **DOSAGE AND ADMINISTRATION:**

One softgel daily; before, during and after pregnancy, with or without food, or as directed by a physician.

## **PRECAUTIONS:**

Ingestion of more than 3 grams per day of omega-3 fatty acids, such as DHA

(docosahexaenoic acid) has been shown to have potential anti-thrombotic effects, including increased bleeding times. Administration of **OBSTETRIX™ DHA** Omega-3 Fatty Acids should be avoided in patients taking anticoagulants.

Sold only in combination pack with **OBSTETRIX™ DHA** prenatal caplets.

# STORAGE:

Optimum storage condition= 30 to 40% Relative Humidity & 59 to 86° F (15-30°C). Protect from extreme temperatures. Freezing & excessive heat may damage softgels.

## Keep out of the reach of Children.

To report a serious adverse event or to obtain product information, contact (888) 444-5072.

Manufactured in the USA for: Seyer Pharmatec, Inc. Caguas, Puerto Rico 00725

# DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Dietary Supplement

• Enteric coated to help prevent burp back

## Obstetrix<sup>™</sup> DHA Caplets

Supplement Facts		
Serving Size: 1 Caplet /		
Servings Per Container: 30		
	Amount	% Daily
	Per,	Value**
Vitamin A (as Beta Carotene)	Serving 810 mcg	62%
Vitamin C (as Ascorbic Acid)	120 mg	100%
Vitamin D (as Cholecalciferol)	10 mcg	67%
Vitamin E (as dl-Alpha Tocopheryl Acetate)	8.1 mg	43%
Thiamin (as Thiamine HCI)	3 mg	214%
Riboflavin	3.4 mg	213%
Niacin (as Niacinamide)	20 mg	111%
Vitamin B6 (as Pyridoxine HCl)	40 mg	2000%
Folate (as L-5- Methyltetrahydrofolate Calcium)	1700 mcg DFE	283%
· ·	(1000 mcg Folic Acid)	
Vitamin B12 (as Cyanocobalamin)	12 mcg	429%

Iron (as Carbonyl Iron)	29 mg	107%
Magnesium (as Magnesium Oxide)	30 mg	8%
Zinc (as Zinc Oxide)	25 mg	192%
Selenium (as Sodium Selenate)	65mcg	93%
* Daily Value (DV) for		
Pregnant and Lactating		
Women.		

# **OTHER INGREDIENTS:**

Microcrystalline Cellulose, Methacrylic Acid Copolymer, Croscarmellose Sodium, Gelatin, Stearic Acid, Hypromellose, Maltodextrin, Crospovidone, Starch, Silica, Titanium Dioxide, Triacetin, Sucrose, Citric Acid, Magnesium Stearate, Sodium Lauryl Sulfate, Polyvinylpyrrolidone, Soy Polysaccharide, Mineral Oil, Riboflavin, FD&C Yellow #6 Lake, Sodium Ascorbate, Polyethylene Glycol, Trisodium Citrate, Tocopherols, Tricalcium Phosphate, Glycerol Monostearate, Triethyl Citrate, Ascorbyl Palmitate, Triglycerides, Sodium Benzoate, Polysorbate 80, Sorbic Acid and Talc. **Contains soy.** 

# **DESCRIPTION:**

**OBSTETRIX™ DHA** is a multivitamin/multi-mineral dietary supplement indicated for use in improving the nutritional status of women throughout pregnancy and in the postnatal period for both lactating and non-lactating mothers.<sup>†</sup>

It is recommended that for all women folate levels should be high for at least one month prior to possible conception and continued at that level for the first three months of pregnancy.<sup>†</sup>

<sup>†</sup>These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

# DOSAGE AND ADMINISTRATION:

One caplet daily or as directed by a physician.

# **CONTRAINDICATIONS:**

**OBSTETRIX**<sup>™</sup> **DHA** is contraindicated in patients with a known hypersensitivity to any of the components of this product. Hemochromatosis and hemosiderosis are contraindications to iron therapy.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or Poison Control Center immediately.

Folic acid alone is improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where Vitamin B-12 is deficient.

## **PRECAUTIONS:**

Folic acid in doses above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations remain progressive. Pernicious anemia should be excluded before using this product since folic acid may mask the symptoms of pernicious anemia.

# **ADVERSE REACTIONS:**

Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

Sold only in combination pack with **OBSTETRIX™ DHA** softgels.

## STORAGE:

Store at room temperature, USP.

#### Keep out of the reach of Children.

To report a serious adverse event or to obtain product information, contact (888) 444-5072.

Manufactured in the USA for: Seyer Pharmatec, Inc. Caguas, Puerto Rico 00725

# DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Dietary Supplement

#### with 100% Bioactive Folate

- Enteric coated
- Well tolerated
- Easy to swallow

## **OBSTETRIX™ DHA** Combo Pack

#### **DOSAGE AND ADMINISTRATION:**

One caplet and one softgel daily or as directed by a physician.

#### HOW SUPPLIED:

**OBSTETRIX**<sup>™</sup> **DHA** is available as a combination pack, which contains: Bottle of 30 DHA translucent soft gels and a Bottle of 30 orange, oval shaped tablets with imprinted "OBX". Available in Box of 30 x 2 Count Bottles 11026-2629-3 and as professional samples in Box of 7 x 2 Count Bottles. The listed product number is not a National Drug Code, but rather a product code assigned, formatted according to standard industry practice to meet the formatting requirements of pharmacy and health insurance computer systems.

Store at room temperature, approximately 15°-30°C (59° to 86°F), avoid excessive heat, light and moisture.

#### Manufactured in the USA for: SEYER PHARMATEC Caguas, Puerto Rico 00725

Dietary Supplement

# **The Preferred Prenatal**

The small, well-tolerated, prenatal duo, formulated to ensure compliance.<sup>†</sup>

Enteric coated to help prevent burp back, well tolerated, and easy to swallow.<sup>†</sup>

# Packaging

#### KIT COMPONENTS LABEL



#### DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Supplement Facts Serving Size: 1 Softgel / Servings Per Container: 30

> Amount % Daily Per Serving Value

DHA (docosahexaenoic acid) 350 mg

\* % Daily Value not established.

**OTHER INGREDIENTS:** Gelatin, Glycerine, Purified Water, Sodium Alginate, Stearic Acid. This product contains purified fish oil.

DESCRIPTION: OBSTETRIX<sup>™</sup> DHA is a multivitamin/multimineral dietary supplement with Omega-3 Fatty Acids indicated for improving the nutritional status of women throughout pregnancy and in the postnatal period for both lactating and non-lactating mothers.<sup>†</sup>

It is recommended that for all women folate levels should be high for at least one month prior to possible conception and continued at that level for the first three months of pregnancy.<sup>†</sup> **PRECAUTIONS:** Ingestion of more than 3 grams per day of omega-3 fatty acids, such as DHA (docosahexaenoic acid) has been shown to have potential anti-thrombotic effects, including increased bleeding times. Administration of **OBSTETRIX™ DHA** Omega-3 Fatty Acids should be avoided in patients taking anticoagulants.

Sold only in combination pack with **OBSTETRIX™ DHA** prenatal caplets.

**STORAGE:** Optimum storage condition= 30 to 40% Relative Humidity & 59 to 86 °F (15-30°C). Protect from extreme temperatures. Freezing & excessive heat may damage softgels.

Keep out of the reach of children.

To report a serious adverse event or to obtain product information, contact (888) 444-5072.

Rev 10/20

Code# SEY2629-3DHAL

Manufactured in the USA for:




<sup>†</sup>These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

**DOSAGE AND ADMINISTRATION:** One softgel daily; before, during and after pregnancy, with or without food, or as directed by a physician.

PEEL HERE FOR MORE INFORMATION

Seyer Pharmatec, Inc. Caguas, Puerto Rico 00725



DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.				
Supplement Serving Size: 1 Caplet / Servings P	nt Fac	cts		
	Amount Per Serving	% Daily Value**		
Vitamin A (as Beta Carotene)	810 mcg	62%		

**DESCRIPTION: OBSTETRIX™ DHA** is a multivitamin/multi-mineral dietary supplement indicated for use in improving the nutritional status of women throughout pregnancy and in the postnatal period for both lactating and non-lactating mothers.<sup>↑</sup>

It is recommended that for all women folate levels should be high for at least one month prior to possible conception

	_	
Vitamin D (as Cholecalciferol)	10 mcg	67%
Vitamin E (as dl-Alpha Tocopheryl Acetate)	8.1 mg	43%
Thiamin (as Thiamine HCI)	3 mg	214%
Riboflavin	3.4 mg	213%
Niacin (as Niacinamide)	20 mg	111%
Vitamin B6 (as Pyridoxine HCI)	40 mg	2000%
Folate (as L-5-Methyltetrahydro- 1700 folate Calcium) (1000 mcg F	mcg DFE olic Acid)	283%
Vitamin B12 (as Cyanocobalamin)	12 mcg	429%
Iron (as Carbonyl Iron)	29 mg	107%
Magnesium (as Magnesium Oxide)	30 mg	8%
Zinc (as Zinc Oxide)	25 mg	192%
Selenium (as Sodium Selenate)	65mcg	93%
* Daily Value (DV) for Pregnant and Lactatin	g Women	

OTHER INGREDIENTS: Microcrystalline Cellulose, Methacrylic Acid Copolymer, Croscarmellose 5t Sodium, Gelatin, Stearic Acid, Hypromellose, # Maltodextrin, Crospovidone, Starch, Silica, 동 Titanium Dioxide, Triacetin, Sucrose, Citric Acid, Date Magnesium Stearate, Sodium Lauryl Sulfate, Polyvinylpyrrolidone, Soy Polysaccharide, Mineral Oil, Riboflavin, FD&C Yellow #6 Lake, Sodium Ascorbate, Polyethylene Glycol, Trisodium Citrate, Tocopherols, Tricalcium Phosphate, Glycerol Monostearate, Triethyl Citrate, Ascorbyl Palmitate, Triglycerides, Sodium Benzoate, Polysorbate 80, Sorbic Acid and Talc. Contains sov.

#### PEEL HERE FOR MORE INFORMATION

and continued at that level for the first three months of pregnancy.

<sup>†</sup>These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

**DOSAGE AND ADMINISTRATION:** One caplet daily or as directed by a physician.

**CONTRAINDICATIONS: OBSTETRIX™ DHA** is contraindicated in patients with a known hypersensitivity to any of the components of this product. Hemochromatosis and hemosiderosis are contraindications to iron therapy.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or Poison Control Center immediately.

Folic acid alone is improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where Vitamin B-12 is deficient.

**PRECAUTIONS:** Folic acid in doses above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations remain progressive. Pernicious anemia should be excluded before using this product since folic acid may mask the symptoms of pernicious anemia.

ADVERSE REACTIONS: Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

Sold only in combination pack with OBSTETRIX™ DHA softgels.

STORAGE: Store at room temperature, USP.

Keep out of the reach of children,

To report a serious adverse event or to obtain product information, contact (888) 444-5072 Code# SEY2629-3PNL Rev. 10/20


Manufactured in the USA for:




PHARMATEC INC. Seyer Pharmatec, Inc. Caguas, Puerto Rico 00725



Obstetrix DHA With Bioactive Folate







# Obstetrix DHA With

**DESCRIPTION: OBSTETRIX<sup>™</sup> DHA** is a multivitamin/multi-mineral dietary supplement indicated for use in improving the nutritional status of women throughout pregnancy and in the postnatal period for both lactating and non-lactating mothers.<sup>†</sup>

These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

DOSAGE AND ADMINISTRATION: One caplet and one softgel daily or as directed by a physician.

**CONTRAINDICATIONS: OBSTETRIX™ DHA** is contraindicated in patients with a known hypersensitivity to any of the components of this product. Hemochromatosis and hemosiderosis are contraindications to iron therapy.

#### WARNINGS:

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or Poison Control Center immediately.

Folic acid alone is improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where Vitamin B-12 is deficient.

PRECAUTIONS: Folic acid in doses above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations remain progressive. Pernicious anemia should be excluded before using this product since folic acid may mask the symptoms of pernicious anemia.

**Bioactive Folate** 

ADVERSE REACTIONS: Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

**HOW SUPPLIED: OBSTETRIX DHA**<sup>TM</sup> is available as a combination pack, which contains: Bottle of 30 DHA translucent soft gels and a Bottle of 30 orange, oval shaped tablets with imprinted "OBX". Available in Box of 30 x 2 Count Bottles 11026-2629-3 and as professional samples in Box of 7 x 2 Count Bottles. The listed product number is not a National Drug Code, but rather a product code assigned, formatted according to standard industry practice to meet the formatting requirements of pharmacy and health insurance computer systems.

Store at room temperature, approximately 15°-30°C (59°-86°F), avoid excessive heat, light and moisture.

Keep out of the reach of children.

To report an adverse event or to obtain product information, contact 1-888-444-5072.





Enteric coated to help prevent burp back, well tolerated, and easy to swallow.<sup>+</sup>

#### **OBSTETRIX DHA COMBO PACK**

docosahexaenoic acid, beta carotene, ascorbic acid, cholecalciferol, dl-alpha tocopheryl acetate, thiamine hydrochloride, riboflavin, niacinamide, pyridoxine hydrochloride, l-5-methyltetrahydrofolate calcium, cyanocobalamin, carbonyl iron, magnesium oxide, zinc oxide, sodium selenate kit

Product Infor	mation					
Product Type	DIETARY S	SUPPLEMENT	Item Code (Source	e)	NHRIC:11026	-2629
,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,				-,		
Packaging						
# Item Code		age Description	Marketing Star	t Date	Marketing	End Date
<b>1</b> NHRIC:11026-262	29-3 1 in 1 K	IT				
Quantity of Pa	arts					
Part #	Package (	Quantity	Tota	I Product	t Quantity	
Part 1 1 BOTTLE			30		-	
Part 2 1 BOTTLE			30			
Part 1 of 2						
OBSTETRIX	DHA SO	FTGEL				
docosahexaeno		-				
Product Infor	mation					
Route of Admini		ORAL				
Route of Admini	stration	URAL				
Active Ingredi	ent/Active	Moiety				
	Ingre	edient Name		Basis of	f Strength	Strength
DOCONEXENT (UN	-	C) (DOCONEXENT - UN	NII:ZAD9OKH9JC)	DOCONEXE	INT	350 mg
-						
Inactive Ingre	dients					
		Ingredient Nan	ne		Str	ength
GELATIN, UNSPEC		586QN327L)				
GLYCERIN (UNII: PD WATER (UNII: 059Q						
SODIUM ALGINATE		lG2Z(0)				
STEARIC ACID (UN						
FISH OIL (UNII: XGF						
Dockoging						
Packaging	Deckers	Description	Markating Chart P	ata a		nd Data
# Item Code	30 in 1 BOTTL	Description	Marketing Start D	ate N	Marketing E	na Date
1						
Marketing	Intormat	lion				

Marketing Category	Category Citation Date Date		-	ting End ate	
ietary supplement			02/22/2023		
Part 2 of 2					
OBSTETRIX D		٢			
		calciferol, dl-alpha tocop	hen/lacet	ate thiamine hydro	chloride
		ydrochloride, l-5-methyl			cinonae,
		· · ·	2		
Cyd110C0Daiai11ii1, Ca	irbonyi iron, mag	gnesium oxide, zinc oxid	e, souium	Selendle lablet	
Product Informa	ition				
Route of Administr	ation ORAL				
Active Ingredien	t/Active Moie	ty			
	Ingredier	nt Name		Basis of Strength	Strength
BETA CAROTENE (UNI	: 01YAE03M7J) (BE	TA CAROTENE - UNII:01YAE03I	M7J)	BETA CAROTENE	810 ug
ASCORBIC ACID (UNII:	PQ6CK8PD0R) (AS	CORBIC ACID - UNII:PQ6CK8PE	DOR)	ASCORBIC ACID	120 mg
CHOLECALCIFEROL (UNII: 1C6V77QF41) (CHOLECALCIFEROL - UNII:1C6V77QF41)				CHOLECALCIFEROL	10 ug
<b>.ALPHATOCOPHEROL ACETATE, DL-</b> (UNII: WR1WPI7EW8) (.ALPHA TOCOPHEROL, DL UNII:7QWA1RIO01)				.ALPHATOCOPHEROL, DL-	8.1 mg
				thiamine Hydrochloride	3 mg
<b>RIBOFLAVIN</b> (UNII: TLM2976OFR) (RIBOFLAVIN - UNII:TLM2976OFR)RIBOFLAVIN				RIBOFLAVIN	3.4 mg
NIACINAMIDE (UNII: 25	X51I8RD4) (NIACINA	AMIDE - UNII:25X51I8RD4)		NIACINAMIDE	20 mg
PYRIDOXINE HYDROCHLORIDE (UNII: 68Y4CF58BV) (PYRIDOXINE - PYRIDOXINE   UNII:KV2JZ1BI6Z) HYDROCHLORIDE					40 mg
EVOMEFOLATE CALCIUM (UNII: A9R10K3F2F) (LEVOMEFOLIC ACID - LEVOMEFOLATE CALCIUM					1700 ug
CYANOCOBALAMIN (L	NII: P6YC3EG204)	CYANOCOBALAMIN - UNII:P6Y	C3EG204)	CYANOCOBALAMIN	12 ug
RON (UNII: E1UOL152H	7) (IRON - UNII:E1U	OL152H7)		IRON	29 mg
MAGNESIUM OXIDE (L	INII: 3A3U0GI71G) (	MAGNESIUM CATION - UNII:T6	V3LHY838)	MAGNESIUM OXIDE	30 mg
ZINC OXIDE (UNII: SOI	2LOH54Z) (ZINC O	(IDE - UNII:SOI2LOH54Z)		ZINC OXIDE	25 mg
SODIUM SELENATE (U	NII: 5DQP25600A)	SELENATE ION - UNII:6X37R1	OB70)	SELENIUM	65 ug
Inactive Insurable	nto				
Inactive Ingredie					Character 11
					Strength
			(UNII: 74G4	K01H13)	
GELATIN, UNSPECIFIE		27L)			
STEARIC ACID (UNII: 4					
HYPROMELLOSE, UNS		NXW29V3WO)			
MALTODEVTDIN / UNIT	7CVR7L4A2D)				
CROSPOVIDONE, UNS					

	•				
	(UNII: ETJ7Z6XBU4)				
	DE (UNII: 15FIX9V2JP)				
<b>TRIACETIN</b> (UNII:					
SUCROSE (UNII: C					
	NOHYDRATE (UNII: 2968PHW8QF	<b>'</b> )			
	<b>ARATE</b> (UNII: 70097M6I30)				
	SULFATE (UNII: 368GB5141J)				
-	JNII: SZR7Z3Q2YH)				
SOYBEAN COTYL	EDON CELL WALL POLYSACC	HARIDES (UNII: 4UL6	5DF56YQ)		
MINERAL OIL (UN	II: T5L8T28FGP)				
FD&C YELLOW N	<b>O. 6</b> (UNII: H77VEI93A8)				
SODIUM ASCORB	<b>ATE</b> (UNII: S033EH8359)				
POLYETHYLENE	GLYCOL, UNSPECIFIED (UNII: 3	WJQ0SDW1A)			
SODIUM CITRATE	, UNSPECIFIED FORM (UNII: 10	Q73Q2JULR)			
TOCOPHEROL (U	NII: R0ZB2556P8)				
FRICALCIUM PHO	SPHATE (UNII: K4C08XP666)				
GLYCERYL MONO	STEARATE (UNII: 2300U9XXE4)				
TRIETHYL CITRA	E (UNII: 8Z96QXD6UM)				
ASCORBYL PALM	ITATE (UNII: QN83US2B0N)				
MEDIUM-CHAIN 1	RIGLYCERIDES (UNII: C9H2L21)	√7U)			
SODIUM BENZOA	TE (UNII: OJ245FE5EU)				
POLYSORBATE 8	<b>0</b> (UNII: 60ZP39ZG8H)				
SORBIC ACID (UN	II: X045WJ989B)				
TALC (UNII: 7SEV7	J4R1U)				
Packaging					
# Item Code	Package Description	Marketing S	Start Date	Mark	eting End Date
1	30 in 1 BOTTLE				
1	30 in 1 BOTTLE				
Marketing	Information				
Marketing	Application Number Citation		Marketing Date		Marketing End Date
Category					

Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
dietary supplement		02/22/2023				

Labeler - Seyer Pharmatec, Inc. (832947126)