

ALTUVIIIIO- antihemophilic factor (recombinant), fc-vwf-xten fusion protein-ehtl

Bioverativ Therapeutics Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ALTUVIIIIO® safely and effectively. See full prescribing information for ALTUVIIIIO.

ALTUVIIIIO [antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-ehtl], lyophilized powder for solution, for intravenous use
Initial U.S. Approval: 2023

----- RECENT MAJOR CHANGES -----

Warnings and Precautions (5.2)

12/2025

----- INDICATIONS AND USAGE -----

ALTUVIIIIO [antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-ehtl] is a recombinant DNA-derived, Factor VIII concentrate indicated for use in adults and children with hemophilia A (congenital factor VIII deficiency) for:

- Routine prophylaxis to reduce the frequency of bleeding episodes
- On-demand treatment & control of bleeding episodes
- Perioperative management of bleeding (1)

Limitation of Use:

ALTUVIIIIO is not indicated for the treatment of von Willebrand disease. (1)

----- DOSAGE AND ADMINISTRATION -----

For intravenous use only.

- Each ALTUVIIIIO vial label states Factor VIII activity in international units (IU or unit). (2.1)
 - For routine prophylaxis: 50 IU/kg once weekly. (2.1)
 - For on-demand treatment and control of bleeding episodes and perioperative management: 50 IU/kg (2.1)
- Estimated Increment of Factor VIII (IU/dL or % of normal) = 50 IU/kg × 2 (IU/dL per IU/kg) (2.1)

To achieve a specific target Factor VIII activity level, use the following formula: Dosage (IU) = Body Weight (kg) × Desired Factor VIII Increase (IU/dL or % normal) × 0.5 (IU/kg per IU/dL). (2.1)

----- DOSAGE FORMS AND STRENGTHS -----

For injection: nominally 250, 500, 750, 1000, 2000, 3000, or 4000 IU, lyophilized powder in single-dose vials for reconstitution. (3)

----- CONTRAINDICATIONS -----

Do not use in patients who have had severe hypersensitivity reactions, including anaphylaxis, to ALTUVIIIIO or excipients of ALTUVIIIIO. (4)

----- WARNINGS AND PRECAUTIONS -----

- Hypersensitivity reactions, including anaphylaxis, have occurred with ALTUVIIIIO. If symptoms occur, immediately discontinue ALTUVIIIIO and initiate appropriate treatment. (5.1)
- Neutralizing antibodies (inhibitors) to Factor VIII have been reported. If expected plasma Factor VIII activity levels are not attained, or if bleeding is not controlled with an appropriate dose, perform an assay that measures Factor VIII inhibitor concentration. (5.2, 5.3)

----- ADVERSE REACTIONS -----

Most common adverse reactions (incidence >10%) are headache and arthralgia. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Bioverativ Therapeutics Inc. (A SANOFI COMPANY) at 1-800-633-1610 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- USE IN SPECIFIC POPULATIONS -----

Pediatric Use: No dosing adjustment is needed in this population. (8.4)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dose
- 2.2 Preparation and Reconstitution
- 2.3 Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Hypersensitivity Reactions
- 5.2 Neutralizing Antibodies
- 5.3 Monitoring Laboratory Tests

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.6 Immunogenicity

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

ALTUVIIIIO is indicated for use in adults and pediatric patients with hemophilia A (congenital factor VIII deficiency) for:

- Routine prophylaxis to reduce the frequency of bleeding episodes

- On-demand treatment and control of bleeding episodes
- Perioperative management of bleeding

Limitation of Use

ALTUVIIIIO is not indicated for the treatment of von Willebrand disease.

2 DOSAGE AND ADMINISTRATION

For intravenous use after reconstitution only.

2.1 Dose

- Each ALTUVIIIIO vial label states the Factor VIII potency in international units (IU). One IU corresponds to the Factor VIII activity contained in one milliliter of normal human plasma, as defined by the current World Health Organization (WHO) international standard for Factor VIII concentrate.
- Potency assignment for ALTUVIIIIO is determined using an activated partial thromboplastin time (aPTT)-based one-stage clotting assay. It is recommended to use a validated one-stage clotting assay to measure ALTUVIIIIO Factor VIII activity in plasma. The ALTUVIIIIO Factor VIII activity level is overestimated by the chromogenic assay and a specific ellagic acid based aPTT reagent in one-stage clotting assay by approximately 2.5-fold [*see Warnings and Precautions (5.3)*].

For the dose of 50 IU/kg, the expected *in vivo* peak increase in Factor VIII level expressed as IU/dL (or % of normal) is estimated using the following formula:

Estimated Increment of Factor VIII (IU/dL or % of normal) = 50 IU/kg × 2 (IU/dL per IU/kg)

To achieve a specific target Factor VIII activity level, use the following formula: Dosage (IU) = Body Weight (kg) × Desired Factor VIII Increase (IU/dL or % normal) × 0.5 (IU/kg per IU/dL).

Routine Prophylaxis

The recommended dosing for routine prophylaxis for adults and children is 50 IU/kg of ALTUVIIIIO administered once weekly.

On-demand Treatment and Control of Bleeding Episodes

ALTUVIIIIO dosing for the on-demand treatment and control of bleeding episodes is provided in Table 1.

Table 1: Dosing for On-demand Treatment and Control of Bleeding Episodes

Type of Bleeding	Recommended Dose	Additional Information
Minor and Moderate For example: Uncomplicated joint		For minor and moderate bleeding episodes occurring within 2 to 3 days after a prophylactic

Uncomplicated joint bleeds, minor muscular bleeds, mucosal or subcutaneous bleeds	Single dose of 50 IU/kg	dose, a lower dose of 30 IU/kg dose may be used. Additional doses of 30 or 50 IU/kg every 2 to 3 days may be considered.
Major For example: Intracranial, retroperitoneal, iliopsoas and neck bleeds, muscle bleeds with compartment syndrome and bleeds associated with a significant decrease in the hemoglobin level	Single dose of 50 IU/kg	Additional doses of 30 or 50 IU/kg every 2 to 3 days can be considered.

For resumption of prophylaxis (if applicable) after treatment of a bleed, it is recommended to allow an interval of at least 72 hours between the last 50 IU/kg dose for treatment of a bleed and resuming prophylaxis dosing. Thereafter, prophylaxis can be continued as usual on the patient's regular schedule.

Perioperative Management

ALTUVIII O dosing for perioperative management is provided in Table 2.

Table 2: Dosing for Perioperative Management

Type of Surgery	Pre-operative Dose	Post-operative Dose
Minor For example: Tooth extraction	Single dose of 50 IU/kg	An additional dose of 30 or 50 IU/kg after 2 to 3 days may be considered.
Major For example: Intracranial, intra-abdominal, joint replacement surgery, or complicated dental procedures.	Single dose of 50 IU/kg	Additional doses of 30 or 50 IU/kg every 2 to 3 days may be administered as clinically needed for perioperative management.

2.2 Preparation and Reconstitution


1. Use aseptic technique and a flat work surface during the reconstitution procedure.
2. Allow the ALTUVIII O vial, containing the white to off-white lyophilized powder, and the

prefilled diluent syringe to reach room temperature before use.


3. Remove the plastic cap from the ALTUVIIIIO vial and wipe the rubber stopper of the vial with an alcohol wipe. Allow the rubber stopper to dry.
4. Completely remove the backing from the vial adapter package by peeling back the lid. **Do not** remove the vial adapter from the package or touch the inside of the package of the adapter.




5. Keep the vial on a flat surface. Hold the vial with one hand and using the other hand, place the vial adapter in its package over the vial. The spike should be placed directly above the center of the rubber stopper. Push the vial adapter straight down until the spike on the vial adapter punctures the center of the vial stopper and is fully inserted.



6. Lift the package cover away from the vial adapter and throw away the cover.



7. Only use the prefilled diluent syringe provided to reconstitute the powdered medicine. Hold the plunger rod by the circular disk. Place the tip of the plunger rod into the end of the prefilled diluent syringe. Turn the plunger rod to the right until it is firmly attached.



8. With one hand, hold the prefilled diluent syringe directly under the cap with the cap pointing up. Make sure you are holding the prefilled diluent syringe by the ridged part directly under the cap. **Do not** use if the cap has been removed or is not securely attached.
9. With your other hand, grasp the cap and bend it at a 90 degree angle until it snaps off. After the cap snaps off, you will see the glass tip of the prefilled diluent syringe. **Do not** touch the glass tip of the prefilled diluent syringe or the inside of the cap.
10. Be sure the vial is sitting on a flat surface. Insert the tip of the prefilled diluent syringe into the vial adapter opening. Turn the prefilled diluent syringe to the right until it is securely attached to the vial adapter.
11. Slowly push down on the plunger rod to inject all of the liquid (diluent) from the prefilled diluent syringe into the vial. The plunger rod may rise slightly afterward. This is normal.
12. With the prefilled diluent syringe still connected to the adapter, gently swirl the vial until the powder is completely dissolved. Check the solution through the vial to make sure the powder is fully dissolved. The solution should look clear and colorless to opalescent. **Do not** shake. **Do not** use the reconstituted ALTUVIIIIO if it contains visible particles or is cloudy.

POOLING: pooling is the process of combining two or more reconstituted vials into a larger luer lock syringe (not provided in the carton). If the dose requires more than one vial, reconstitute each vial as described above (See Steps 3–12) with the prefilled diluent syringe provided. **Do not** detach the prefilled diluent syringe until you are ready to attach the larger luer lock syringe to the next vial. Keep the vial adapter attached to the vial as you will need it for attaching a larger luer lock syringe. Use a larger luer lock plastic syringe to combine the contents of the reconstituted vials into the syringe, similar as described in Steps 13–14. Repeat this pooling procedure with each vial you will be using. Once you have pooled the required dose, proceed with the administration steps using the larger luer lock syringe.

13. Make sure the plunger rod is pressed all the way down and the diluent syringe is firmly attached to the vial adapter. Turn the vial upside-down. Slowly pull down on the plunger rod to draw all the solution from the vial into the diluent syringe. Be careful not to pull the plunger rod completely out of the diluent syringe.
14. Gently unscrew the diluent syringe from the vial adapter by turning it to the right. Dispose of the vial with the adapter still attached. If you are not ready to inject, put the syringe cap carefully back onto the syringe tip. **Do not** touch the syringe tip or the inside of the cap.
15. Use the reconstituted ALTUVIIIIO as soon as possible, but no later than **3 hours** after reconstitution. Do not touch the glass tip of the syringe if not used immediately after reconstitution. Protect from direct sunlight. Do not refrigerate after reconstitution.

2.3 Administration

For intravenous use only.

- Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. **Do not** use ALTUVIIIIO solution if particulate matter or discoloration is observed.
- **Do not** administer reconstituted ALTUVIIIIO in the same tubing or container with other medications.

Administration Steps:

1. Attach the syringe to the connector end of the infusion set tubing by turning it to the right until it is securely attached.
2. Push the plunger rod until all air is removed from the syringe and ALTUVIIIIO has filled the infusion set needle. **Do not** push ALTUVIIIIO solution through the needle.
3. Remove the protective needle cover from the infusion set needle.
4. Perform intravenous injection slowly over 1 to 10 minutes, based on the patient's comfort level.
5. After infusing ALTUVIIIIO, remove and properly discard the infusion set.

3 DOSAGE FORMS AND STRENGTHS

ALTUVIIIIO is available as a white to off-white lyophilized powder for reconstitution in single-dose vials containing nominally 250, 500, 750, 1000, 2000, 3000, or 4000 international units (IU) per vial.

4 CONTRAINDICATIONS

ALTUVIIIIO is contraindicated in patients who have had severe hypersensitivity reactions, including anaphylaxis, to the product or its excipients [see *Description (11)*].

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions

Hypersensitivity reactions, including anaphylaxis, have occurred with ALTUVIIIIO [see *Postmarketing Experience (6.2)*]. Signs and symptoms include, but not limited to, hives, shortness of breath, chest tightness, wheezing, hypotension, nausea, vomiting, and itching. Discontinue ALTUVIIIIO if hypersensitivity reaction occurs and manage symptoms as appropriate.

5.2 Neutralizing Antibodies

Formation of neutralizing antibodies (inhibitors) to Factor VIII has been reported following administration of ALTUVIIIIO [see *Postmarketing Experience (6.2)*]. Monitor all patients for the development of Factor VIII inhibitors by appropriate clinical observations and laboratory tests. If the patient's plasma Factor VIII level fails to increase as expected or if bleeding is not controlled after ALTUVIIIIO administration, the presence of an inhibitor (neutralizing antibodies) should be suspected, and appropriate testing performed [see *Warnings and Precautions (5.3)*].

5.3 Monitoring Laboratory Tests

If assessment of plasma Factor VIII activity is needed, it is recommended to use a validated one-stage clotting assay [see *Dosage and Administration (2)*]. The ALTUVIIIIO Factor VIII activity level is overestimated by the chromogenic assay and a specific ellagic acid based aPTT reagent in one-stage clotting assay by approximately 2.5-fold. If these assays are used, divide the result by 2.5 to approximate the patient's ALTUVIIIIO Factor VIII activity level. Use of a reference laboratory is recommended when a qualified one-stage clotting assay or chromogenic assay is not available locally.

Monitor for the development of Factor VIII inhibitors. If bleeding is not controlled with ALTUVIIIIO and the expected factor VIII activity plasma levels are not attained, perform an assay to determine if Factor VIII inhibitors are present (use Bethesda Units to titer inhibitors).

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety data described in this section reflects exposure to ALTUVIIIIO in two clinical studies, Study 1 and Study 2 and are pooled for analysis. In Study 1, a total of 159 previously treated patients (PTPs) (134 adults and 25 adolescents) with severe Hemophilia A) received at least one dose of ALTUVIIIIO for either routine prophylaxis, on-demand treatment of bleeding episodes or perioperative management. A total of 152

(96%) patients achieved at least 25 exposure days and 115 (72%) patients achieved at least 50 exposure days with a median of 53.0 (range 2–63) for both exposure days and injections per patient. Overall exposure was monitored for a total of 151.5 patient-years [see *Clinical Studies (14)*].

In Study 2, the safety of ALTUVIIIIO was evaluated in 74 male PTPs <12 years of age with severe hemophilia A who received at least one dose of ALTUVIIIIO. Sixty-six (89.2%) patients achieved at least 50 exposure days with a median of 53.0 (range 3–72).

Adverse events were monitored for a total of 210.7 patient-years in 2 completed clinical studies in PTPs. Adverse drug reactions (ADRs) (summarized in Table 3) were reported in 79 (33.9%) of the 233 patients treated with routine prophylaxis or on-demand therapy. The most common ADRs (>10%) in adults and adolescents were headache (20.1%) and arthralgia (16.4%). In children below 12 years, pyrexia (12.2%) was the most common ADR (>10%). In the studies, no inhibitors to FVIII were detected and no ADRs of anaphylaxis were reported.

The most common adverse reactions (>10% of patients) reported in clinical trials were headache and arthralgia.

Table 3: Adverse Reactions with Frequency of ≥3% Reported in ALTUVIIIIO Studies*

MedDRA System Organ Class	Adverse Drug Reactions	Number of Patients n (%) (N = 233)*
Nervous system disorders	Headache	35 (15)
Musculoskeletal and connective tissue disorders	Arthralgia	31 (13)
	Pain in extremity	10 (4)
	Back pain	9 (4)
General disorders and administration	Pyrexia	10 (4)
Gastrointestinal disorders	Vomiting	7 (3)

* Pooled data from Study 1 and Study 2 including 233 patients across the adult and adolescent and pediatric studies.

Thromboembolic events occurred in 1% (3/261) of patients in the long-term safety extension study; these three patients had pre-existing risk factors.

6.2 Postmarketing Experience

The following adverse reactions have been identified during the post approval use of ALTUVIIIIO. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Blood and lymphatic system disorders: Factor VIII inhibitor development [see *Warnings and Precautions (5.2)*].

Immune system disorders: Hypersensitivity reactions, including anaphylaxis [see *Warnings and Precautions (5.1)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no data with ALTUVIIIIO use in pregnant women to inform a drug-associated risk. Animal developmental and reproductive studies have not been conducted with ALTUVIIIIO. Therefore, it is not known whether ALTUVIIIIO can affect reproductive capacity or cause fetal harm when given to pregnant women.

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2–4% and 15–20%, respectively.

8.2 Lactation

Risk Summary

There is no information regarding the presence of ALTUVIIIIO in human milk, its effects on the breastfed infant, or its effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for ALTUVIIIIO and any potential adverse effects on the breastfed infant from ALTUVIIIIO or from the underlying maternal condition.

8.4 Pediatric Use

The safety and effectiveness of ALTUVIIIIO for routine prophylaxis, on-demand treatment, and perioperative management of bleeding episodes have been established in pediatric patients <18 years old. The use of ALTUVIIIIO for these indications is supported by evidence from two clinical studies which enrolled 99 previously treated patients <18 years of age who received at least one dose of ALTUVIIIIO as part of routine prophylaxis, treatment of bleeding episodes, or perioperative management. Thirty-eight patients (38.4%) were <6 years of age, 36 (36.4%) patients were 6 to <12 years of age, and 25 patients (25.2%) were adolescents (12 to <18 years of age). Data from the pediatric study (74 patients) <12 years of age showed that no dosing adjustment was required [see *Adverse Reactions (6.1)*, *Clinical Pharmacology (12)*, and *Clinical Studies (14)*].

8.5 Geriatric Use

Clinical studies of ALTUVIIIIO did not include sufficient numbers of patients 65 years of age and older to determine whether or not such patients respond differently from younger patients. However, clinical experience with other Factor VIII products has not identified differences between the elderly and younger patients.

11 DESCRIPTION

ALTUVIIIIO [antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-ehtl] is a sterile, non-pyrogenic, white to off-white lyophilized powder for reconstitution for intravenous injection. The product is supplied in single-dose vials containing nominal

potencies of 250, 500, 750, 1000, 2000, 3000, or 4000 international units (IU). Each vial of ALTUVIIIIO is labeled with the actual Factor VIII activity content in IU. The powder for injection is reconstituted with 3 mL sterile water for injection (sWFI) supplied in a sterile prefilled syringe. The reconstituted solution should be essentially free of particles. The final product contains the excipients: arginine hydrochloride (250 mM), calcium chloride dihydrate (5 mM), histidine (10 mM), polysorbate 80 (0.05% w/v), and sucrose (5% w/v).

The active ingredient in ALTUVIIIIO is a fully recombinant fusion protein comprising a single chain B-domain deleted (BDD) analogue of human FVIII covalently fused to the Fc domain of human immunoglobulin G1 (IgG1), the FVIII-binding D'D3 domain of human von Willebrand factor (VWF), and 2 XTEN polypeptides. ALTUVIIIIO contains 2829 amino acids with an apparent molecular weight of 312 kDa. ALTUVIIIIO is synthesized as 2 polypeptide chains which are covalently linked by 2 Fc hinge disulfide bonds. The first FVIII-XTEN-Fc polypeptide chain contains the A1A2 domain of FVIII along with 5 amino acids from B-domain (1-745 amino acids) fused to the 288-XTEN polypeptide (in place of the natural FVIII B-domain), the A3C1C2 domain of FVIII (1649-2332), and the Fc domain of human IgG1. The second VWF-XTEN-a2-Fc polypeptide chain contains the D'D3 domain of VWF (1-477 amino acids) fused to the 144-XTEN polypeptide, a thrombin cleavable acidic region 2 sequence from FVIII and the Fc domain of human IgG1. The Fc domain includes the hinge, CH₂, and CH₃ domains of IgG1. The Fc, VWF, and XTEN polypeptide portions of the molecule extend the half-life of ALTUVIIIIO in plasma.

ALTUVIIIIO is produced by recombinant DNA technology in a human embryonic kidney (HEK) cell line, which has been extensively characterized. ALTUVIIIIO is manufactured without addition of human- or animal-derived components and purified by a combination of multiple chromatography steps, a detergent or solvent/detergent viral inactivation step, a nano filtration step for viral clearance, and ultrafiltration steps.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

ALTUVIIIIO temporarily replaces the missing coagulation factor VIII needed for effective hemostasis. ALTUVIIIIO has demonstrated 3- to 4-fold prolonged half-life relative to other standard and extended half-life FVIII products.

Mechanism of Half-life Extension

ALTUVIIIIO is a recombinant FVIII analogue fusion protein that is independent of endogenous VWF in order to overcome the half-life limit imposed by FVIII-VWF interactions. The D'D3 domain of VWF is the region that interacts with FVIII. Appending the D'D3 domain of VWF to a recombinant FVIII-Fc fusion protein provides protection and stability to FVIII, and prevents FVIII interaction with endogenous VWF, thus overcoming the limitation on FVIII half-life imposed by VWF clearance.

The Fc region of human immunoglobulin G1 (IgG1) binds to the neonatal Fc receptor (FcRn). FcRn is part of a naturally occurring pathway that delays lysosomal degradation of immunoglobulins by recycling them back into circulation, thus prolonging the plasma half-life of the fusion protein.

ALTUVIIIIO contains 2 XTEN polypeptides, which alter the hydrodynamic radius of the

fusion protein, thus reducing rates of clearance and degradation, and improving pharmacokinetic properties. In ALTUVIII O, the natural FVIII B domain (except 5 amino acids) is replaced with the first XTEN polypeptide, inserted in between FVIII N745 and E1649 amino acid residues; and the second XTEN polypeptide is inserted in between the D'D3 domain and Fc.

12.2 Pharmacodynamics

Hemophilia A is a bleeding disorder characterized by a deficiency of functional coagulation factor VIII (FVIII), which leads to a prolonged clotting time in the activated partial thromboplastin time (aPTT)-based one-stage clotting assay. Administration of ALTUVIII O increases plasma levels of FVIII, temporarily correcting the coagulation defect in hemophilia A patients.

Based on FVIII pharmacokinetic/pharmacodynamic analyses, the risk of bleeding is negatively correlated with FVIII activity. Once weekly 50 IU/kg ALTUVIII O provided factor VIII activity levels that were associated with a low bleed risk.

12.3 Pharmacokinetics

The PK of ALTUVIII O were evaluated in prospective, open-label clinical studies, enrolling 159 adults and adolescents, and 74 children <12 years old, respectively, receiving weekly IV injections of 50 IU/kg. Among children <12 years old, 37 patients had ALTUVIII O single dose PK profiles available.

PK parameters following a single dose of ALTUVIII O are presented in Table 4. The PK parameters were based on plasma FVIII activity measured by the aPTT-based one-stage clotting assay. After a single dose of 50 IU/kg, ALTUVIII O exhibited high sustained FVIII activity with prolonged half-life across age cohorts. There was a trend of increasing area under the curve (AUC), and decreasing clearance, with increasing age in the pediatric cohorts. The PK profile at steady state (Week 26) was comparable with the PK profile obtained after the first dose.

Table 4: Pharmacokinetic Parameters Following a Single Dose of ALTUVIII O by age (one-stage clotting assay)

PK Parameters (mean SD)	Pediatric Study 1 to <6 Years N = 18	Pediatric Study 6 to <12 Years N = 18	Adult and Adolescent Study 12 to <18 years N = 25	Adult and Adolescent Study Adults N = 134
AUC (IU×h/dL)	6800 (1120)*	7190 (1450)	8350 (1550)	9850 (2010)†
t_{1/2} (h)	38.0 (3.7)	42.4 (3.7)	44.6 (5.0)	48.2 (9.3)
CL (mL/h/kg)	0.742 (0.121)	0.681 (0.139)	0.582 (0.115)	0.493 (0.121)†
V_{SS} (mL/kg)	36.6 (5.6)	38.1 (6.8)	34.9 (7.4)	31.0 (7.3)†
MRT (hr)	49.6 (5.5)	56.3 (5.1)	60.0 (5.5)	63.9 (10.2)†

AUC_{0-tau} = area under the activity-time curve over the dosing interval, CL = clearance, MRT = mean residence time, SD = standard deviation, t_{1/2z} = terminal half-life, V_{SS} = volume of distribution at steady state.

* N = 17

† Calculation based on 128 profiles.

ALTUVIIIIO at steady state maintained normal to near normal (>40 IU/dL) FVIII activity for a mean (SD) of 4.1 (0.7) days with once weekly prophylaxis in adults. The FVIII activity over 10 IU/dL was maintained in 83.5% of adults and adolescent patients throughout the study. In children <12 years ALTUVIIIIO maintained normal to near normal (>40 IU/dL) FVIII activity for 2 to 3 days and >10 IU/dL FVIII activity for approximately 7 days (see Table 5).

Table 5: Pharmacokinetic Parameters at Steady State of ALTUVIIIIO by age (one-stage clotting assay)

PK Parameters Mean (SD)	Pediatric Study* 1 to <6 years N = 37	Pediatric Study* 6 to <12 years N = 36	Adult and Adolescent Study* 12 to <18 years N = 24	Adult and Adolescent Study* Adults N = 125
Peak (IU/dL)	136 (49) (N = 35)	131 (36) (N = 35)	124 (31)	150 (35) (N = 124)
IR (kg×IU/dL/IU)	2.22 (0.83) (N = 35)	2.10 (0.73) (N = 35)	2.25 (0.61) (N = 22)	2.64 (0.61) (N = 120)
Time to 40 IU/dL (h)	68.0 (10.5)†	80.6 (12.3)†	81.5 (12.1)‡	98.1 (20.1)‡
Time to 20 IU/dL (h)	109 (14)†	127 (15)†	130 (16)‡	150 (28)‡
Time to 10 IU/dL (h)	150 (18)†	173 (17)†	179 (20)‡	201 (36)‡
Trough (IU/dL)	10.9 (19.7) (N = 36)	16.5 (23.7)	9.23 (4.77) (N = 22)	18.0 (16.6) (N = 123)

Peak = 15 min post dose at steady state, IR = incremental recovery, Trough – predose FVIII activity value at steady state, SD = standard deviation.

* Steady state peak, trough and IR were computed using available measurements at week 52/End of study PK sampling visit.

† Time to FVIII activity was predicted using population PK model for pediatric study.

‡ Time to FVIII activity was predicted using population PK model for adult study.

Specific Populations

The following factors have no clinically meaningful effect on the pharmacokinetics of ALTUVIIIIO: age (1.4 to 72 years), sex, race (White, Asian), VWF activity (40 to 339 IU/dL), hematocrit level (28% to 57%), blood type, HCV status, or HIV status. Body weight (12.5 to 133 kg) is expected to alter weight normalized clearance (dL/h/kg) by 79% to -18% compared to a typical patient.

12.6 Immunogenicity

The observed incidence of anti-drug antibodies (ADAs) is highly dependent on the sensitivity and specificity of the assay. Differences in assay methods preclude

meaningful comparisons of the incidence of anti-drug antibodies in the studies described below with the incidence of anti-drug antibodies in other studies, or of other Factor VIII products.

In Study 1 and Study 2, all patients were monitored for neutralizing antibodies (inhibitors) to Factor VIII in the clinical program. No patients developed neutralizing antibodies to Factor VIII. Factor VIII inhibitor development has occurred in the postmarketing setting [see *Postmarketing Experience (6.2)*].

During ALTUVIIIIO clinical studies (median treatment duration 96.3 weeks), 4/276 (1.4%) of evaluable patients developed transient treatment emergent anti-drug antibodies.

No impact of ADAs on the FVIII activity levels or PK exposure parameters was observed.

No impact of ADAs with respect to bleeding episodes pharmacodynamic response, or safety was noted.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No carcinogenicity or mutagenicity studies have been conducted with of ALTUVIIIIO. No studies have been conducted to evaluate the effects of ALTUVIIIIO on fertility.

14 CLINICAL STUDIES

The safety, efficacy, and pharmacokinetics of ALTUVIIIIO were evaluated in two multicenter, prospective, open-label clinical studies, Study 1 (NCT04161495) and Study 2 (NCT04759193), as described below.

All studies evaluated the efficacy of routine prophylaxis with a weekly dose of 50 IU/kg and determined hemostatic efficacy in the treatment of bleeding episodes and during perioperative management in patients undergoing major or minor surgical procedures.

Routine Prophylaxis to Reduce Bleeding Episodes

Study 1 (Adult and Adolescent Study)

Study 1 enrolled a total of 159 previously treated patients (PTPs) including 158 male and 1 female patients with severe hemophilia A (<1% endogenous Factor VIII activity or a documented genetic mutation). Patients were aged 12 to 72 years and included 25 adolescent patients aged 12 to 17 years. All 159 enrolled patients received at least one dose of ALTUVIIIIO and were evaluable for efficacy. A total of 149 patients (93.7%) completed the study.

The efficacy of weekly 50 IU/kg ALTUVIIIIO as routine prophylaxis was evaluated as estimated by the mean annualized bleed rate (ABR) and by comparing the ABR during on-study prophylaxis vs. the ABR during pre-study FVIII prophylaxis. A total of 133 adults and adolescents, who were on pre-study FVIII prophylaxis, were assigned to receive ALTUVIIIIO for routine prophylaxis at a dose of 50 IU/kg IV once weekly for 52 weeks (Arm A). An additional 26 patients, who were on pre-study episodic (on-demand) treatment with FVIII, received episodic (on-demand) treatment with ALTUVIIIIO at doses of 50 IU/kg IV for 26 weeks, followed by routine prophylaxis at a dose of 50 IU/kg IV

once weekly for 26 weeks (Arm B). Overall, 115 patients received at least a total number of 50 exposure days (EDs) in Arm A and 17 patients completed at least 25 EDs of routine prophylaxis in Arm B.

The ABR in patients evaluable for efficacy with at least 26 weeks of exposure are summarized in Table 6.

Table 6: Summary of Annualized Bleeding Rate (ABR) with ALTUVIII O Prophylaxis, ALTUVIII O On-demand Treatment, and After Switch to ALTUVIII O Prophylaxis in Patients ≥ 12 Years of Age (Study 1)

Endpoint*	Arm A Prophylaxis†	Arm B On-demand‡	Arm B Prophylaxis‡
	N = 128	N = 26	N = 26
Treated bleeds			
Mean ABR (95% CI)§	0.7 (0.5, 1.0)	21.4 (18.8, 24.4)	0.7 (0.3, 1.5)
Median ABR (Q1, Q3)	0 (0, 1.0)	21.1 (15.1, 27.1)	0 (0, 0)
% patients with zero bleeds, n (%)	82 (64.1)	0	20 (76.9)
Treated spontaneous bleeds			
Mean ABR (95% CI)§	0.3 (0.2, 0.4)	15.8 (12.3, 20.4)	0.4 (0.2, 1.2)
Median ABR (Q1, Q3)	0 (0, 0)	16.7 (8.6, 23.8)	0 (0, 0)
% patients with zero bleeds, n (%)	103 (80.5)	1 (3.8)	22 (84.6)
Treated joint bleeds			
Mean ABR (95% CI)§	0.5 (0.4, 0.7)	17.5 (14.9, 20.5)	0.6 (0.3, 1.5)
Median ABR (Q1, Q3)	0 (0, 1.0)	18.4 (10.8, 23.9)	0 (0, 0)
% patients with zero bleeds, n (%)	92 (71.9)	0	21 (80.8)
All Bleeds (treated and untreated)*			
Mean ABR (95% CI)§	1.1 (0.8, 1.5)	22.2 (19.4, 25.4)	0.9 (0.4, 1.8)
Median ABR (Q1, Q3)	0 (0, 1.2)	21.1 (16.8, 27.1)	0 (0, 1.9)
% patients with zero bleeds, n (%)	71 (55.5)	0	19 (73.1)

ABR = annualized bleed rate; CI = confidence interval; Q1 = 25th percentile, Q3 = 75th percentile.

* Reflects all bleeds reported by patients including those where no ALTUVIIIIO was administered.

† Patients assigned to receive ALTUVIIIIO prophylaxis for 52 weeks.

‡ Patients assigned to receive ALTUVIIIIO for 26 weeks.

§ Based on negative binomial model.

An intra-patient comparison (N = 78) between mean ABR during on-study prophylaxis with ALTUVIIIIO and that during pre-study FVIII prophylaxis yielded a 77% reduction in treated bleeds (95% CI: 58%, 87%).

All patients with target joints at baseline (defined as ≥ 3 spontaneous bleeding episodes in a major joint which occurred in a consecutive 6-month period) achieved resolution of all target joints (45/45, 100%) with 12 months of prophylactic treatment with ALTUVIIIIO (defined as ≤ 2 bleeding episodes in the target joint in 12 months).

Study 2 (Pediatric Study)

Study 2 enrolled 74 male PTPs <12 years of age with severe hemophilia A (38 patients were 1 to 5 years of age and 36 were 6 to 11 years of age). Of the 74 enrolled patients, all received at least 1 dose of ALTUVIIIIO. Seventy-two patients were evaluable for efficacy.

The efficacy of weekly 50 IU/kg ALTUVIIIIO as routine prophylaxis in children <12 years was evaluated as estimated by the mean annualized bleed rate (ABR). A total of 74 children (38 children <6 years of age and 36 children 6 to <12 years of age) were enrolled to receive ALTUVIIIIO for routine prophylaxis at a dose of 50 IU/kg IV once weekly for 52 weeks.

The ABR in patients evaluable for efficacy with at least 26 weeks of exposure are summarized in Table 7.

Table 7: Summary of Annualized Bleeding Rate (ABR) with ALTUVIIIIO Prophylaxis in Patients <12 Years of Age (Study 2)

Endpoint*	<6 years N = 37 [†]	6 to <12 years N = 35 [‡]	Overall N = 72 ^{†‡}
Treated bleeds			
Mean ABR (95% CI) [§]	0.5 (0.3, 0.8)	0.8 (0.4, 1.4)	0.6 (0.4, 0.9)
Median ABR (Q1, Q3)	0 (0, 1.0)	0 (0, 1.1)	0 (0, 1.0)
% patients with zero bleeds, n (%)	23 (62.2)	23 (65.7)	46 (63.9)
Treated spontaneous bleeds			
Mean ABR (95% CI) [§]	0.2 (0.1, 0.4)	0.2 (0, 0.6)	0.2 (0.1, 0.3)
Median ABR (Q1, Q3)	0 (0, 0)	0 (0, 0)	0 (0, 0)
% patients with zero bleeds, n (%)	31 (83.8)	32 (91.4)	63 (87.5)
Treated joint bleeds			

Mean ABR (95% CI) [§]	0.2 (0.1, 0.6)	0.4 (0.2, 0.9)	0.3 (0.2, 0.6)
Median ABR (Q1, Q3)	0 (0, 0)	0 (0, 0)	0 (0, 0)
% patients with zero bleeds, n (%)	33 (89.2)	27 (77.1)	60 (83.3)
All Bleeds (treated and untreated)*			
Mean ABR (95% CI) [§]	2.8 (1.4, 5.6)	2.3 (1.3, 4.1)	2.6 (1.6, 4.0)
Median ABR (Q1, Q3)	0 (0, 2.0)	1.0 (0, 2.9)	0.5 (0, 2.1)
% patients with zero bleeds, n (%)	20 (54.1)	16 (45.7)	36 (50.0)

ABR = annualized bleed rate; CI = confidence interval; Q1 = 25th percentile, Q3 = 75th percentile.

* Reflects all bleeds reported by patients including those where no ALTUVIIIIO was administered.

† A patient in the <6 years old age group who had a positive inhibitor result at baseline (pre-exposure to study drug) and was withdrawn after 3 EDs was excluded.

‡ A patient in the 6 to <12 years old age group who received an intense consolidation treatment (2 to 3 injections per week) for 15 weeks, after 2 traumatic hip joint bleeds was excluded from the efficacy analysis as the patient did not receive the weekly prophylaxis treatment as specified in the protocol for an extended period.

§ Based on negative binomial model.

Efficacy in Control of Bleeding

In the adult and adolescent study, a total of 362 bleeding episodes were treated with ALTUVIIIIO, most occurring during on-demand treatment in Arm B. Majority of bleeding episodes were localized in joints. Response to the first injection was assessed by patients at least 8 hours after treatment. A 4-point rating scale of excellent, good, moderate, and no response was used to assess response. Bleeding was resolved with a single 50 IU/kg injection of ALTUVIIIIO in 96.7% of bleeding episodes. The median (Q1; Q3) total dose to treat a bleeding episode was 50.9 IU/kg (50.0; 51.9). Control of bleeding episodes was similar across the treatment arms.

The efficacy of ALTUVIIIIO in control of bleeding in children <12 years of age was assessed in the pediatric study. A total of 43 bleeding episodes were treated with ALTUVIIIIO. Bleeding was resolved with a single 50 IU/kg injection of ALTUVIIIIO in 95.3% of bleeding episodes. The median (Q1; Q3) total dose to treat a bleeding episode was 52.6 IU/kg (50.0; 55.8).

Perioperative Management of Bleeding

Perioperative hemostasis was assessed in 14 major surgeries in 13 patients (10 adults and 3 children) across the adult and adolescent and pediatric clinical studies. Of the 14 major surgeries, 13 surgeries required a single pre-operative dose to maintain hemostasis during surgery; for 1 major surgery during routine prophylaxis no pre-operative loading dose was administered on the day of/or before surgery. The median pre-operative dose per surgery was 50 IU/kg (range 12.7 – 61.9).

The clinical evaluation of hemostatic response during major surgery was assessed using a 4-point scale of excellent, good, moderate, or poor/none. The hemostatic effect of ALTUVIIIIO was rated as "excellent" in 14 of 14 surgeries (100%).

Types of major surgeries assessed include major orthopedic procedures such as arthroplasties (joint replacements of knee, hip, and elbow), joint revisions and ankle fusion. Other major surgeries included molar tooth extractions, dental restoration and tooth extraction, circumcision, and rhinoplasty/mentoplasty.

Perioperative hemostasis was assessed in 27 minor surgeries in 23 patients (12 adults and 11 adolescents and children). The hemostatic response was evaluated by the investigator/surgeon in 21 of these minor surgeries; an excellent response was reported in all (100%).

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

ALTUVIIIIO is supplied in kits comprising a single-dose vial containing nominally, 250, 500, 750, 1000, 2000, 3000, or 4000 international units (IU) of Factor VIII potency, a prefilled syringe with 3 mL sterile water for injection, and a sterile vial adapter (reconstitution device). The actual amount of ALTUVIIIIO in IU is stated on the label and carton of each vial.

Not made with natural rubber latex.

Strength	Potency Color Code	Kit NDC Number
250 IU	Yellow	71104-978-01
500 IU	Red	71104-979-01
750 IU	Garnet	71104-980-01
1000 IU	Green	71104-981-01
2000 IU	Royal Blue	71104-982-01
3000 IU	Mist Grey	71104-983-01
4000 IU	Orange	71104-984-01

Not all pack sizes may be marketed.

Storage and Handling

Prior to reconstitution:

- Store ALTUVIIIIO in the original package to protect the ALTUVIIIIO vials from light.
- Store ALTUVIIIIO in powder form at 2°C to 8°C (36°F to 46°F). Do not freeze to avoid damage to the prefilled diluent syringe.
- ALTUVIIIIO may be stored at room temperature, not to exceed 30°C (86°F), for a single period of up to 6 months, within the expiration date printed on the label.
- If stored at room temperature, record the date that ALTUVIIIIO is removed from refrigeration on the carton in the area provided. After storage at room temperature, do not return the product to the refrigerator.
- Do not use beyond the expiration date printed on the vial or 6 months after the date that was written on the carton, whichever is earlier.

After Reconstitution:

- The reconstituted product may be stored at room temperature, not to exceed 30°C

(86°F), for up to 3 hours. Protect from direct sunlight. After reconstitution, if the product is not used within 3 hours, it must be discarded.

- Do not use ALTUVIIIIO if the reconstituted solution is cloudy or has particulate matter.
- Discard any unused ALTUVIIIIO.

17 PATIENT COUNSELING INFORMATION

Advise the patients to:

- Read the FDA-approved patient labeling (Patient Information and Instructions for Use).
- Call their healthcare provider or go to the emergency department right away if a hypersensitivity reaction occurs. Early signs of hypersensitivity reactions may include rash, hives, itching, facial swelling, tightness of the chest, and wheezing.
- Contact their healthcare provider or treatment facility for further treatment and/or assessment if they experience a lack of a clinical response to Factor VIII therapy because this may be a sign of inhibitor development.

Manufactured by:

Bioverativ Therapeutics Inc.

Waltham, MA 02451

A SANOFI COMPANY

US License Number 2078

©2025 Bioverativ Therapeutics Inc. All rights reserved.

For patent information: <https://www.sanofi.us/en/products-and-resources/patents>

ALTUVIIIIO is a registered trademark of Bioverativ Therapeutics Inc.

Patient Information

ALTUVIIIIO® (al too'vee oh)

[antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-ehl]

for intravenous use after reconstitution only

Single-dose vial

Please read this Patient Information carefully before using ALTUVIIIIO and each time you get a refill, as there may be new information. This Patient Information does not take the place of talking with your healthcare provider about your medical condition or your treatment.

What is the most important information I need to know about ALTUVIIIIO?

Do not attempt to give yourself an injection unless you have been taught how by your healthcare provider or hemophilia center.

You must carefully follow your healthcare provider's instructions regarding the dose and schedule for injecting ALTUVIIIIO so that your treatment will work best for you.

What is ALTUVIIIIO?

ALTUVIIIIO is an injectable medicine that is used to control and reduce the number of

bleeding episodes in people with Hemophilia A (congenital Factor VIII deficiency).

Your healthcare provider may give you ALTUVIIIIO when you have surgery.

Who should not use ALTUVIIIIO?

You should not use ALTUVIIIIO if you had an allergic reaction to it in the past.

What should I tell my healthcare provider before using ALTUVIIIIO?

Talk to your healthcare provider about:

- Any medical problems that you have or had.
- All prescription and non-prescription medicines that you take, including over-the-counter medicines, supplements or herbal medicines.
- Pregnancy or if you are planning to become pregnant. It is not known if ALTUVIIIIO may harm your unborn baby.
- Breastfeeding. It is not known if ALTUVIIIIO passes into the milk and if it can harm your baby.

How should I use ALTUVIIIIO?

You get ALTUVIIIIO as an injection into your vein. Your healthcare provider will instruct you on how to do injections on your own and may watch you give yourself the first dose of ALTUVIIIIO.

Contact your healthcare provider right away if bleeding is not controlled after using ALTUVIIIIO.

What are the possible side effects of ALTUVIIIIO?

You can have an allergic reaction to ALTUVIIIIO. Call your healthcare provider or emergency department right away if you have any of the following symptoms: difficulty breathing, chest tightness, swelling of the face, rash or hives.

Your body can also make antibodies called "inhibitors" against ALTUVIIIIO. This can stop ALTUVIIIIO from working properly. Your healthcare provider may give you blood tests to check for inhibitors.

The common side effects of ALTUVIIIIO are headache and joint pain.

These are not the only possible side effects of ALTUVIIIIO. Tell your healthcare provider about any side effect that bothers you or does not go away.

What are the ALTUVIIIIO dosage strengths?

ALTUVIIIIO comes in seven different dosage strengths with 3 mL sterile water for injection (sWFI). The actual number of international units (IU) of Factor VIII activity in the vial will be imprinted on the label and on the box. The seven different strengths are as follows:

Strength	Cap Color
250 IU	Yellow
500 IU	Red
750 IU	Garnet
1000 IU	Green
2000 IU	Royal Blue

3000 IU	Mist Grey
4000 IU	Orange

Always check the actual dosage strength printed on the label to make sure you are using the strength prescribed by your healthcare provider.

How should I store ALTUVIIIIO?

- Keep ALTUVIIIIO in its original package.
- Protect it from light.
- Do not freeze.
- Store refrigerated 2°C to 8°C (36°F to 46°F) up to 48 months or at room temperature [not to exceed 30°C (86°F)], for a single period up to 6 months. Do not use ALTUVIIIIO after the expiration date printed on the label and carton of each vial.
- When storing at room temperature:
 - Note on the carton the date on which the product is removed from refrigeration.
 - Use the product before the end of this 6-month period or discard it.
 - Do not return the product to the refrigerator.

After mixing with the diluent:

- Do not use ALTUVIIIIO if the mixed solution is not clear and colorless to slightly yellowish.
- Use mixed product as soon as possible.
- You may store mixed ALTUVIIIIO at room temperature, not to exceed 30°C (86°F), for up to **3 hours**. Protect the mixed ALTUVIIIIO from direct sunlight. Discard any mixed ALTUVIIIIO not used within 3 hours.

What else should I know about ALTUVIIIIO?

Medicines are sometimes prescribed for purposes other than those listed here. Do not use ALTUVIIIIO for a condition for which it was not prescribed. Do not share ALTUVIIIIO with other people, even if they have the same symptoms that you have.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Manufactured by:
Bioverativ Therapeutics Inc.
Waltham, MA 02451
A SANOFI COMPANY
US License Number 2078

©2024 Bioverativ Therapeutics Inc. All rights reserved.

ALTUVIIIIO is a registered trademark of Bioverativ Therapeutics Inc.

Revised: SEP 2024

INSTRUCTIONS FOR USE

ALTUVIIIIO® (al too'vee oh)

**[antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-ehI]
for intravenous use after reconstitution only
Single-dose vial**

Read the Instructions for Use before you start using ALTUVIIIIO and each time you get a

refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

Your healthcare provider should show you or your caregiver how to mix (reconstitute) and give ALTUVIIIIO the first time ALTUVIIIIO is used.

Any further questions? Ask your healthcare provider or call 1-800-633-1610.

Important Information You Need to Know Before Injecting ALTUVIIIIO

It is important that you do not try to inject ALTUVIIIIO unless you have received training from a healthcare provider.

- Read all of the instructions carefully before using the ALTUVIIIIO.
- ALTUVIIIIO is supplied as a single-dose vial of powder for mixing (reconstitution). Before you inject ALTUVIIIIO, you must mix (reconstitute) ALTUVIIIIO powder with liquid (diluent) that comes in a prefilled diluent syringe.
- After the powder and diluent are mixed the medicine is given in the vein (intravenous injection).
- The vial and the diluent syringe are not made with natural rubber latex.
- **Do not** use ALTUVIIIIO if it has been dropped on a hard surface or damaged.
- **Do not** use if the syringe cap has been removed or is not securely attached.
- **Do not** use the mixed ALTUVIIIIO if it contains visible particles or is cloudy.
- **Do not** give mixed ALTUVIIIIO in the same tubing or container with other medicines.

Storing ALTUVIIIIO

- Keep unused ALTUVIIIIO kit in the original carton and store in the refrigerator between 2°C and 8°C (36°F and 46°F). The product may be stored at room temperature up to 30°C (86°F). If stored at room temperature, the product (prior to mixing) expires after 6 months or after the expiration date on the product vial, whichever is earlier.
- Remove the product kit from the refrigerator and allow the ALTUVIIIIO vial and the prefilled diluent syringe to come to room temperature prior to injection.
- **Do not** use external heat sources such as putting the vial or prefilled diluent syringe in hot water.
- **Do not** return room temperature ALTUVIIIIO to refrigerator.
- Keep away from direct sunlight.
- **Do not** freeze.


Storing of Mixed (Reconstituted) ALTUVIIIIO

- ALTUVIIIIO should be given within **3 hours** after mixing.
- Keep away from direct sunlight.
- Do not refrigerate after mixing.
- Keep ALTUVIIIIO and all medicines out of the reach of children.

Preparing to Inject ALTUVIIIIO

Mixing (Reconstitution)

Step 1:



Look at the ALTUVIII O kit:

- Check that you have the correct medicine and dose.
- Check the expiration date.

⚠ Do not use ALTUVIII O if the expiration date has passed.

Step 2:




vial adapter



vial with
powdered
medicine



plunger rod



prefilled diluent syringe

**not included in
the carton:**



Wash your hands with soap and water.

Find a clean, flat work surface. Remove the supplies from the carton:


- Vial adapter in its package
- Vial with powdered medicine
- Plunger rod
- Prefilled diluent syringe

Also ensure you have the following supplies (not included in the carton):

- Infusion set
- Tourniquet
- 2 alcohol wipes
- 1 cotton ball or gauze pad
- 1 adhesive bandage, if required (See Step 22)
- 1 tape, if required (See Step 20)
- 1 larger luer lock syringe, if required (See Step 13)
- FDA-cleared sharps disposal container (See Step 23)

 **Do not** use ALTUVIIIIO (vial or prefilled diluent syringe) if it has been dropped on a hard surface or damaged.

Step 3:



Allow the ALTUVIIIIO vial and the prefilled diluent syringe to come to room temperature before use.

⚠ Do not use external heat sources such as putting the vial or prefilled diluent syringe in hot water.

⚠ Do not put ALTUVIIIIO in direct sunlight.

⚠ Do not return room temperature ALTUVIIIIO to the refrigerator.

Step 4:



Remove the plastic cap from the ALTUVIIIIO vial.

Wipe the rubber stopper of the vial with an alcohol wipe and allow it to dry.

⚠ After cleaning, **do not** touch the rubber stopper with your hand or allow it to touch any surface.


Step 5:



Completely remove the backing from the vial adapter package by peeling back the lid.

⚠ Do not remove the vial adapter from the package or touch the inside of the vial adapter.

Step 6:




Keep the vial on a flat surface. Hold the vial with one hand and using the other hand, place the vial adapter in its package over the vial.

Note: The spike should be placed directly above the center of the rubber stopper.


- Push the vial adapter straight down until the spike on the vial adapter punctures the center of the vial stopper and is fully inserted.

Step 7:



Lift the package cover away from the vial adapter and throw away the cover.


Step 8:



Note: Only use the prefilled diluent syringe provided to mix (reconstitute) the powdered medicine.

- Hold the plunger rod by the circular disk.
- Place the tip of the plunger rod into the end of the prefilled diluent syringe.
- Turn the plunger rod to the right until it is firmly attached.

Step 9:




With one hand, hold the prefilled diluent syringe directly under the cap with the cap pointing up.

Note: Make sure you are holding the prefilled diluent syringe by the ridged part directly under the cap.

 **Do not** use if the cap has been removed or is not securely attached.

Step 10:




With your other hand, grasp the cap and bend it at a 90-degree angle until it snaps off.

Note: After the cap snaps off, you will see the glass tip of the prefilled diluent syringe.

 **Do not** touch the glass tip of the prefilled diluent syringe or the inside of the cap.


Step 11:



Note: Be sure the vial is sitting on a flat surface.

- Insert the tip of the prefilled diluent syringe into the vial adapter opening.
- Turn the prefilled diluent syringe to the right until it is securely attached to the vial adapter.

Step 12:



Slowly push down on the plunger rod to inject all of the liquid (diluent) from the prefilled diluent syringe into the vial.

Note: The plunger rod may rise slightly afterward. This is normal.

Step 13:



With the prefilled diluent syringe still connected to the adapter, gently swirl the vial until the powder is completely dissolved.

Check the solution through the vial to make sure the powder is fully dissolved.


Note: The solution should look clear and colorless to slightly yellowish.

 **Do not** shake.

 **Do not** use the mixed (reconstituted) ALTUVIIIIO if it contains visible particles or is cloudy.

POOLING: pooling is the process of combining two or more reconstituted vials into a larger luer lock syringe (not provided in the carton). If the dose requires more than one vial, reconstitute each vial as described above (See Steps 4–13) with the prefilled diluent syringe provided. **Do not** detach the prefilled diluent syringe until you are ready to attach the larger luer lock syringe to the next vial. Keep the vial adapter attached to the vial as you will need it for attaching a larger luer lock syringe. Use a larger luer lock plastic syringe to combine the contents of the reconstituted vials into the syringe, similar as described in Steps 14–15. Repeat this pooling procedure with each vial you will be using. Once you have pooled the required dose, proceed with Step 16 (Injecting ALTUVIIIIO) using the larger luer lock syringe.

Step 14:




Make sure the plunger rod is pressed all the way down and the syringe is firmly attached to the vial adapter.

Turn the vial upside-down.

Slowly pull down on the plunger rod to draw all the solution from the vial into the syringe.

Note: Be careful not to pull the plunger rod completely out of the syringe.

Step 15:



Gently unscrew the syringe from the vial adapter by turning it to the right.

Throw away (dispose of) the vial with the adapter still attached (See Step 23).

If you are not ready to inject, put the syringe cap carefully back onto the syringe tip.

⚠ Do not touch the syringe tip or the inside of the cap.

Note: Mixed (reconstituted) ALTUVIII[®] should be given within **3 hours** after mixing. Protect from direct sunlight. Do not refrigerate after reconstitution.


Injecting ALTUVIIIIO

ALTUVIIIIO is given in the vein (intravenous injection) after mixing (reconstitution) of the powdered medicine with the diluent.

These are very generic instructions so your healthcare provider should teach you how to inject ALTUVIIIIO. After you have been taught to self-inject, you can follow these instructions.

⚠ Do not give mixed (reconstituted) ALTUVIIIIO if it contains particles, is discolored, or is cloudy.

Step 16:




Using aseptic technique (clean and germ free), attach the syringe to the connector end of the infusion set tubing by turning it to the right until securely attached.

⚠ Do not give mixed (reconstituted) ALTUVIIIIO in the same tubing or container with other medicines.


Note: Ask your healthcare provider which infusion set can be used with ALTUVIIIIO.

Step 17:



Apply a tourniquet to the upper arm. Then, using a new alcohol wipe, clean the skin where you will insert the needle and wait for it to dry.


Step 18:



Prime the syringe and the tubing. Push the plunger rod until all air is removed from the syringe and ALTUVIIIIO has filled the infusion set needle.

 **Do not** push ALTUVIIIIO through the needle.

Step 19:




Remove the protective needle cover from the infusion set needle and throw away (discard it).

 **Do not** touch the needle and do not put the needle cover back onto needle after use.

- Insert the needle on the infusion set tubing into the vein as instructed by your healthcare provider.


Step 20:



Use tape to secure the plastic wings of the needle in place at the injection site if needed.
Remove the tourniquet.

Note: Always make sure you have correctly inserted the needle into a vein when you perform an intravenous injection.

Step 21:




Slowly push the plunger rod on the syringe all the way down to give ALTUVIIIIO.

Your healthcare provider will provide your rate of administration based on your comfort level and the minimum injection time recommendation per vial.

Note: A small amount of medicine will be left in the infusion set after injection. This is normal.


Step 22:



After delivering ALTUVIIIIO, remove the tape and the needle from the vein. Use a cotton ball or gauze pad to put pressure on the injection site for several minutes to stop any bleeding.

Note: You may apply an adhesive bandage if needed.

Step 23:



Put the infusion set and the syringe in FDA-cleared sharps disposal container right away after use.

Dispose of all unused solution according to your local regulations.

Dispose of all empty vial(s), and other used medical supplies in your household trash.

⚠ Do not throw away (dispose of) the infusion set or syringe in your household trash.

Disposing of ALTUVIIIIO

Throw away (dispose of) the infusion set and the syringe in FDA-cleared sharps disposal container right away after use.

If you do not have FDA-cleared sharps disposal container you may use a household container that is:

- made of a heavy-duty plastic,

- can be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out,
- upright and stable during use,
- leak-resistant, and
- properly labeled to warn of hazardous waste inside the container.

When your FDA-cleared sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of FDA-cleared sharps disposal container. There may be state or local laws about how you should throw away used ALTUVIIIIO supplies.

 **Do not** dispose of your used FDA-cleared sharps disposal container in your household trash unless your community guidelines permit this.

 **Do not** recycle your used FDA-cleared sharps disposal container.

Keep your FDA-cleared sharps disposal container out of the reach of children.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Manufactured by:
Bioverativ Therapeutics Inc.
Waltham, MA 02451
A SANOFI COMPANY
US License Number 2078

©2025 Bioverativ Therapeutics Inc. All rights reserved.

For more information go to www.ALTUVIIIIO.com or call 1-800-633-1610

Revised: DEC 2025

PRINCIPAL DISPLAY PANEL - 250 IU Kit Carton

250 IU Nominal

NDC 71104-978-01

ALTUVIIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use

This package contains:

- (1) Single-dose vial containing lyophilized powder for reconstitution for intravenous injection
- (1) Prefilled diluent syringe containing 3 mL sterile water for injection (sWFI) with plunger rod
- (1) Sterile vial adapter reconstitution device
- (1) Package Insert

Rx Only


Manufactured by:
Bioverativ Therapeutics Inc.
Waltham, MA 02451

A SANOFI COMPANY
US License Number 2078

www.ALTUVIIIIO.com
1-800-633-1610

Diluent origin Germany
Plunger Rod origin France
Vial Adapter origin Israel

sanofi



250 IU Nominal

NDC 71104-978-01

ALTUVIIIIO™
[Antihemophilic Factor (Recombinant),
Fe-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use



Scan this code with your
smartphone camera for
more information.
Visit www.ALTUVIIIIO.com

sanofi

250 IU Nominal

ALTUVIIIIO™
[Antihemophilic Factor (Recombinant),
Fe-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use

Do not freeze.
Protect from light.
Reconstitute ALTUVIIIIO™ with accompanying
diluent.
Do not use past the expiration date stated on
the label.
The reconstituted product may be stored at
room temperature, not to exceed 30°C (86°F),
for up to 3 hours. Protect from direct sunlight.
Discard unused portion.
877998



Store unopened vials in a refrigerator at 2°C to 8°C
(36°F to 46°F) or at room temperature (not to
exceed 30°C (86°F)), for a single period up to
6 months. Do not return the product to the
refrigerator.
See prescribing information.

Dosage and Administration:
For Intravenous Use

ALTUVIIIIO™
[Antihemophilic Factor (Recombinant),
Fe-VWF-XTEN Fusion Protein-ehtl]

One single-dose vial with
3 mL pre-filled diluent syringe
GTIN: 00371104978015
LOT:
EXP:
IU:
SN:

250 IU Nominal



ALTUVIIIIO™
[Antihemophilic Factor (Recombinant),
Fe-VWF-XTEN Fusion Protein-ehtl]
When reconstituted with 3 mL sterile water for injection (SWFI),
the product contains arginine hydrochloride (250 mM), calcium
chloride dihydrate (5 mM), histidine (10 mM), polysorbate 80
(0.05% w/v), and sucrose (5% w/v).
Contains no preservatives.
Record when the carton is removed from refrigeration and
placed at room temperature (not to exceed 30°C (86°F)).
Date: / /


250 IU Nominal

NDC 71104-678-01

ALTUVIIIIO™
[Antihemophilic Factor (Recombinant),

250 IU Nominal

Rx Only
Manufactured by:
Bioverativ Therapeutics Inc.
Waltham, MA 01981



PRINCIPAL DISPLAY PANEL - 500 IU Kit Carton

500 IU Nominal

NDC 71104-979-01

ALTUVIII O™

[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use

This package contains:

- (1) Single-dose vial containing lyophilized powder for reconstitution for intravenous injection
- (1) Prefilled diluent syringe containing 3 mL sterile water for injection (sWFI) with plunger rod
- (1) Sterile vial adapter reconstitution device
- (1) Package Insert

Rx Only


Manufactured by:
Bioverativ Therapeutics Inc.
Waltham, MA 02451

A SANOFI COMPANY
US License Number 2078

www.ALTUVIII O.com
1-800-633-1610

Diluent origin Germany
Plunger Rod origin France
Vial Adapter origin Israel

sanofi



500 IU Nominal

NDC 71104-979-01

ALTUVIIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use



Scan this code with your
smartphone camera for
more information.
Visit www.ALTUVIIIIO.com

sanofi

500 IU Nominal

ALTUVIIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use

Do not freeze.
Reconstitute ALTUVIIIIO™ with accompanying
diluent.
Do not use past the expiration date stated on
the label.
The reconstituted product may be stored at
room temperature, not to exceed 30°C (86°F),
for up to 3 hours. Protect from direct sunlight.
Discard unused portion.
877999



Store unopened vials in a refrigerator at 2°C to 8°C
(36°F to 46°F) or at room temperature (not to
exceed 30°C (86°F)), for a single period up to
6 months. Do not return the product to the
refrigerator.

For Intravenous Use


[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

ALTUVIIIIO™

One single-dose vial with
3 mL prefilled diluent syringe
GTIN: 00371104979012

LOT:
EXP:
IU:
SN:

500 IU Nominal



[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]
When reconstituted with 3 mL sterile water for injection (SWFI),
the product contains arginine hydrochloride (250 mM), calcium
chloride dihydrate (5 mM), histidine (10 mM), polysorbate 80
(0.05% w/v), and sucrose (5% w/v).
Contains no preservatives.
Record when the carton is removed from refrigeration and
placed at room temperature (not to exceed 30°C (86°F)).
Date: ___/___/___


500 IU Nominal

NDC 71104-979-01

ALTUVIIIIO™
[Antihemophilic Factor (Recombinant)]

500 IU Nominal

Rx Only
Manufactured by:
Bioverativ Therapeutics Inc.



PRINCIPAL DISPLAY PANEL - 1000 IU Kit Carton

1000 IU Nominal

NDC 71104-981-01

ALTUVIIIO™

[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehl]

For Intravenous Use

This package contains:

- (1) Single-dose vial containing lyophilized powder for reconstitution for intravenous injection
- (1) Prefilled diluent syringe containing 3 mL sterile water for injection (sWFI) with plunger rod
- (1) Sterile vial adapter reconstitution device
- (1) Package Insert

Rx Only


Manufactured by:
Bioverativ Therapeutics Inc.
Waltham, MA 02451

A SANOFI COMPANY
US License Number 2078

www.ALTUVIIIO.com
1-800-633-1610

Diluent origin Germany
Plunger Rod origin France
Vial Adapter origin Israel

sanofi



1000 IU Nominal

NDC 71104-981-01

ALTUVIIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use



Scan this code with your
smartphone camera for
more information.
Visit www.ALTUVIIIIO.com

sanofi

1000 IU Nominal

ALTUVIIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use

Do not freeze.
Reconstitute ALTUVIIIIO™ with accompanying
diluent.
Do not use past the expiration date stated on
the label.
The reconstituted product may be stored at
room temperature, not to exceed 30°C (86°F),
for up to 3 hours. Protect from direct sunlight.
Discard unused portion.
878000



See prescribing information.
Store unopened vials in a refrigerator at 2°C to 8°C
(36°F to 46°F) or at room temperature (not to
exceed 30°C (86°F)), for a single period up to
6 months. Do not return the product to the
refrigerator.

Dosage and Administration:
For Intravenous Use

[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

ALTUVIIIIO™

One single-dose vial with
3 mL pre-filled diluent syringe
GTIN: 0037104981015
LOT:
EXP:
IU:
SN:

1000 IU Nominal



[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]
When reconstituted with 3 mL sterile water for injection (SWFI),
the product contains arginine hydrochloride (250 mM), calcium
chloride dihydrate (5 mM), histidine (10 mM), polysorbate 80
(0.05% w/v), and sucrose (5% w/v).
Contains no preservatives.
Record when the carton is removed from refrigeration and
placed at room temperature (not to exceed 30°C (86°F)).
Date: ___/___/___


1000 IU Nominal

NDC 71104-981-01

ALTUVIIIIO™
[Antihemophilic Factor (Recombinant)]

1000 IU Nominal

Rx Only
Manufactured by:
Bioverativ Therapeutics Inc.



PRINCIPAL DISPLAY PANEL - 2000 IU Kit Carton

2000 IU Nominal

NDC 71104-982-01

ALTUVIIIO™

[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehl]

For Intravenous Use

This package contains:

- (1) Single-dose vial containing lyophilized powder for reconstitution for intravenous injection
- (1) Prefilled diluent syringe containing 3 mL sterile water for injection (sWFI) with plunger rod
- (1) Sterile vial adapter reconstitution device
- (1) Package Insert

Rx Only


Manufactured by:
Bioverativ Therapeutics Inc.
Waltham, MA 02451

A SANOFI COMPANY
US License Number 2078

www.ALTUVIIIO.com
1-800-633-1610

Diluent origin Germany
Plunger Rod origin France
Vial Adapter origin Israel

sanofi



2000 IU Nominal

NDC 71104-982-01

ALTUVIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use



Scan this code with your
smartphone camera for
more information.
Visit www.ALTUVIIIO.com

sanofi

2000 IU Nominal

ALTUVIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use

Do not freeze.
Protect from light.
Reconstitute ALTUVIIIO™ with accompanying
diluent.
Do not use past the expiration date stated on
the label.
The reconstituted product may be stored at
room temperature, not to exceed 30°C (86°F),
for up to 3 hours. Protect from direct sunlight.
Discard unused portion. 878001



Store unopened vials in a refrigerator at 2°C to 8°C
(36°F to 46°F) or at room temperature (not to
exceed 30°C (86°F)), for a single period up to
6 months. Do not return the product to the
refrigerator.


ALTUVIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use

Dosage and Administration:

One single-dose vial with
3 mL pre-filled diluent syringe
GTIN: 00371104982012
LOT:
EXP:
IU:
SN:

2000 IU Nominal



3 71104 98201 2

ALTUVIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]
When reconstituted with 3 mL sterile water for injection (SWI),
the product contains arginine hydrochloride (250 mM), calcium
chloride dihydrate (5 mM), histidine (110 mM), polysorbate 80
(0.05% w/v), and sucrose (5% w/v).
Contains no preservatives.
Record when the carton is removed from refrigeration and
placed at room temperature (not to exceed 30°C (86°F)).
Date: ___/___/___

2000 IU Nominal

NDC 71104-982-01

ALTUVIIIO™
[Antihemophilic Factor (Recombinant),

2000 IU Nominal

Rx Only
Manufactured by:
Bioverativ Therapeutics Inc.
Waltham, MA 02451



PRINCIPAL DISPLAY PANEL - 3000 IU Kit Carton

3000 IU Nominal

NDC 71104-983-01

ALTUVIIIIO™

[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use

This package contains:

- (1) Single-dose vial containing lyophilized powder for reconstitution for intravenous injection
- (1) Prefilled diluent syringe containing 3 mL sterile water for injection (sWFI) with plunger rod
- (1) Sterile vial adapter reconstitution device
- (1) Package Insert

Rx Only


Manufactured by:
Bioverativ Therapeutics Inc.
Waltham, MA 02451

A SANOFI COMPANY
US License Number 2078

www.ALTUVIIIIO.com
1-800-633-1610

Diluent origin Germany
Plunger Rod origin France
Vial Adapter origin Israel

sanofi



3000 IU Nominal

NDC 71104-983-01

ALTUVIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehf]

For Intravenous Use



Scan this code with your
smartphone camera for
more information.
Visit www.ALTUVIIIO.com

sanofi

3000 IU Nominal

ALTUVIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehf]

For Intravenous Use

Do not freeze.
Protect from light.
Reconstitute ALTUVIIIO™ with accompanying
diluent.
Do not use past the expiration date stated on
the label.
The reconstituted product may be stored at
room temperature, not to exceed 30°C (86°F),
for up to 3 hours. Protect from direct sunlight.
Discard unused portion.
878002




Store unopened vials in a refrigerator at 2°C to 8°C
(36°F to 46°F) or at room temperature (not to
exceed 30°C (86°F)), for a single period up to
6 months. Do not return the product to the
refrigerator.
Dosage and Administration:
See prescribing information.

ALTUVIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehf]
For Intravenous Use

One single-dose vial with
3 mL pre-filled diluent syringe
GTIN: 00371104983019
LOT:
EXP:
IU:
SN:

3000 IU Nominal



ALTUVIIIO™
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehf]
When reconstituted with 3 mL sterile water for injection (SWFI),
the product contains arginine hydrochloride (250 mM), calcium
chloride dihydrate (5 mM), histidine (110 mM), poly sorbate 80
(0.05% w/v), and sucrose (5% w/v).
Contains no preservatives.
Record when the carton is removed from refrigeration and
placed at room temperature (not to exceed 30°C (86°F)).
Date: ___/___/___


3000 IU Nominal

3000 IU Nominal

NDC 71104-983-01

ALTUVIIIO™
[Antihemophilic Factor (Recombinant)]

Rx Only
Manufactured by:
Bioverativ Therapeutics Inc.



PRINCIPAL DISPLAY PANEL - 4000 IU Kit Carton

4000 IU Nominal

NDC 71104-984-01

ALTUVIIIIO™

[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl]

For Intravenous Use

This package contains:

- (1) Single-dose vial containing lyophilized powder for reconstitution for intravenous injection
- (1) Prefilled diluent syringe containing 3 mL sterile water for injection (sWFI) with plunger rod
- (1) Sterile vial adapter reconstitution device
- (1) Package Insert

Rx Only


Manufactured by:
Bioverativ Therapeutics Inc.
Waltham, MA 02451

A SANOFI COMPANY
US License Number 2078

www.ALTUVIIIIO.com
1-800-633-1610

Diluent origin Germany
Plunger Rod origin France
Vial Adapter origin Israel

sanofi



4000 IU Nominal

NDC 71104-984-01

ALTUVIII[®]
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehf)]

For Intravenous Use



Scan this code with your
smartphone camera for
more information.
Visit www.ALTUVIII.com

sanofi

4000 IU Nominal

ALTUVIII[®]
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehf)]

For Intravenous Use

Do not freeze.
Reconstitute ALTUVIII[®] with accompanying
diluent.
Do not use past the expiration date stated on
the label.
The reconstituted product may be stored at
room temperature, not to exceed 30°C (86°F),
for up to 3 hours. Protect from direct sunlight.
Discard unused portion.
878003



Store unopened vials in a refrigerator [not to
(36°F to 46°F) or at room temperature [not to
exceed 30°C (86°F)], for a single period up to
6 months. Do not return the product to the
refrigerator.

Dosage and Administration:
See prescribing information.


For Intravenous Use
Fc-VWF-XTEN Fusion Protein-ehf]

ALTUVIII[®]
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehf)]

One single-dose vial with
3 mL prefilled diluent syringe
GTIN: 0037104984016

LOT:
EXP:
IU:
SN:

4000 IU Nominal



ALTUVIII[®]
[Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehf)]
When reconstituted with 3 mL sterile water for injection (SWI),
the product contains arginine hydrochloride (250 mM), calcium
chloride dihydrate (5 mM), histidine (10 mM), polysorbate 80
(0.05% w/v), and sucrose (5% w/v).
Contains no preservatives.
Record when the carton is removed from refrigeration and
placed at room temperature (not to exceed 30°C (86°F)).
Date: ___/___/___

4000 IU Nominal

4000 IU Nominal

NDC 71104-984-01

ALTUVIII[®]
[Antihemophilic Factor (Recombinant)]

Rx Only
Manufactured by:
Bioverativ Therapeutics Inc.



ALTUVIIIIO

antihemophilic factor (recombinant), fc-vwf-xten fusion protein-eh1 kit

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71104-978
---------------------	-------------------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-978-01	1 in 1 KIT	02/22/2023	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 VIAL	3 mL
Part 2	1 SYRINGE	3 mL

Part 1 of 2

ALTUVIIIIO

efanesoctocog alfa powder, for solution

Product Information

Item Code (Source)	NDC:71104-985
Route of Administration	INTRAVENOUS

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EFANESOCTOCOG ALFA (UNII: WH7BHQ0RB4) (EFANESOCTOCOG ALFA - UNII:WH7BHQ0RB4)	EFANESOCTOCOG ALFA	250 [iU] in 3 mL

Inactive Ingredients

Ingredient Name	Strength
ARGININE HYDROCHLORIDE (UNII: F7LTH1E20Y)	
CALCIUM CHLORIDE (UNII: M4I0D6VV5M)	
HISTIDINE (UNII: 4QD397987E)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SUCROSE (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-985-08	3 mL in 1 VIAL; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Part 2 of 2

STERILE WATER

sterile water solution

Product Information

Item Code (Source)	NDC:71104-035
Route of Administration	INTRAVENOUS

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-035-01	3 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

ALTUVIIIIO

antihemophilic factor (recombinant), fc-vwf-xten fusion protein-ehtl kit

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71104-979
---------------------	-------------------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-979-01	1 in 1 KIT	02/22/2023	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 VIAL	3 mL
Part 2	1 SYRINGE	3 mL

Part 1 of 2

ALTUVIIIIO

efanesoctocog alfa powder, for solution

Product Information

Item Code (Source)	NDC:71104-986
Route of Administration	INTRAVENOUS

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EFANESCTOCOG ALFA (UNII: WH7BHQ0RB4) (EFANESCTOCOG ALFA - UNII:WH7BHQ0RB4)	EFANESCTOCOG ALFA	500 [iU] in 3 mL

Inactive Ingredients

Ingredient Name	Strength
ARGININE HYDROCHLORIDE (UNII: F7LTH1E20Y)	
CALCIUM CHLORIDE (UNII: M4I0D6VV5M)	
HISTIDINE (UNII: 4QD397987E)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SUCROSE (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-986-08	3 mL in 1 VIAL; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Part 2 of 2

STERILE WATER

sterile water solution

Product Information

Item Code (Source)	NDC:71104-035
Route of Administration	INTRAVENOUS

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-035-01	3 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

ALTUVIIIIO

antihemophilic factor (recombinant), fc-vwf-xten fusion protein-ehtl kit

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71104-981
---------------------	-------------------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-981-01	1 in 1 KIT	02/22/2023	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 VIAL	3 mL
Part 2	1 SYRINGE	3 mL

Part 1 of 2

ALTUVIIIIO

efanesoctocog alfa powder, for solution

Product Information

Item Code (Source)	NDC:71104-988
Route of Administration	INTRAVENOUS

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EFANESOCTOCOG ALFA (UNII: WH7BHQ0RB4) (EFANESOCTOCOG ALFA - UNII:WH7BHQ0RB4)	EFANESOCTOCOG ALFA	1000 [iU] in 3 mL

Inactive Ingredients

Ingredient Name	Strength
ARGININE HYDROCHLORIDE (UNII: F7LTH1E20Y)	
CALCIUM CHLORIDE (UNII: M4I0D6VV5M)	
HISTIDINE (UNII: 4QD397987E)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SUCROSE (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-988-08	3 mL in 1 VIAL; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Part 2 of 2

STERILE WATER

sterile water solution

Product Information

Item Code (Source)	NDC:71104-035
Route of Administration	INTRAVENOUS

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-035-01	3 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

ALTUVIIIIO

antihemophilic factor (recombinant), fc-wvf-xten fusion protein-ehtl kit

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71104-982
---------------------	-------------------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-982-01	1 in 1 KIT	02/22/2023	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 VIAL	3 mL
Part 2	1 SYRINGE	3 mL

Part 1 of 2

ALTUVIIIIO

efanesoctocog alfa powder, for solution

Product Information

Item Code (Source)	NDC:71104-989
Route of Administration	INTRAVENOUS

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EFANESOCTOCOG ALFA (UNII: WH7BHQ0RB4) (EFANESOCTOCOG ALFA - UNII:WH7BHQ0RB4)	EFANESOCTOCOG ALFA	2000 [iU] in 3 mL

Inactive Ingredients

Ingredient Name	Strength
ARGININE HYDROCHLORIDE (UNII: F7LTH1E20Y)	
CALCIUM CHLORIDE (UNII: M4I0D6VV5M)	
HISTIDINE (UNII: 4QD397987E)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SUCROSE (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-989-08	3 mL in 1 VIAL; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Part 2 of 2

STERILE WATER

sterile water solution

Product Information

Item Code (Source)	NDC:71104-035
Route of Administration	INTRAVENOUS

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-035-01	3 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

ALTUVIIIIO

antihemophilic factor (recombinant), fc-vwf-xten fusion protein-ehtl kit

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71104-983
--------------	-------------------------	--------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-983-01	1 in 1 KIT	02/22/2023	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 VIAL	3 mL
Part 2	1 SYRINGE	3 mL

Part 1 of 2

ALTUVIIIIO

efanesoctocog alfa powder, for solution

Product Information

Item Code (Source)	NDC:71104-990
Route of Administration	INTRAVENOUS

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EFANESOCTOCOG ALFA (UNII: WH7BHQ0RB4) (EFANESOCTOCOG ALFA - UNII:WH7BHQ0RB4)	EFANESOCTOCOG ALFA	3000 [iU] in 3 mL

Inactive Ingredients

Ingredient Name	Strength
ARGININE HYDROCHLORIDE (UNII: F7LTH1E20Y)	
CALCIUM CHLORIDE (UNII: M4I0D6VV5M)	
HISTIDINE (UNII: 4QD397987E)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SUCROSE (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-990-08	3 mL in 1 VIAL; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Part 2 of 2

STERILE WATER

sterile water solution

Product Information

Item Code (Source)	NDC:71104-035
Route of Administration	INTRAVENOUS

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-035-01	3 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

ALTUVIIIIO

antihemophilic factor (recombinant), fc-vwf-xten fusion protein-ehtl kit

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71104-984
---------------------	-------------------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-984-01	1 in 1 KIT	02/22/2023	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 VIAL	3 mL
Part 2	1 SYRINGE	3 mL

Part 1 of 2

ALTUVIIIIO

efanesoctocog alfa powder, for solution

Product Information

Item Code (Source)	NDC:71104-991
Route of Administration	INTRAVENOUS

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EFANESCTOCOG ALFA (UNII: WH7BHQ0RB4) (EFANESCTOCOG ALFA - UNII:WH7BHQ0RB4)	EFANESCTOCOG ALFA	4000 [iU] in 3 mL

Inactive Ingredients

Ingredient Name	Strength
ARGININE HYDROCHLORIDE (UNII: F7LTH1E20Y)	
CALCIUM CHLORIDE (UNII: M4I0D6VV5M)	
HISTIDINE (UNII: 4QD397987E)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SUCROSE (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-991-08	3 mL in 1 VIAL; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Part 2 of 2

STERILE WATER

sterile water solution

Product Information

Item Code (Source)	NDC:71104-035
Route of Administration	INTRAVENOUS

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71104-035-01	3 mL in 1 SYRINGE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125771	02/22/2023	

Labeler - Bioverativ Therapeutics Inc. (070517011)

Registrant - PPD Development Ireland Ltd. (985036175)

Establishment

Name	Address	ID/FEI	Business Operations
PPD Development Ireland Ltd.		985036175	ANALYSIS(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984)

Establishment

Name	Address	ID/FEI	Business Operations
Eurofins Lancaster Laboratories, Inc		069777290	ANALYSIS(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984)

Establishment

Name	Address	ID/FEI	Business Operations
Biogen U.S. Corporation		078734950	ANALYSIS(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984) , MANUFACTURE(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984)

Establishment

Name	Address	ID/FEI	Business Operations
Rechon Life Science AB		775207769	PACK(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984) , LABEL(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984)

Establishment

Name	Address	ID/FEI	Business Operations
Genzyme Corporation		050424395	PACK(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984) , LABEL(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984)

Establishment

Name	Address	ID/FEI	Business Operations
Vetter Pharma Fertigung GmbH & Co. KG (Langenargen Eisenbahnstrasse)		344217323	ANALYSIS(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984) , MANUFACTURE(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984)

Establishment

Name	Address	ID/FEI	Business Operations
Vetter Pharma Fertigung GmbH & Co. KG (Ravensburg Schuetzenstrasse)		316126754	ANALYSIS(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984)

Establishment

Name	Address	ID/FEI	Business Operations
Vetter Pharma Fertigung GmbH & Co. KG (Ravensburg Mooswiesen)		312670654	ANALYSIS(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984) , PACK(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984)

Establishment

Name	Address	ID/FEI	Business Operations
Genzyme Ireland Limited		985127419	ANALYSIS(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984) , MANUFACTURE(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984)

Establishment

Name	Address	ID/FEI	Business Operations
Biogen MA Inc.		841087823	API MANUFACTURE(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984)

Establishment

Name	Address	ID/FEI	Business Operations
Genzyme Flanders		372153895	ANALYSIS(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984) , API MANUFACTURE(71104-978, 71104-979, 71104-981, 71104-982, 71104-983, 71104-984)