

## **CHILDRENS PAIN AND FEVER- acetaminophen 160mg/5ml solution CalmCo LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **DrKids Children's Pain and Fever**

#### **Active Ingredient (in each 5mL)**

Acetaminophen, 160mg

#### **Purpose**

Pain reliever/fever reducer

#### **Uses**

Temporarily: • reduces fever

• relieves minor aches and pains due to: • the common cold • flu • headache • sore throat • toothache

#### **Warnings**

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes • more than 5 doses in 24 hours, which is the maximum daily amount for this product • With other drugs containing acetaminophen

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

**Ask a doctor before use if your child has** liver disease

**Ask a doctor or pharmacist before use if** your child is taking the blood thinning drug warfarin.

#### **Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical even if you do not notice any signs or symptoms.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- If your child is allergic to acetaminophen or any of the inactive ingredients in this product

### **Directions**

- **this product does not contain directions or complete warnings for adult use.**
- do not give more than directed (see overdose warning)
- use the chart below to confirm dose. If possible, use weight to confirm, otherwise use age
- give the recommended number of vials below following the instructions under "Easy to Use" on the side panel.
- empty each Single-Use Vial required of all liquid when dosing
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

### **Dosing Chart**

<b>Weight (lbs)</b>	<b>Age (Years)</b>	<b>Dose (mL)</b>
under 24 lbs	under 2 years	Talk to a doctor
24 -35 lbs	2 - 3 years	1 Single-Use Vial (5 mL)
36 - 47 lbs	4-5 years	Talk to a doctor
48 - 71 lbs	6-10 years	2 Single-Use Vial (10 mL)
72 - 95 lbs	11 years	3 Single-Use Vial (15 mL)

### **Other information**

- each Single-Use vial contains 8mg of sodium
- store at 68-77°F (20-25°C)
- tamper evident: do not use if top flap of carton is open. Do not use if any vials are open or broken

### **Inactive ingredients**

citric acid, cherry flavor, edetate disodium, glycerin, maltitol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium metabisulfite, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

## Questions or Comments?

+1 941-202-3270 or visit [www.DrKids.com](http://www.DrKids.com)

## Package



Single  
Use  
Vials



Ages 2-11  
years



# DrKids<sup>®</sup>

Children's

## Pain + Fever

Acetaminophen 160 mg / 5 mL

-  **Relieves pain**
-  **Reduces fevers**
-  **Soothes aches**
-  **Pre-measured dose**





**4** Pre-Measured  
Single-Use Vials  
0.17 fl.oz (5 mL) Each  
0.68 fl.oz (20 mL) Total

**Free From:**

- ✓ Alcohol
- ✓ Dyes
- ✓ Sugar



Made in the USA: Distributed by CalmCo LLC, Sarasota FL 34240

DKPF-04P-001-01

**Questions or Comments?** 941-202-3270 or visit [www.DrKids.com](http://www.DrKids.com)

**Inactive Ingredients**  
citric acid, cherry flavor, edetate disodium, glycerin, maltitol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium metabisulfite, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum.

**Other Information**  
■ Each Single-Use Vial contains 8mg of sodium. ■ Store at 68-77°F (20-25°C).  
■ Tamper evident: do not use if top flap of carton is open. Do not use if any vials are open or broken.

Weight (lbs)	Age (Years)	Dose (mL)
72-95 lbs.	11 years	3 single use vials (15 mL)
48-71 lbs.	6-10 years	2 single use vials (10 mL)
36-47 lbs.	4-5 years	Talk to a doctor
24-35 lbs.	2-3 years	1 single use vial (5 mL)
under 24 lbs.	under 2 years	Talk to a doctor

**Dosing Chart**  
Do not give for more than 5 days unless directed by a doctor.  
■ Do not give more than 5 times in 24 hours. ■ Symptoms last. ■ If needed, repeat dose every 4 hours while liquid when dosing. ■ Empty each Single Use Vial required of all recommended number of vials below following the instructions above "Easy to Use". ■ Give the possible, use weight to confirm, otherwise use age. ■ Use the chart below to confirm dose. ■ Do not give more than directed dose. If (see overdose warning). ■ Use the chart below to confirm dose. ■ Do not give more than directed dose or complete warnings for adult use. ■ This product does not contain directions or complete directions.

**Directions**  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical even if you do not notice any signs or symptoms.

**Drug Facts** (continued)

**Drug Facts**

Keep Outer Package for Complete Product Information

**Active Ingredient**  
Acetaminophen, 160 mg.....Pain reliever/fever reducer (in each 5 mL)

**Purpose**

**Uses** Temporarily: ■ Reduces fever ■ Relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache.

**Warnings**  
Liver Warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ More than 5 doses in 24 hours, which is the maximum daily amount ■ With other drugs containing acetaminophen.  
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**Stop use and ask doctor if:**  
■ Pain gets worse or last more than 5 days. ■ Fever gets worse or last more than 3 days. ■ New symptoms occur. ■ Redness or swelling is present. These could be signs of a serious condition.

**Drug Facts Continued** ▶

## Easy to Use

1 Tap the vial to eliminate air bubbles



2 Twist the top of the vial off



3 Squeeze the liquid into child's mouth



CHOKING HAZARD  
Small parts. Keep out  
of reach of children



## CHILDRENS PAIN AND FEVER

acetaminophen 160mg/5ml solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73282-3164(NDC:70302-130)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
WATER (UNII: 059QF0KO0R)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SORBITOL (UNII: 506T60A25R)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
MALTITOL (UNII: D65DG142WK)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
XANTHAN GUM (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73282-3164-2	4 in 1 POUCH	09/01/2022	
1		5 mL in 1 VIAL, SINGLE-DOSE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/01/2022	

**Labeler** - CalmCo LLC (079308308)**Registrant** - CalmCo LLC (079308308)

Revised: 7/2023

CalmCo LLC