

**DULCOLAX LAXATIVE- bisacodyl suppository**  
Chattem, Inc.

-----  
**Dulcolax Medicated Laxative Suppository**

***Dulcolax®***

***Medicated Laxative Suppositories***

***Drug Facts***

**Active ingredient (*in each suppository*)**

Bisacodyl (USP) 10 mg

**Purpose**

Stimulant laxative

**Use**

- for relief of occasional constipation and irregularity
- this product generally produces bowel movement in 15 minutes to 1 hour

**Warnings**

**For rectal use only**

**Ask a doctor before use if you have**

- stomach pain, nausea or vomiting
- noticed a sudden change in bowel habits that lasts more than 2 weeks

**When using this product**

it may cause stomach discomfort, faintness, rectal burning and mild cramps

**Stop use and ask a doctor if**

- you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- **Peel open plastic.** Insert suppository into rectum, pointed end first. Retain about 15 to 20 minutes.

---

adults and children 12 years of age and over	1 suppository in a single daily dose.
children 6 to under 12 years of age	½ suppository in a single daily dose
children under 6 years of age	ask a doctor

---

**Other information**

- store at 20°-25°C (68°-77°F)

**Inactive ingredient**

hard fat

**Questions?**

Call **1-866-844-2798** or visit **www.Dulcolax.com**

**Keep carton as it contains important product information.**

**PRINCIPAL DISPLAY PANEL**

Dulcolax  
Bisacodyl (USP) 10 mg Suppository/ Stimulant Laxative  
MEDICATED LAXATIVE  
SUPPOSITORY  
4 COMFORT SHAPED  
SUPPOSITORIES  
DO NOT SWALLOW

#1 DOCTOR RECOMMENDED\*

Easy-to-insert, comfort shaped Dulcolax® Suppositories provide fast relief from constipation. Trust Dulcolax® Suppositories to work within 15 minutes to 1 hour when inserted rectally.

Safety sealed: Do not use if individual blister unit is open or torn.

# Dulcolax®

Bisacodyl (USP) 10 mg Suppository/Stimulant Laxative

## MEDICATED LAXATIVE SUPPOSITORY

Effective, predictable relief

**FAST RELIEF IN MINUTES**



**4** COMFORT-SHAPED SUPPOSITORIES

\*Among OTC Stimulant Laxative brands

**DO NOT SWALLOW**

Dulcolax® does not supply store brands.

76/E

Dul. by: Owens, Inc.  
P.O. Box 22718  
Ocala, FL 32749-22718  
©2025 Owens by  
333-145



Questions? Call 1-866-844-2798 or visit [www.Dulcolax.com](http://www.Dulcolax.com)  
**Inactive ingredient** hard fat  
**Drug Facts** (continued)

**Drug Facts**  
**Active ingredient (in each suppository)** Bisacodyl (USP) 10 mg  
**Purpose** Stimulant laxative

**Use** ■ for relief of occasional constipation and irregularity  
■ this product generally produces bowel movement in 15 minutes to 1 hour

**Warnings**  
**For rectal use only**

**Ask a doctor before use if you have** ■ stomach pain, nausea or vomiting  
■ noticed a sudden change in bowel habits that lasts more than 2 weeks

**When using this product** it may cause stomach discomfort, flatness, rectal burning and mild cramps

**Stop use and ask a doctor if** ■ you have rectal bleeding or fail to have a bowel movement after using this product. These could be signs of a serious condition.  
■ you need to use a laxative for more than 1 week

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children**. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** ■ **Peel open plastic**. Insert suppository into rectum, pointed end first. Retain about 15 to 20 minutes.

adults and children 12 years of age and over 1 suppository in a single daily dose  
children 6 to under 12 years of age ½ suppository in a single daily dose  
children under 6 years of age ask a doctor

**Other information** ■ store at 20°-25°C (68°-77°F)

Keep carton as it contains important product information.

## DULCOLAX LAXATIVE

bisacodyl suppository

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41167-0210
<b>Route of Administration</b>	RECTAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISACODYL</b> (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	10 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>FAT, HARD</b> (UNII: 8334LX7S21)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	
<b>Shape</b>	BULLET	<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41167-0210-3	4 in 1 CARTON	06/04/2019	
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:41167-0210-5	7 in 1 CARTON	06/04/2019	
2		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:41167-0210-1	1 in 1 CARTON	06/04/2019	
3		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:41167-0210-2	2 in 1 CARTON	06/04/2019	
4		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	06/04/2019	

**Labeler** - Chattem, Inc. (003336013)

