

ACETAMINOPHEN- acetaminophen tablet

Major Pharmaceuticals

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Major Pharmaceuticals Acetaminophen Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 caplets every 6 hours while symptoms last• do not take more than 6 caplets in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

carnauba wax, corn starch*, croscarmellose sodium*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid

*may contain one or more of these ingredients

Questions or comments?

1-800-616-2471

Principal Display Panel

EXTRA STRENGTH

Acetaminophen

PAIN RELIEVER/FEVER REDUCER

ASPIRIN FREE

FOR ADULTS

ACTUAL SIZE

CAPLETS

Compare to the active ingredient in Extra Strength Tylenol® Caplets

1000 Caplets 500 mg. Each

MAJOR[®] NDC 0904-6720-80
EXTRA STRENGTH
Acetaminophen
PAIN RELIEVER/FEVER REDUCER
ASPIRIN FREE
FOR ADULTS
ACTUAL SIZE
CAPLETS
Compare to the active ingredient in Extra Strength Tylenol® Caplets**
1000 Caplets
500 mg. Each



DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts	
Active ingredient (in each caplet)	Purpose
Acetaminophen 500 mg.....	Pain reliever/fever reducer
Uses	
■ temporarily relieves minor aches and pains due to: ■ the common cold ■ headache ■ backache ■ minor pain of arthritis ■ muscular aches ■ toothache ■ premenstrual and menstrual cramps ■ temporarily reduces fever	
Warnings	
■ Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product ■ Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.	
Do not use	
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you have ever had an allergic reaction to this product or any of its ingredients	
Ask a doctor before use if you have liver disease	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin	
Stop use and ask a doctor if	
■ pain gets worse or lasts more than 10 days ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions	
■ do not take more than directed (see overdose warning)	
adults and children 12 years and over	■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor
Other information ■ store at 20-25°C (68-77°F)	
Inactive ingredients carnauba wax, corn starch*, croscarmellose sodium*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain one or more of these ingredients	
Questions or comments? 1-800-616-2471	

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Extra Strength Tylenol® Caplets.

12 25 616 2471

Distributed by
MAJOR® PHARMACEUTICALS
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152
M-05 REV. 06/18
Re-Order No. 7 00897



ACETAMINOPHEN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-6720
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	L484
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-6720-80	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/26/2018	
2	NDC:0904-6720-51	1 in 1 CARTON	07/30/2018	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0904-6720-59	1 in 1 CARTON	07/30/2018	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0904-6720-60	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/27/2018	
5	NDC:0904-6720-40	500 in 1 BOTTLE; Type 0: Not a Combination Product	08/28/2018	
6	NDC:0904-6720-24	1 in 1 CARTON	08/28/2018	
6		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	07/26/2018	

