

LEVOCETIRIZINE DIHYDROCHLORIDE- levocetirizine dihydrochloride tablet, coated

BJWC (Berkley & Jensen / BJ'S)

Levocetirizine Dihydrochloride Tablets USP, 5 mg

Active ingredient (in each tablet)

Levocetirizine dihydrochloride USP, 5 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings

Do not use

- if you have kidney disease
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have

- ever had trouble urinating or emptying your bladder

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask doctor if

- you have trouble urinating or emptying your bladder
- an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- if breast-feeding: not recommended

- if pregnant: ask a health professional before use

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

adults 65 years of age and older	<ul style="list-style-type: none"> • ask a doctor
adults and children 12-64 years of age	<ul style="list-style-type: none"> • take 1 tablet (5 mg) once daily in the evening • do not take more than 1 tablet (5 mg) in 24 hours • 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms
children 6-11 years of age	<ul style="list-style-type: none"> • take 1/2 tablet (2.5 mg) once daily in the evening • do not take more than 1/2 tablet (2.5 mg) in 24 hours
children under 6 years of age	<ul style="list-style-type: none"> • do not use
consumers with kidney disease	<ul style="list-style-type: none"> • do not use

Other information

- store between 20° and 25°C (68° and 77°F)
- safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients

colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions or comments?

Call **1-888-375-3784**

Levocetirizine Dihydrochloride Tablets, USP 5 mg carton label

**Levocetirizine
Dihydrochloride
Tablets USP, 5 mg**

Antihistamine

ORIGINAL PRESCRIPTION STRENGTH
ALLERGY RELIEF
LEVOCETIRIZINE DIHYDROCHLORIDE TABLETS USP, 5 mg

NDC 68391-965-55

24 HOUR

berkley jensen®

ANTIHISTAMINE
55 TABLETS

Active ingredient (in each tablet) Levocetirizine dihydrochloride USP, 5 mg.....**Purpose** Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings Do not use ■ if you have kidney disease ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine Ask a doctor before use if you have ■ ever had trouble urinating or emptying your bladder

When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if ■ you have trouble urinating or

(Continued On Back Of Label)

DISTRIBUTED BY:
BJ'S WHOLESALE CLUB, INC.
350 CAMPUS DRIVE
MARLBOROUGH, MA 01752 Made in India
800-BJ'S-CLUB
REV 06/24

150100627

LOT
EXP

Unprinted/Uncoated
For Lot & Exp

PEEL HERE

emptying your bladder ■ an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:** ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions adults 65 years of age and older ■ ask a doctor adults and children 12-64 years of age ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours ■ 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms children 6-11 years of age ■ take 1/2 tablet (2.5 mg) once daily in

the evening ■ do not take more than 1/2 tablet (2.5 mg) in 24 hours children under 6 years of age ■ do not use consumers with kidney disease ■ do not use

Other information ■ store between 20° and 25°C (68° and 77°F) ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions and comments? call 1-888-375-3784

NDC 68391-955-11

Compare to Xyzal® Allergy 24HR active ingredient*

ORIGINAL
PRESCRIPTION STRENGTH

ALLERGY RELIEF



LEVOCETIRIZINE DIHYDROCHLORIDE
TABLETS USP, 5 mg

ANTIHISTAMINE

- 24 HOUR RELIEF OF:**
- SNEEZING • RUNNY NOSE
 - ITCHY NOSE OR THROAT
 - ITCHY, WATERY EYES



110 TABLETS
— TWO .55 COUNT BOTTLES



ACTUAL SIZE

NDC 68391-955-11

Compare to Xyzal® Allergy 24HR active ingredient*

ORIGINAL
PRESCRIPTION STRENGTH

ALLERGY RELIEF



LEVOCETIRIZINE DIHYDROCHLORIDE
TABLETS USP, 5 mg

ANTIHISTAMINE

- 24 HOUR RELIEF OF:**
- SNEEZING • RUNNY NOSE
 - ITCHY NOSE OR THROAT
 - ITCHY, WATERY EYES



110 TABLETS
— TWO .55 COUNT BOTTLES



ACTUAL SIZE

- Clinically Proven Relief in 60 Minutes
- Clinically Proven Relief From Indoor And Outdoor Allergies
- Clinically Proven Relief For 24 Hours

Drug Facts		Drug Facts (continued)	
Active Ingredient (in each tablet) Levocetirizine dihydrochloride USP, 5 mg	Purpose Antihistamine	Directions	
Uses temporarily relieves these symptoms due to hay fever or other respiratory allergies: • runny nose • sneezing • itchy, watery eyes • itching of the nose or throat	Warnings Do not use • if you have kidney disease • if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine Ask a doctor before use if you have • ever had trouble urinating or emptying your bladder	adults 65 years of age and older • ask a doctor	adults and children 12-64 years of age • take 1 tablet (5 mg) once daily in the evening • do not take more than 1 tablet (5 mg) in 24 hours • 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms
When using this product • drowsiness may occur • avoid alcoholic drinks • alcohol, sedatives, and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery Stop use and ask a doctor if • you have trouble urinating or emptying your bladder • an allergic reaction to this product occurs. Seek medical help right away.	Other information • store between 20° and 25°C (68° and 77°F) • safety sealed; do not use if carton was opened or if printed foil inner seal on bottle is torn or missing	children 6-11 years of age • take 1/2 tablet (2.5 mg) once daily in the evening • do not take more than 1/2 tablet (2.5 mg) in 24 hours	children under 6 years of age • do not use
If pregnant or breast-feeding: • if breast-feeding: not recommended • if pregnant: ask a health professional before use Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).	Inactive ingredients colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide	consumers with kidney disease • do not use	• do not use
84F 320234 100 80% PRINTER MUST GENERATE UPC CODE WITH BELOW DIGITS 8 88670 18504 9 000000 000000	Questions or comments? Call 1-888-376-3784	Read directions and warnings before use. Keep this carton. It has important information.	

*This product is not manufactured or distributed by Chatham, Inc. (part of the Sanofi Group), distributor of Xyzal® Allergy 24HR Tablets. Xyzal® is a registered trademark of UCB Group of companies.

DISTRIBUTED BY: BJS WHOLESALE CLUB, INC.
350 CAMPUS DRIVE, MARLBOROUGH, MA 01752
800-BJS-CLUB

Made in India 1 5 0 1 0 0 6 2 6

REV 06/24

- Clinically Proven Relief in 60 Minutes
- Clinically Proven Relief From Indoor And Outdoor Allergies
- Clinically Proven Relief For 24 Hours

Drug Facts		Drug Facts (continued)	
Active ingredient (in each tablet) Levocetirizine dihydrochloride USP, 5 mg	Purpose Antihistamine	Directions	
Uses temporarily relieves these symptoms due to hay fever or other respiratory allergies: • runny nose • sneezing • itchy, watery eyes • itching of the nose or throat	Warnings Do not use • if you have kidney disease • if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine Ask a doctor before use if you have • ever had trouble urinating or emptying your bladder	adults 65 years of age and older • ask a doctor	adults and children 12-64 years of age • take 1 tablet (5 mg) once daily in the evening • do not take more than 1 tablet (5 mg) in 24 hours • 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms
When using this product • drowsiness may occur • avoid alcoholic drinks • alcohol, sedatives, and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery Stop use and ask a doctor if • you have trouble urinating or emptying your bladder • an allergic reaction to this product occurs. Seek medical help right away.	Other information • store between 20° and 25°C (68° and 77°F) • safety sealed; do not use if carton was opened or if printed foil inner seal on bottle is torn or missing	children 6-11 years of age • take 1/2 tablet (2.5 mg) once daily in the evening • do not take more than 1/2 tablet (2.5 mg) in 24 hours	children under 6 years of age • do not use
If pregnant or breast-feeding: • if breast-feeding: not recommended • if pregnant: ask a health professional before use Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).	Inactive ingredients colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide	consumers with kidney disease • do not use	• do not use
84F 320234 100 80% PRINTER MUST GENERATE UPC CODE WITH BELOW DIGITS 8 88670 18504 9 000000 000000	Questions or comments? Call 1-888-376-3784	Read directions and warnings before use. Keep this carton. It has important information.	

*This product is not manufactured or distributed by Chatham, Inc. (part of the Sanofi Group), distributor of Xyzal® Allergy 24HR Tablets. Xyzal® is a registered trademark of UCB Group of companies.

DISTRIBUTED BY: BJS WHOLESALE CLUB, INC.
350 CAMPUS DRIVE, MARLBOROUGH, MA 01752
800-BJS-CLUB

Made in India 1 5 0 1 0 0 6 2 6

REV 06/24

LEVOCETIRIZINE DIHYDROCHLORIDE

levocetirizine dihydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68391-955(NDC:43598-735)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
levocetirizine dihydrochloride (UNII: SOD6A38AGA) (levocetirizine - UNII:6U5EA9RT2O)	levocetirizine dihydrochloride	5 mg

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
Polyethylene Glycol, Unspecified (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	white	Score	2 pieces
Shape	OVAL	Size	9mm
Flavor		Imprint Code	L
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68391-955-11	2 in 1 PACKAGE	08/15/2024	
1	NDC:68391-955-55	55 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210375	06/24/2020	

Labeler - BJWC (Berkley & Jensen / BJ'S) (159082692)

Revised: 12/2025

BJWC (Berkley & Jensen / BJ'S)