

**EQUALINE TUSSIN MULTI SYMPTOM- dextromethorphan hydrobromide,  
guaifenesin, phenylephrine hydrochloride solution  
United Natural Foods, Inc. dba UNFI**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**SuperValu Inc. Tussin Multi-Symptom Drug Facts**

**Active ingredients (in each 10 mL)**

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg

Phenylephrine HCl, USP 10 mg

**Purposes**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

**Warnings**

**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**When using this product**

**do not use more than directed**

**Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

**Other information**

- **each 10 mL contains:** sodium 5 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients**

anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

**Questions or comments?**

**1-877-932-7948**

## **Principal Display Panel**

compare to Robitussin® Multi-Symptom Cold active ingredients

adult

tussin

multi-symptom

cold

dextromethorphan HBr (cough suppressant)

guaifenesin (expectorant)

phenylephrine HCl (nasal decongestant)

non-drowsy

peak cold

CF

relieves:

cough

nasal congestion

mucus

for ages 12 & over

ALCOHOL FREE

8 FL OZ (237mL)

NDC 41163-388-34

compare to  
Robitussin®  
Multi-Symptom Cold  
active ingredients\*

**EQUALINE®**

adult  
**tussin**  
multi-symptom  
cold

dextromethorphan HBr  
(cough suppressant)  
guaifenesin (expectorant)  
phenylephrine HCl  
(nasal decongestant)

non-drowsy  
peak cold

**CF**

relieves:

- cough
- nasal congestion
- mucus

for ages 12 & over  
ALCOHOL FREE



NDC 41163-388-34

compare to  
Robitussin®  
Multi-Symptom Cold  
active ingredients\*

**EQUALINE®**

adult  
**tussin**  
multi-symptom  
cold

dextromethorphan HBr  
(cough suppressant)  
guaifenesin (expectorant)  
phenylephrine HCl  
(nasal decongestant)

non-drowsy  
peak cold

**CF**

relieves:

- cough
- nasal congestion
- mucus


for ages 12 & over  
ALCOHOL FREE



0.5 FL OZ (14.8 mL)

8 FL OZ (237mL)

8 FL OZ (237mL)



(nasal decongestant)  
 phenylephrine HCl  
 guaifenesin (expectorant)  
 cough suppressant  
 dextromethorphan HBr  
 cold  
 multi-symptom  
**tussin**  
 adult  
**®EVALUINE**

### Drug Facts

**Active ingredients**      **Purposes**  
*(in each 10 mL)*  
 Dextromethorphan HBr,  
 USP 20 mg.....Cough suppressant  
 Guaifenesin, USP 200 mg.....Expectorant  
 Phenylephrine HCl,  
 USP 10 mg.....Nasal decongestant

#### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - nasal congestion
  - cough due to minor throat and bronchial irritation

#### Warnings

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- heart disease      ■ high blood pressure
- thyroid disease    ■ diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**When using this product do not use more than directed**

#### Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition. ▶

### Drug Facts (continued)

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

#### Other information

- each 10 mL contains: sodium 5 mg
- store at 20-25°C (68-77°F).  
Do not refrigerate.

**Inactive ingredients** anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

#### Questions or comments?

1-877-932-7948

\*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Multi-Symptom Cold.

**GLUTEN FREE**

Distributed by SUPERVALU INC.

Eden Prairie, MN 55344 USA

Contact us at 1-877-932-7948, or  
[www.supervalu-ourownbrands.com](http://www.supervalu-ourownbrands.com)

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



0 41163 46036 2

LOT NO.

EXP.

51634 EL C10

## EQUALINE TUSSIN MULTI SYMPTOM

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41163-388
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

**Product Characteristics**

<b>Color</b>	RED (clear)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	FRUIT	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-388-26	1 in 1 CARTON	07/11/2014	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41163-388-34	1 in 1 CARTON	07/11/2014	
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/11/2014	

**Labeler** - United Natural Foods, Inc. dba UNFI (943556183)

Revised: 8/2023

United Natural Foods, Inc. dba UNFI