

CITROMA- magnesium citrate liquid
Consumer Product Partners, LLC

Swan 166.000/166AA-166AB
Lemon Magnesium Citrate

Tamper Evident Statement

Do not use if tamper evident twist-off cap is missing, broken or separated from neckring

Active Ingredient (in each fl oz)

Magnesium citrate 1.745g

Purpose

Saline laxative

Uses

- for relief of occasional constipation (irregularity).
- Generally produces bowel movement in 1/2 to 6 hours

Warnings

for this product

Ask a doctor before use if you have

- kidney disease
- a magnesium restricted diet
- abdominal pain, nausea, or vomiting
- noticed a sudden change in bowel habits that persists over a period of 2 weeks
- already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are

- taking any other drug.
- Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding

ask help professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Store at temperatures between 46° and 86°F (8° and 30° C)

Directions

- shake well before using
- drink a full glass (8 ounces) of liquid with each dose
- may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over - 6.5 to 10 fl oz in 24 hours

children 6 to under 12 years of age - 3 to 7 fl oz maximum 7 fl oz in 24 hours

children 2 to under 6 years of age - 2 to 3 fl oz in 24 hours

children under 2 years of age - ask a doctor

other information

- **each fl oz contains:** magnesium 290 mg
- **each fl oz contains:** sodium 1 mg

Inactive ingredients

benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Disclaimer

Non-USP: Non-Carbonated, Non-Pasteurized; Contains Preservatives

Adverse Reaction

Distributed by:

Consumer Product Partners, LLC

St. Louis, MO 63114

1-888-593-0593

Principal display panel

swan®

Magnesium Citrate

Saline Laxative
 Oral Solution
 Low Sodium
 Lemon Flavor
 10 FL OZ (296 mL)

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKRING

swan[®]
Magnesium Citrate
 Saline Laxative Oral Solution

Low Sodium
 Lemon Flavor



10 FL OZ
 (296 mL)

Drug Facts
Active ingredient (in each fl oz) Purpose
 Magnesium citrate 1,745 g Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
 ■ Generally produces bowel movement in ½ to 6 hours

Warnings
 Ask a doctor before use if you have
 ■ kidney disease ■ a magnesium restricted diet
 ■ abdominal pain, nausea, or vomiting
 ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
 ■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
 ■ taking any other drug.
 ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help ▶

Drug Facts (continued)
 or contact a Poison Control Center right away.
 Store at temperatures between 46° and 86° F (8° and 32° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information
 ■ each fl oz contains: magnesium 290 mg
 ■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Non-USP: Non-Carbonated, Non-Pasteurized; Contains Preservatives

Distributed by:
 Consumer Product Partners, LLC
 St. Louis, MO 63114
 1-888-593-0593

0 7278514388 5



L0017544SD

Principal display panel

NO USAR SI LA ROSCA CON PRECINTO DE SEGURIDAD
FALTA, ESTÁ ROTA O ESTÁ SEPARADA DEL
ANILLO ALREDEDOR DEL CUELLO DE LA BOTELLA

CRUZ BLANCA

CITRATO DE MAGNESIO

LAXANTE SALINO, LIMÓN
INGREDIENTE ACTIVO (POR ONZA FLUIDA):
1.745 G DE CITRATO DE MAGNESIO
CONTENIDO DE ALCOHOL: 0.0125%
(PROVENIENTE DE LOS SABORIZANTES)
SOLUCION ORAL

**CONTENIDO NETO
10 ONZAS FLUIDAS
(296 ML)**

Fabricado por Consumer Product Partners, LLC
St. Louis, MO 63114 U.S.A. Dist.

INGREDIENTE ACTIVO (POR ONZA FLUIDA): 1.745 g DE CITRATO DE MAGNESIO.
PROPÓSITO: LAXANTE SALINO.

USO: PARA ALIVIAR EL ESTREÑIMIENTO OCASIONAL (IRREGULARIDAD).
GENERALMENTE PRODUCE MOVIMIENTO INTESTINAL EN UN PERÍODO DE 1/4 A 6 HORAS.
INSTRUCCIONES: *AGITAR BIEN ANTES DE USAR *BEBER UN VASO (8 ONZAS) DE LÍQUIDO CON
CADA DOSIS. PUEDE SER INGERIDO EN UNA SOLA DOSIS DIARIA O EN DOSIS DIVIDIDAS.
ADULTOS Y NIÑOS MAYORES DE 12 AÑOS: DE 6.5 A 10 ONZAS FLUIDAS, MÁXIMO 10 ONZAS
FLUIDAS EN 24 HORAS. NIÑOS DE 6 A 12 AÑOS: DE 3 A 7 ONZAS FLUIDAS, MÁXIMO 7 ONZAS
FLUIDAS EN 24 HORAS. NIÑOS DE 2 A 6 AÑOS: DE 2 A 3 ONZAS FLUIDAS, MÁXIMO 3 ONZAS
FLUIDAS EN 24 HORAS. CONSULTE A UN MÉDICO ANTES DE ADMINISTRAR A NIÑOS MENORES
DE DOS (2) AÑOS DE EDAD.

INGREDIENTES INACTIVOS: ÁCIDO BENZOICO, ÁCIDO CÍTRICO, ETILÉNDIAMINOTETRAACETATO
SODÍCO (EDTA), SABORIZANTE, SUCRALOSA, AGUA.

ADVERTENCIAS: CONSULTE A UN MÉDICO ANTES DE USAR SI USTED *PADOCE DE
ENFERMEDADES DE LOS RÍÑONES *SIGUE UNA DIETA RESTRINGIDA EN MAGNESIO
*SUFRE DE DOLOR, NÁUSEAS O VÓMITOS *OBSERVA UN CAMBIO REPENTINO EN SUS HÁBITOS
INTESTINALES QUE PERSISTE POR MÁS DE 2 SEMANAS *HA USADO UN LAXANTE POR MÁS DE
UNA SEMANA. CONSULTE A UN MÉDICO O UN FARMACÉUTICO ANTES DE USAR, SI USTED
*TOMA CUALQUIER OTRO MEDICAMENTO *TOMA ESTE PRODUCTO 2 O MÁS HORAS ANTES O
DESPUÉS DE OTROS MEDICAMENTOS. LOS LAXANTES PUEDEN AFECTAR CÓMO FUNCIONAN
OTROS MEDICAMENTOS. DEJE DE USAR Y CONSULTE A UN MÉDICO SI TIENE SANGRADO
RECTAL (RECTORRAGIA) O NO DEFECAR DESPUÉS DEL USO, ESTO PODRÍAN SER SÍNTOMAS DE
UNA ENFERMEDAD GRAVE. SI ESTÁ EMBARAZADA O LACTANDO, CONSULTE A UN PROFESIONAL
DE LA SALUD ANTES DE USAR, MANTENGA FUERA DEL ALCANCE DE LOS NIÑOS. EN CASO DE
SORBODOSIS, BUSQUE ASISTENCIA MÉDICA O LLAME DE INMEDIATO A UN CENTRO DE
TOXICOLOGÍA. ALMACENE A TEMPERATURAS ENTRE 46 °F Y 86 °F (8 °C Y 30 °C).
VENTA SIN PRESCRIPCIÓN MÉDICA.



REG. SAN. PAN. 107853

L0019614SD



CITROMA

magnesium citrate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11344-166
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM CITRATE (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM CITRATE	1.745 g in 29.6 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
EDETATE DISODIUM ANHYDROUS (UNII: 8NLQ36F6MM)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11344-166-38	296 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/18/2018	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	02/18/2018	

Labeler - Consumer Product Partners, LLC (119091520)

Registrant - Consumer Product Partners, LLC (119091520)

Establishment			
Name	Address	ID/FEI	Business Operations
Consumer Product Partners, LLC		119091514	manufacture(11344-166)

Revised: 3/2025

Consumer Product Partners, LLC