

DOCUSATE SODIUM- docusate sodium capsule
Cellchem Pharmaceutical Inc

Active Ingredient (in each capsule)	Purpose
Docusate sodium 100 mg.....	Laxative, stool softener
Stool Softener	

Uses

- Relief of occasional constipation
- softens the stools

Warnings

Do not use

- if you have
 - abdominal pain
 - nausea
 - fever
 - vomiting
- with other medicines or if you are presently taking mineral oil
- if you are on a low salt diet
- longer than 7 days

When using this product

- frequent or prolonged use may result in dependence on laxatives
- it may take up to 3 days for you to have a bowel movement

Stop use and ask a doctor if

- rectal bleeding occurs
- symptoms persist, last for more than 7 days

Keep out of reach of children. In case of overdose, call a poison control centre or get medical help right away, even if you do not notice any signs or symptoms.

Directions

- **Adults and children 12 years and over:**
- Take 1 - 2 capsules daily or as directed by physician
- Do not take more than 4 capsules per day.
- Drink lots of fluids (one full glass or more)

Other information

- Questions ?



1(844) 481-8884

Inactive ingredients

Polyethylene glycol 400, propylene glycol, gelatin, glycerin, sodium methyl paraben, sodium propyl paraben, sorbitol sorbitan solution, purified water, FD&C Red No. 40, FD&C Yellow No. 6.

Manufactured for :

CellChem Pharmaceuticals Inc.
Ottawa, ON, Canada K2E 7V7

<p>NDC 73147-8273-1</p> <h1>Docusate Sodium</h1> <p>Docusate Sodium Capsules USP 100 mg</p> <p>Stool Softener</p> <p>100 Capsules</p> <p>Manufactured for :</p>  <p>CellChem Pharmaceuticals Inc. Ottawa, ON, Canada K2E 7V7</p>	<table border="1"> <tr> <th colspan="2">Drug Facts</th> </tr> <tr> <th>Active ingredients (in each capsule)</th> <th>Purpose</th> </tr> <tr> <td>Docusate sodium 100 mg</td> <td>Laxative, stool softener</td> </tr> <tr> <td colspan="2">Uses •Relief of occasional constipation •softens the stools</td> </tr> <tr> <td colspan="2">Warnings Do not use •if you have •abdominal pain •nausea •fever •vomiting •with other medicines or if you are presently taking mineral oil •if you are on a low salt diet •longer than 7 days</td> </tr> <tr> <td colspan="2">Ask a doctor or pharmacist before use if you •noticed a sudden change in bowel habits that has lasted longer than 2 weeks (this may be a sign of a serious condition) •are pregnant or breastfeeding</td> </tr> <tr> <td colspan="2">When using this product •frequent or prolonged use may result in dependence on laxatives •it may take up to 3 days for you to have a bowel movement</td> </tr> <tr> <td colspan="2">Stop use and ask a doctor if •rectal bleeding occurs •symptoms persist, last for more than 7 days</td> </tr> <tr> <td colspan="2">Keep out of reach of children. In case of overdose, call a poison control centre or get medical help right away, even if you do not notice any signs or symptoms.</td> </tr> <tr> <td colspan="2">Directions •Adults and children 12 years and over: •Take 1 - 2 capsules daily or as directed by physician •Do not take more than 4 capsules per day. •Drink lots of fluids (one full glass or more)</td> </tr> <tr> <td colspan="2">Inactive ingredients Polyethylene glycol 400, propylene glycol, gelatin, glycerin, sodium methyl paraben, sodium propyl paraben, sorbitol sorbitan solution, purified water, FD&C Red No. 40, FD&C Yellow No. 6.</td> </tr> <tr> <td colspan="2">Questions? 1(844) 481-8884</td> </tr> </table>	Drug Facts		Active ingredients (in each capsule)	Purpose	Docusate sodium 100 mg	Laxative, stool softener	Uses •Relief of occasional constipation •softens the stools		Warnings Do not use •if you have •abdominal pain •nausea •fever •vomiting •with other medicines or if you are presently taking mineral oil •if you are on a low salt diet •longer than 7 days		Ask a doctor or pharmacist before use if you •noticed a sudden change in bowel habits that has lasted longer than 2 weeks (this may be a sign of a serious condition) •are pregnant or breastfeeding		When using this product •frequent or prolonged use may result in dependence on laxatives •it may take up to 3 days for you to have a bowel movement		Stop use and ask a doctor if •rectal bleeding occurs •symptoms persist, last for more than 7 days		Keep out of reach of children. In case of overdose, call a poison control centre or get medical help right away, even if you do not notice any signs or symptoms.		Directions •Adults and children 12 years and over: •Take 1 - 2 capsules daily or as directed by physician •Do not take more than 4 capsules per day. •Drink lots of fluids (one full glass or more)		Inactive ingredients Polyethylene glycol 400, propylene glycol, gelatin, glycerin, sodium methyl paraben, sodium propyl paraben, sorbitol sorbitan solution, purified water, FD&C Red No. 40, FD&C Yellow No. 6.		Questions? 1(844) 481-8884		<p>LOT:XXXXXXXX EXPIRY:MM/YYYY</p> <p>Product Code</p>  <p>73147-82731 2</p> <p>N 3</p>
Drug Facts																										
Active ingredients (in each capsule)	Purpose																									
Docusate sodium 100 mg	Laxative, stool softener																									
Uses •Relief of occasional constipation •softens the stools																										
Warnings Do not use •if you have •abdominal pain •nausea •fever •vomiting •with other medicines or if you are presently taking mineral oil •if you are on a low salt diet •longer than 7 days																										
Ask a doctor or pharmacist before use if you •noticed a sudden change in bowel habits that has lasted longer than 2 weeks (this may be a sign of a serious condition) •are pregnant or breastfeeding																										
When using this product •frequent or prolonged use may result in dependence on laxatives •it may take up to 3 days for you to have a bowel movement																										
Stop use and ask a doctor if •rectal bleeding occurs •symptoms persist, last for more than 7 days																										
Keep out of reach of children. In case of overdose, call a poison control centre or get medical help right away, even if you do not notice any signs or symptoms.																										
Directions •Adults and children 12 years and over: •Take 1 - 2 capsules daily or as directed by physician •Do not take more than 4 capsules per day. •Drink lots of fluids (one full glass or more)																										
Inactive ingredients Polyethylene glycol 400, propylene glycol, gelatin, glycerin, sodium methyl paraben, sodium propyl paraben, sorbitol sorbitan solution, purified water, FD&C Red No. 40, FD&C Yellow No. 6.																										
Questions? 1(844) 481-8884																										

DOCUSATE SODIUM

docusate sodium capsule

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73147-8273
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)	DOCUSATE SODIUM	100 mg

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN SODIUM (UNII: CR6K9C2NHK)	
PROPYLPARABEN SODIUM (UNII: 625NNB0G9N)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	red (Dark Red)	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73147-8273-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	10/10/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	10/10/2023	

Labeler - Cellchem Pharmaceutical Inc (111518618)

Establishment

Name	Address	ID/FEI	Business Operations
Softgel Healthcare Private Limited		675584180	manufacture(73147-8273)

Revised: 12/2023

Cellchem Pharmaceutical Inc