

# IMATINIB MESYLATE- imatinib mesylate tablet, film coated

## Apotex Corp

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use IMATINIB MESYLATE TABLETS safely and effectively. See full prescribing information for IMATINIB MESYLATE TABLETS.

#### IMATINIB MESYLATE tablets, for oral use

Initial U.S. Approval: 2001

#### INDICATIONS AND USAGE

Imatinib mesylate is a kinase inhibitor indicated for the treatment of:

- Newly diagnosed adult and pediatric patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase (1.1)
- Patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in blast crisis (BC), accelerated phase (AP), or in chronic phase (CP) after failure of interferon-alpha therapy (1.2)
- Adult patients with relapsed or refractory Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) (1.3)
- Pediatric patients with newly diagnosed Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) in combination with chemotherapy (1.4)
- Adult patients with myelodysplastic/myeloproliferative diseases (MDS/MPD) associated with platelet-derived growth factor receptor (PDGFR) gene re-arrangements. (1.5)
- Adult patients with aggressive systemic mastocytosis (ASM) without the D816V c-Kit mutation or with c-Kit mutational status unknown (1.6)
- Adult patients with hypereosinophilic syndrome (HES) and/or chronic eosinophilic leukemia (CEL) who have the FIP1L1-PDGFR $\alpha$  fusion kinase (mutational analysis or fluorescence in situ hybridization [FISH] demonstration of CHIC2 allele deletion) and for patients with HES and/or CEL who are FIP1L1-PDGFR $\alpha$  fusion kinase negative or unknown. (1.7)
- Adult patients with unresectable, recurrent and/or metastatic dermatofibrosarcoma protuberans (DFSP) (1.8)
- Patients with Kit (CD117) positive unresectable and/or metastatic malignant gastrointestinal stromal tumors (GIST). (1.9)
- Adjuvant treatment of adult patients following resection of Kit (CD117) positive GIST. (1.10)

#### DOSAGE AND ADMINISTRATION

|                                                           |                            |
|-----------------------------------------------------------|----------------------------|
| Adults with Ph+ CML CP (2.2):                             | 400 mg/day                 |
| Adults with Ph+ CML AP or BC (2.2):                       | 600 mg/day                 |
| Pediatrics with Ph+ CML CP (2.3):                         | 340 mg/m <sup>2</sup> /day |
| Adults with Ph+ ALL (2.4):                                | 600 mg/day                 |
| Pediatrics with Ph+ ALL (2.5)                             | 340 mg/m <sup>2</sup> /day |
| Adults with MDS/MPD (2.6):                                | 400 mg/day                 |
| Adults with ASM (2.7):                                    | 100 mg/day or 400 mg/day   |
| Adults with HES/CEL (2.8):                                | 100 mg/day or 400 mg/day   |
| Adults with DFSP (2.9):                                   | 800 mg/day                 |
| Adults with metastatic and/or unresectable GIST (2.10):   | 400 mg/day                 |
| Adjuvant treatment of adults with GIST (2.11):            | 400 mg/day                 |
| Patients with mild to moderate hepatic impairment (2.12): | 400 mg/day                 |
| Patients with severe hepatic impairment (2.12):           | 300 mg/day                 |

All doses of imatinib mesylate tablets should be taken with a meal and a large glass of water. Doses of 400 mg or 600 mg (imatinib as free base) should be administered once daily, whereas a dose of 800 mg (imatinib as free base) should be administered as 400 mg (imatinib as free base) twice a day. Imatinib mesylate tablets can be dissolved in water or apple juice for patients having difficulty swallowing. Daily dosing of 800 mg (imatinib as free base) and above should be accomplished using the 400 mg tablet (imatinib as free base) to reduce exposure to iron.

#### DOSAGE FORMS AND STRENGTHS

Tablets (with functional scoring): 100 mg and 400 mg (3)

#### CONTRAINDICATIONS

None (4).

#### WARNINGS AND PRECAUTIONS

- Edema and severe fluid retention have occurred. Weigh patients regularly and manage unexpected rapid weight gain by drug interruption and diuretics. (5.1, 6.1)
- Cytopenias, particularly anemia, neutropenia, and thrombocytopenia, have occurred. Manage with dose reduction, dose interruption, or discontinuation of treatment. Perform complete blood counts weekly for the first month, biweekly for the second month, and periodically thereafter. (5.2)
- Severe congestive heart failure and left ventricular dysfunction have been reported, particularly in patients with comorbidities and risk factors. Monitor and treat patients with cardiac disease or risk factors for cardiac failure. (5.3)
- Severe hepatotoxicity, including fatalities may occur. Assess liver function before initiation of treatment and monthly thereafter or as clinically indicated. Monitor liver function when combined with chemotherapy known to be associated with liver dysfunction. (5.4)
- Grade 3/4 hemorrhage has been reported in clinical studies in patients with newly diagnosed CML and with GIST. GI tumor sites may be the source of GI bleeds in GIST. (5.5)
- Gastrointestinal (GI) perforations, some fatal, have been reported. (5.6)
- Cardiogenic shock/left ventricular dysfunction has been associated with the initiation of imatinib mesylate in patients with conditions associated with high eosinophil levels (e.g., HES, MDS/MPD, and ASM). (5.7)
- Bullous dermatologic reactions (e.g., erythema multiforme and Stevens-Johnson syndrome) have been reported with the use of imatinib mesylate. (5.8)
- Hypothyroidism has been reported in thyroidectomy patients undergoing levothyroxine replacement. Closely monitor TSH levels in such patients. (5.9)
- Fetal harm can occur when administered to a pregnant woman. Apprise women of the potential harm to the fetus, and to use effective contraception. (5.10, 8.1)
- Growth retardation occurring in children and pre-adolescents receiving imatinib mesylate has been reported. Close monitoring of growth in children under imatinib mesylate treatment is recommended. (5.11, 6.2)

- Tumor Lysis Syndrome. Close monitoring is recommended. (5.12)
- Reports of motor vehicle accidents have been received in patients receiving imatinib mesylate. Caution patients about driving a car or operating machinery. (5.13)
- Renal Toxicity. A decline in renal function may occur in patients receiving imatinib mesylate tablets. Evaluate renal function at baseline and during therapy, with attention to risk factors for renal dysfunction. (5.14)

#### ADVERSE REACTIONS

The most frequently reported adverse reactions (greater than or equal to 30%) were edema, nausea, vomiting, muscle cramps, musculoskeletal pain, diarrhea, rash, fatigue and abdominal pain. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Apotex Corp. at 1-800-706-5575 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

#### DRUG INTERACTIONS

- CYP3A4 inducers may decrease imatinib mesylate  $C_{max}$  and area under curve (AUC). (2.12, 7.1, 12.3)
- CYP3A4 inhibitors may increase imatinib mesylate  $C_{max}$  and AUC. (7.2, 12.3)
- Imatinib mesylate is an inhibitor of CYP3A4 and CYP2D6 which may increase the  $C_{max}$  and AUC of other drugs. (7.3, 7.4, 12.3)
- Patients who require anticoagulation should receive low-molecular weight or standard heparin and not warfarin. (7.3)
- Imatinib mesylate tablets delay the clearance of methotrexate when methotrexate is used at high doses ( $> 500 \text{ mg/m}^2$ ). (7.5)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 2/2026

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

- 1.1 Newly Diagnosed Philadelphia Positive Chronic Myeloid Leukemia (Ph+ CML)
- 1.2 Ph+ CML in Blast Crisis (BC), Accelerated Phase (AP) or Chronic Phase (CP) After Interferon-alpha (IFN) Therapy
- 1.3 Adult Patients With Ph+ Acute Lymphoblastic Leukemia (ALL)
- 1.4 Pediatric Patients With Ph+ Acute Lymphoblastic Leukemia (ALL)
- 1.5 Myelodysplastic/Myeloproliferative Diseases (MDS/MPD)
- 1.6 Aggressive Systemic Mastocytosis (ASM)
- 1.7 Hypereosinophilic Syndrome (HES) and/or Chronic Eosinophilic Leukemia (CEL)
- 1.8 Dermatofibrosarcoma Protuberans (DFSP)
- 1.9 Kit+ Gastrointestinal Stromal Tumors (GIST)
- 1.10 Adjuvant Treatment of GIST

### 2 DOSAGE AND ADMINISTRATION

- 2.1 Drug Administration
- 2.2 Adult Patients With Ph+ CML CP, AP, or BC
- 2.3 Pediatric Patients With Ph+ CML CP
- 2.4 Adult Patients With Ph+ ALL
- 2.5 Pediatric Patients With Ph+ ALL
- 2.6 Adult Patients With MDS/MPD
- 2.7 Adult Patients With ASM
- 2.8 Adult Patients With HES/CEL
- 2.9 Adult Patients With DFSP
- 2.10 Adult Patients With Metastatic and/or Unresectable GIST
- 2.11 Adult Patients With Adjuvant GIST
- 2.12 Dose Modification Guidelines
- 2.13 Dose Adjustment for Hepatotoxicity and Non-Hematologic Adverse Reactions
- 2.14 Dose Adjustment for Hematologic Adverse Reactions

### 3 DOSAGE FORMS AND STRENGTHS

### 4 CONTRAINDICATIONS

### 5 WARNINGS AND PRECAUTIONS

- 5.1 Fluid Retention and Edema
- 5.2 Hematologic Toxicity
- 5.3 Congestive Heart Failure and Left Ventricular Dysfunction
- 5.4 Hepatotoxicity
- 5.5 Hemorrhage
- 5.6 Gastrointestinal Disorders
- 5.7 Hypereosinophilic Cardiac Toxicity
- 5.8 Dermatologic Toxicities
- 5.9 Hypothyroidism
- 5.10 Embryo-Fetal Toxicity
- 5.11 Growth Retardation in Children and Adolescents
- 5.12 Tumor Lysis Syndrome
- 5.13 Impairments Related to Driving and Using Machinery
- 5.14 Renal Toxicity

### 6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

### 7 DRUG INTERACTIONS

- 7.1 Agents Inducing CYP3A Metabolism
- 7.2 Agents Inhibiting CYP3A Metabolism
- 7.3 Interactions With Drugs Metabolized by CYP3A4
- 7.4 Interactions With Drugs Metabolized by CYP2D6
- 7.5 Interactions With Methotrexate

## **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment
- 8.7 Renal Impairment

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

## **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

## **14 CLINICAL STUDIES**

- 14.1 Chronic Myeloid Leukemia
- 14.2 Pediatric CML
- 14.3 Acute Lymphoblastic Leukemia
- 14.4 Pediatric ALL
- 14.5 Myelodysplastic/Myeloproliferative Diseases
- 14.6 Aggressive Systemic Mastocytosis
- 14.7 Hypereosinophilic Syndrome/Chronic Eosinophilic Leukemia
- 14.8 Dermatofibrosarcoma Protuberans
- 14.9 Gastrointestinal Stromal Tumors

## **15 REFERENCES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

#### **1.1 Newly Diagnosed Philadelphia Positive Chronic Myeloid Leukemia (Ph+ CML)**

Newly diagnosed adult and pediatric patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase.

#### **1.2 Ph+ CML in Blast Crisis (BC), Accelerated Phase (AP) or Chronic Phase (CP) After Interferon-alpha (IFN) Therapy**

Patients with Philadelphia chromosome positive chronic myeloid leukemia in blast crisis, accelerated phase, or in chronic phase after failure of interferon-alpha therapy.

#### **1.3 Adult Patients With Ph+ Acute Lymphoblastic Leukemia (ALL)**

Adult patients with relapsed or refractory Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL).

#### **1.4 Pediatric Patients With Ph+ Acute Lymphoblastic Leukemia (ALL)**

Pediatric patients with newly diagnosed Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) in combination with chemotherapy.

#### **1.5 Myelodysplastic/Myeloproliferative Diseases (MDS/MPD)**

Adult patients with myelodysplastic/myeloproliferative diseases associated with platelet-derived growth factor receptor (PDGFR) gene re-arrangements.

#### **1.6 Aggressive Systemic Mastocytosis (ASM)**

Adult patients with aggressive systemic mastocytosis without the D816V c-Kit mutation or with c-Kit mutational status unknown.

#### **1.7 Hypereosinophilic Syndrome (HES) and/or Chronic Eosinophilic Leukemia**

## **(CEL)**

Adult patients with hypereosinophilic syndrome and/or chronic eosinophilic leukemia who have the FIP1L1-PDGFR $\alpha$  fusion kinase (mutational analysis or fluorescence in situ hybridization [FISH] demonstration of CHIC2 allele deletion) and for patients with HES and/or CEL who are FIP1L1-PDGFR $\alpha$  fusion kinase negative or unknown.

### **1.8 Dermatofibrosarcoma Protuberans (DFSP)**

Adult patients with unresectable, recurrent and/or metastatic dermatofibrosarcoma protuberans.

### **1.9 Kit+ Gastrointestinal Stromal Tumors (GIST)**

Patients with Kit (CD117) positive unresectable and/or metastatic malignant gastrointestinal stromal tumors.

### **1.10 Adjuvant Treatment of GIST**

Adjuvant treatment of adult patients following complete gross resection of Kit (CD117) positive GIST.

## **2 DOSAGE AND ADMINISTRATION**

### **2.1 Drug Administration**

The prescribed dose should be administered orally, with a meal and a large glass of water. Doses of 400 mg or 600 mg (imatinib as free base) should be administered once daily, whereas a dose of 800 mg (imatinib as free base) should be administered as 400 mg (imatinib as free base) twice a day.

For patients unable to swallow the film-coated tablets, the tablets may be dispersed in a glass of water or apple juice. The required number of tablets should be placed in the appropriate volume of beverage (approximately 50 mL for a 100-mg tablet, and 200 mL for a 400-mg tablet) and stirred with a spoon. The suspension should be administered immediately after complete disintegration of the tablet(s).

For daily dosing of 800 mg (imatinib as free base) and above, dosing should be accomplished using the 400-mg (imatinib as free base) tablet to reduce exposure to iron.

Treatment may be continued as long as there is no evidence of progressive disease or unacceptable toxicity.

### **2.2 Adult Patients With Ph+ CML CP, AP, or BC**

The recommended dose of imatinib mesylate tablets is 400 mg/day (imatinib as free base) for adult patients in chronic phase CML and 600 mg/day (imatinib as free base) for adult patients in accelerated phase or blast crisis.

In CML, a dose increase from 400 mg to 600 mg (imatinib as free base) in adult patients with chronic phase disease, or from 600 mg to 800 mg (imatinib as free base) (given as 400 mg twice daily) in adult patients in accelerated phase or blast crisis may be considered in the absence of severe adverse drug reaction and severe non-leukemia related neutropenia or thrombocytopenia in the following circumstances: disease progression (at any time), failure to achieve a satisfactory hematologic response after at least 3 months of treatment, failure to achieve a cytogenetic response after 6 to 12 months of treatment, or loss of a previously achieved hematologic or cytogenetic response.

### **2.3 Pediatric Patients With Ph+ CML CP**

The recommended dose of imatinib mesylate tablets for children with newly diagnosed Ph+ CML is 340 mg/m<sup>2</sup>/day (imatinib as free base) (not to exceed 600 mg). Imatinib mesylate tablets treatment can be given as a once daily dose or the daily dose may be split into two—one portion dosed in the morning and one portion in the evening. There is no experience with imatinib mesylate treatment in children under 1 year of age.

### **2.4 Adult Patients With Ph+ ALL**

The recommended dose of imatinib mesylate tablets is 600 mg/day (imatinib as free base) for adult patients with relapsed/refractory Ph+ ALL.

### **2.5 Pediatric Patients With Ph+ ALL**

The recommended dose of imatinib mesylate tablets to be given in combination with chemotherapy to children with newly diagnosed Ph+ ALL is 340 mg/m<sup>2</sup>/day (not to exceed 600 mg). Imatinib mesylate tablets treatment can be given as a once daily dose.

## 2.6 Adult Patients With MDS/MPD

Determine PDGFRb gene rearrangements status prior to initiating treatment.

The recommended dose of imatinib mesylate tablets is 400 mg/day (imatinib as free base) for adult patients with MDS/MPD.

## 2.7 Adult Patients With ASM

Determine D816V c-Kit mutation status prior to initiating treatment.

The recommended dose of imatinib mesylate tablets is 400 mg/day (imatinib as free base) for adult patients with ASM without the D816V c-Kit mutation. If c-Kit mutational status is not known or unavailable, treatment with imatinib mesylate tablets 400 mg/day (imatinib as free base) may be considered for patients with ASM not responding satisfactorily to other therapies. For patients with ASM associated with eosinophilia, a clonal hematological disease related to the fusion kinase FIP1L1-PDGFR $\alpha$ , a starting dose of 100 mg/day (imatinib as free base) is recommended. Dose increase from 100 mg to 400 mg (imatinib as free base) for these patients may be considered in the absence of adverse drug reactions if assessments demonstrate an insufficient response to therapy.

## 2.8 Adult Patients With HES/CEL

The recommended dose of imatinib mesylate tablets is 400 mg/day (imatinib as free base) for adult patients with HES/CEL. For HES/CEL patients with demonstrated FIP1L1-PDGFR $\alpha$  fusion kinase, a starting dose of 100 mg/day (imatinib as free base) is recommended. Dose increase from 100 mg to 400 mg (imatinib as free base) for these patients may be considered in the absence of adverse drug reactions if assessments demonstrate an insufficient response to therapy.

## 2.9 Adult Patients With DFSP

The recommended dose of imatinib mesylate tablets is 800 mg/day (imatinib as free base) for adult patients with DFSP.

## 2.10 Adult Patients With Metastatic and/or Unresectable GIST

The recommended dose of imatinib mesylate tablets is 400 mg/day (imatinib as free base) for adult patients with unresectable and/or metastatic, malignant GIST. A dose increase up to 800 mg daily (imatinib as free base) (given as 400 mg twice daily) may be considered, as clinically indicated, in patients showing clear signs or symptoms of disease progression at a lower dose and in the absence of severe adverse drug reactions.

## 2.11 Adult Patients With Adjuvant GIST

The recommended dose of imatinib mesylate tablets is 400 mg/day (imatinib as free base) for the adjuvant treatment of adult patients following complete gross resection of GIST. In clinical trials, one year of imatinib mesylate tablets and three years of imatinib mesylate tablets were studied. In the patient population defined in Study 2, three years of imatinib mesylate tablets is recommended [see *Clinical Studies* (14.8)]. The optimal treatment duration with imatinib mesylate tablets is not known.

## 2.12 Dose Modification Guidelines

*Concomitant Strong CYP3A4 inducers:* The use of concomitant strong CYP3A4 inducers should be avoided (e.g., dexamethasone, phenytoin, carbamazepine, rifampin, rifabutin, rifampacin, phenobarbital). If patients must be coadministered a strong CYP3A4 inducer, based on pharmacokinetic studies, the dosage of imatinib mesylate tablets should be increased by at least 50%, and clinical response should be carefully monitored [see *Drug Interactions* (7.1)].

*Hepatic Impairment:* Patients with mild and moderate hepatic impairment do not require a dose adjustment and should be treated per the recommended dose. A 25% decrease in the recommended dose should be used for patients with severe hepatic impairment [see *Use in Specific Populations* (8.6)].

*Renal Impairment:* Patients with moderate renal impairment (creatinine clearance [CrCL] = 20 to 39 mL/min) should receive a 50% decrease in the recommended starting dose and future doses can be increased as tolerated. Doses greater than 600 mg (imatinib as free base) are not recommended in patients with mild renal impairment (CrCL = 40 to 59 mL/min). For patients with moderate renal impairment doses greater than 400 mg (imatinib as free base) are not recommended.

Imatinib should be used with caution in patients with severe renal impairment. A dose of 100 mg/day was tolerated in two patients with severe renal impairment [see *Warnings and Precautions* (5.3), *Use in Specific Populations* (8.7)].

## 2.13 Dose Adjustment for Hepatotoxicity and Non-Hematologic Adverse Reactions

If elevations in bilirubin greater than 3 times the institutional upper limit of normal (IULN) or in liver transaminases greater than 5 times the IULN occur, imatinib mesylate tablets should be withheld until bilirubin levels have returned to a less than 1.5 times the IULN and transaminase levels to less than 2.5 times the IULN. In adults, treatment with imatinib mesylate tablets may then be continued at a reduced daily dose (i.e., 400 mg to 300 mg, 600 mg to 400 mg, or 800 mg to 600 mg (imatinib as free base). In children, daily doses can be reduced under the same circumstances from 340 mg/m<sup>2</sup>/day to 260 mg/m<sup>2</sup>/day (imatinib as free base).

If a severe non-hematologic adverse reaction develops (such as severe hepatotoxicity or severe fluid retention), imatinib mesylate tablets should be withheld until the event has resolved. Thereafter, treatment can be resumed as appropriate depending on the initial severity of the event.

## 2.14 Dose Adjustment for Hematologic Adverse Reactions

Dose reduction or treatment interruptions for severe neutropenia and thrombocytopenia are recommended as indicated in Table 1.

**Table 1: Dose Adjustments for Neutropenia and Thrombocytopenia**

|                                                                                                      |                                                                                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| ASM associated with eosinophilia (starting dose 100 mg)                                              | ANC <sup>1</sup> less than $1 \times 10^9/L$ and/or platelets less than $50 \times 10^9/L$ | 1. Stop imatinib mesylate until ANC greater than or equal to $1.5 \times 10^9/L$ and platelets greater than or equal to $75 \times 10^9/L$<br>2. Resume treatment with imatinib mesylate at previous dose (i.e., dose before severe adverse reaction)                                                                                                                                                                                                                            |
| HES/CEL with FIP1L1-PDGFR $\alpha$ fusion kinase (starting dose 100 mg)                              | ANC less than $1 \times 10^9/L$ and/or platelets less than $50 \times 10^9/L$              | 1. Stop imatinib mesylate until ANC greater than or equal to $1.5 \times 10^9/L$ and platelets greater than or equal to $75 \times 10^9/L$<br>2. Resume treatment with imatinib mesylate at previous dose (i.e., dose before severe adverse reaction)                                                                                                                                                                                                                            |
| Chronic Phase CML (starting dose 400 mg)                                                             | ANC less than $1 \times 10^9/L$ and/or platelets less than $50 \times 10^9/L$              | 1. Stop imatinib mesylate until ANC greater than or equal to $1.5 \times 10^9/L$ and platelets greater than or equal to $75 \times 10^9/L$                                                                                                                                                                                                                                                                                                                                       |
| MDS/MPD, ASM and HES/CEL (starting dose 400 mg)<br>GIST (starting dose 400 mg)                       |                                                                                            | 2. Resume treatment with imatinib mesylate at the original starting dose of 400 mg<br>3. If recurrence of ANC less than $1 \times 10^9/L$ and/or platelets less than $50 \times 10^9/L$ , repeat step 1 and resume imatinib mesylate at a reduced dose of 300 mg                                                                                                                                                                                                                 |
| Ph+ CML: Accelerated Phase and Blast Crisis (starting dose 600 mg)<br>Ph+ ALL (starting dose 600 mg) | ANC less than $0.5 \times 10^9/L$ and/or platelets less than $10 \times 10^9/L$            | 1. Check if cytopenia is related to leukemia (marrow aspirate or biopsy)<br>2. If cytopenia is unrelated to leukemia, reduce dose of imatinib mesylate to 400 mg<br>3. If cytopenia persists 2 weeks, reduce further to 300 mg<br>4. If cytopenia persists 4 weeks and is still unrelated to leukemia, stop imatinib mesylate until ANC greater than or equal to $1 \times 10^9/L$ and platelets greater than or equal to $20 \times 10^9/L$ and then resume treatment at 300 mg |
| DFSP (starting dose 800 mg)                                                                          | ANC less than $1 \times 10^9/L$ and/or platelets less than $50 \times 10^9/L$              | 1. Stop imatinib mesylate until ANC greater than or equal to $1.5 \times 10^9/L$ and platelets greater than or equal to $75 \times 10^9/L$ .<br>2. Resume treatment with imatinib mesylate at 600 mg<br>3. In the event of recurrence of ANC less than $1 \times 10^9/L$ and/or platelets less than $50 \times 10^9/L$ , repeat step 1 and resume imatinib mesylate at reduced dose of 400 mg.                                                                                   |
| Pediatric newly diagnosed chronic phase CML (starting dose 340 mg/m <sup>2</sup> )                   | ANC less than $1 \times 10^9/L$ and/or platelets less than $50 \times 10^9/L$              | 1. Stop imatinib mesylate until ANC greater than or equal to $1.5 \times 10^9/L$ and platelets greater than or equal to $75 \times 10^9/L$ .<br>2. Resume treatment with imatinib mesylate at previous dose (i.e., dose before severe adverse reaction)<br>3. In the event of recurrence of ANC less than $1 \times 10^9/L$ and/or platelets less than $50 \times 10^9/L$ , repeat step 1 and resume imatinib mesylate at reduced                                                |

Abbreviations: ANC, absolute neutrophil count; ASM, aggressive systemic mastocytosis; CEL, chronic eosinophilic leukemia; CML, chronic myeloid leukemia; DFSP, dermatofibrosarcoma protuberans; HES, hypereosinophilic syndrome; MDS/MPD, myelodysplastic/myeloproliferative diseases; PDGFR, platelet-derived growth factor receptor; Ph+ CML, Philadelphia chromosome positive chronic myeloid leukemia; Ph+ ALL, Philadelphia chromosome positive acute lymphoblastic leukemia.

### 3 DOSAGE FORMS AND STRENGTHS

100 mg film coated tablets

Brownish orange, slightly biconvex, round film-coated tablets with functional scoring, engraved "IMA" over score "100" on one side, "APO" on the other side

400 mg film coated tablets

Brownish orange, capsule shaped, biconvex film-coated tablets with functional scoring, engraved "IMA" score "400" on one side, "APO" on the other side.

### 4 CONTRAINDICATIONS

None.

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Fluid Retention and Edema

Imatinib mesylate is often associated with edema and occasionally serious fluid retention [see *Adverse Reactions* (6.1)]. Weigh and monitor patients regularly for signs and symptoms of fluid retention. Investigate unexpected rapid weight gain carefully and provide appropriate treatment. The probability of edema was increased with higher imatinib mesylate dose and age greater than 65 years in the CML studies. Severe superficial edema was reported in 1.5% of newly diagnosed CML patients taking imatinib mesylate, and in 2% to 6% of other adult CML patients taking imatinib mesylate. In addition, other severe fluid retention (e.g., pleural effusion, pericardial effusion, pulmonary edema, and ascites) reactions were reported in 1.3% of newly diagnosed CML patients taking imatinib mesylate, and in 2% to 6% of other adult CML patients taking imatinib mesylate. Severe fluid retention was reported in 9% to 13.1% of patients taking imatinib mesylate for GIST [see *Adverse Reactions* (6.1)].

In a randomized trial in patients with newly diagnosed Ph+CML in chronic phase comparing imatinib mesylate and nilotinib, severe (Grade 3 or 4) fluid retention occurred in 2.5% of patients receiving imatinib mesylate and in 3.9% of patients receiving nilotinib 300 mg twice daily. Effusions (including pleural effusion, pericardial effusion, ascites) or pulmonary edema were observed in 2.1% (none were Grade 3 or 4) of patients in the imatinib mesylate arm and 2.2% (0.7% Grade 3 or 4) of patients in the nilotinib 300 mg twice daily arm.

#### 5.2 Hematologic Toxicity

Treatment with imatinib mesylate tablets is associated with anemia, neutropenia, and thrombocytopenia. Perform complete blood counts weekly for the first month, biweekly for the second month, and periodically thereafter as clinically indicated (for example, every 2 to 3 months). In CML, the occurrence of these cytopenias is dependent on the stage of disease and is more frequent in patients with accelerated phase CML or blast crisis than in patients with chronic phase CML. In pediatric CML patients the most frequent toxicities observed were Grade 3 or 4 cytopenias, including neutropenia, thrombocytopenia, and anemia. These generally occur within the first several months of therapy [see *Dosage and Administration* (2.14)].

#### 5.3 Congestive Heart Failure and Left Ventricular Dysfunction

Congestive heart failure and left ventricular dysfunction have been reported in patients taking imatinib mesylate. Cardiac adverse reactions were more frequent in patients with advanced age or co-morbidities, including previous medical history of cardiac disease. In an international randomized Phase 3 study in 1106 patients with newly diagnosed Ph+ CML in chronic phase, severe cardiac failure and left ventricular dysfunction were observed in 0.7% of patients taking imatinib mesylate compared to 0.9% of patients taking IFN + Ara-C. In another randomized trial with newly diagnosed Ph+ CML patients in chronic phase that compared imatinib mesylate and nilotinib, cardiac failure was observed in 1.1% of patients in the imatinib mesylate arm and 2.2% of patients in the nilotinib 300 mg twice daily arm and severe (Grade 3 or 4) cardiac failure occurred in

0.7% of patients in each group. Carefully monitor patients with cardiac disease or risk factors for cardiac or history of renal failure. Evaluate and treat any patient with signs or symptoms consistent with cardiac or renal failure.

## **5.4 Hepatotoxicity**

Hepatotoxicity, occasionally severe, may occur with imatinib mesylate [see *Adverse Reactions* (6.1)]. Cases of fatal liver failure and severe liver injury requiring liver transplants have been reported with both short-term and long-term use of imatinib mesylate. Monitor liver function (transaminases, bilirubin, and alkaline phosphatase) before initiation of treatment and monthly, or as clinically indicated. Manage laboratory abnormalities with imatinib mesylate interruption and/or dose reduction [see *Dosage and Administration* (2.13)]. When imatinib mesylate is combined with chemotherapy, liver toxicity in the form of transaminase elevation and hyperbilirubinemia has been observed. Additionally, there have been reports of acute liver failure. Monitoring of hepatic function is recommended.

## **5.5 Hemorrhage**

In a trial of imatinib mesylate versus IFN+ Ara-C in patients with the newly diagnosed CML, 1.8% of patients had Grade 3/4 hemorrhage. In the Phase 3 unresectable or metastatic GIST studies, 211 patients (12.9%) reported Grade 3/4 hemorrhage at any site. In the Phase 2 unresectable or metastatic GIST study, 7 patients (5%) had a total of 8 CTC Grade 3/4 hemorrhages; gastrointestinal (GI) (3 patients), intra-tumoral (3 patients) or both (1 patient). Gastrointestinal tumor sites may have been the source of GI hemorrhages. In a randomized trial in patients with newly diagnosed Ph+ CML in chronic phase comparing imatinib mesylate and nilotinib, GI hemorrhage occurred in 1.4% of patients in the imatinib mesylate arm, and in 2.9% of patients in the nilotinib 300 mg twice daily arm. None of these events were Grade 3 or 4 in the imatinib mesylate arm; 0.7% were Grade 3 or 4 in the nilotinib 300 mg twice daily arm. In addition, gastric antral vascular ectasia has been reported in postmarketing experience.

## **5.6 Gastrointestinal Disorders**

Imatinib mesylate is sometimes associated with GI irritation. Imatinib mesylate tablets should be taken with food and a large glass of water to minimize this problem. There have been rare reports, including fatalities, of GI perforation.

## **5.7 Hypereosinophilic Cardiac Toxicity**

In patients with hypereosinophilic syndrome with occult infiltration of HES cells within the myocardium, cases of cardiogenic shock/left ventricular dysfunction have been associated with HES cell degranulation upon the initiation of imatinib mesylate therapy. The condition was reported to be reversible with the administration of systemic steroids, circulatory support measures and temporarily withholding imatinib mesylate.

Myelodysplastic/myeloproliferative disease and systemic mastocytosis may be associated with high eosinophil levels. Consider performing an echocardiogram and determining serum troponin in patients with HES/CEL, and in patients with MDS/MPD or ASM associated with high eosinophil levels. If either is abnormal, consider prophylactic use of systemic steroids (1 to 2 mg/kg) for one to two weeks concomitantly with imatinib mesylate at the initiation of therapy.

## **5.8 Dermatologic Toxicities**

Bullous dermatologic reactions, including erythema multiforme and Stevens-Johnson syndrome, have been reported with use of imatinib mesylate. In some cases of bullous dermatologic reactions, including erythema multiforme and Stevens-Johnson syndrome reported during postmarketing surveillance, a recurrent dermatologic reaction was observed upon rechallenge. Several foreign postmarketing reports have described cases in which patients tolerated the reintroduction of imatinib mesylate therapy after resolution or improvement of the bullous reaction. In these instances, imatinib mesylate was resumed at a dose lower than that at which the reaction occurred and some patients also received concomitant treatment with corticosteroids or antihistamines.

## **5.9 Hypothyroidism**

Clinical cases of hypothyroidism have been reported in thyroidectomy patients undergoing levothyroxine replacement during treatment with imatinib mesylate tablets. Monitor TSH levels in such patients.

## **5.10 Embryo-Fetal Toxicity**

Imatinib mesylate can cause fetal harm when administered to a pregnant woman. Imatinib mesylate was teratogenic in rats when administered during organogenesis at doses approximately equal to the maximum human dose of 800 mg/day based on body surface area (BSA). Significant post-implantation loss was seen in female rats administered imatinib mesylate at doses approximately one-half the maximum human

dose of 800 mg/day based on BSA. Advise sexually active female patients of reproductive potential to use effective contraception (methods that result in less than 1% pregnancy rates) when using imatinib mesylate tablets and for 14 days after stopping imatinib mesylate tablets. If this drug is used during pregnancy or if the patient becomes pregnant while taking this drug, apprise the patient of the potential hazard to a fetus [see *Use in Specific Populations* (8.1)].

### 5.11 Growth Retardation in Children and Adolescents

Growth retardation has been reported in children and pre-adolescents receiving imatinib mesylate tablets. The long-term effects of prolonged treatment with imatinib mesylate on growth in children are unknown. Therefore, monitor growth in children under imatinib mesylate treatment [see *Adverse Reactions* (6.1)].

### 5.12 Tumor Lysis Syndrome

Cases of Tumor Lysis Syndrome (TLS), including fatal cases, have been reported in patients with CML, GIST, ALL, and eosinophilic leukemia receiving imatinib mesylate tablets. The patients at risk of TLS are those with tumors having a high proliferative rate or high tumor burden prior to treatment. Monitor these patients closely and take appropriate precautions. Due to possible occurrence of TLS, correct clinically significant dehydration and treat high uric acid levels prior to initiation of imatinib mesylate.

### 5.13 Impairments Related to Driving and Using Machinery

Motor vehicle accidents have been reported in patients receiving imatinib mesylate tablets. Advise patients that they may experience side effects, such as dizziness, blurred vision, or somnolence during treatment with imatinib mesylate. Recommend caution when driving a car or operating machinery.

### 5.14 Renal Toxicity

A decline in renal function may occur in patients receiving imatinib mesylate tablets. Median estimated glomerular filtration rate (eGFR) values in patients on imatinib mesylate tablets 400 mg daily for newly-diagnosed CML (four randomized trials) and malignant GIST (one single-arm trial) declined from a baseline value of 85 mL/min/1.73 m<sup>2</sup> (N = 1190) to 75 mL/min/1.73 m<sup>2</sup> at 12 months (N = 1082) and 69 mL/min/1.73 m<sup>2</sup> at 60 months (N = 549). Evaluate renal function prior to initiating imatinib mesylate tablets and monitor during therapy, with attention to risk factors for renal dysfunction, such as preexisting renal impairment, diabetes mellitus, hypertension, and congestive heart failure.

## 6 ADVERSE REACTIONS

The following serious adverse reactions are described elsewhere in the labeling:

- Fluid Retention and Edema [see *Warnings and Precautions* (5.1)]
- Hematologic Toxicity [see *Warnings and Precautions* (5.2)]
- Congestive Heart Failure and Left Ventricular Dysfunction [see *Warnings and Precautions* (5.3)]
- Hepatotoxicity [see *Warnings and Precautions* (5.4)]
- Hemorrhage [see *Warnings and Precautions* (5.5)]
- Gastrointestinal Disorders [see *Warnings and Precautions* (5.6)]
- Hypereosinophilic Cardiac Toxicity [see *Warnings and Precautions* (5.7)]
- Dermatologic Toxicities [see *Warnings and Precautions* (5.8)]
- Hypothyroidism [see *Warnings and Precautions* (5.9)]
- Growth Retardation in Children and Adolescents [see *Warnings and Precautions* (5.11)]
- Tumor Lysis Syndrome [see *Warnings and Precautions* (5.12)]
- Impairments Related to Driving and Using Machinery [see *Warnings and Precautions* (5.13)]
- Renal Toxicity [see *Warnings and Precautions* (5.14)]

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

#### *Chronic Myeloid Leukemia*

The majority of imatinib mesylate-treated patients experienced adverse reactions at some time. Imatinib was discontinued due to drug-related adverse reactions in 2.4% of patients receiving imatinib mesylate in the randomized trial of newly diagnosed patients with Ph+ CML in chronic phase comparing imatinib mesylate versus IFN+ Ara-C, and in 12.5% of patients receiving imatinib mesylate in the randomized trial of newly diagnosed patients with Ph+ CML in chronic phase comparing imatinib mesylate and nilotinib.

Imatinib mesylate was discontinued due to drug-related adverse reactions in 4% of patients in chronic phase after failure of interferon-alpha therapy, in 4% of patients in accelerated phase and in 5% of patients in blast crisis.

The most frequently reported drug-related adverse reactions were edema, nausea and vomiting, muscle cramps, musculoskeletal pain, diarrhea and rash (Table 2 and Table 3 for newly diagnosed CML, Table 4 for other CML patients). Edema was most frequently periorbital or in lower limbs and was managed with diuretics, other supportive measures, or by reducing the dose of Imatinib [see *Dosage and Administration* (2.13)]. The frequency of severe superficial edema was 1.5% to 6%.

A variety of adverse reactions represent local or general fluid retention, including pleural effusion, ascites, pulmonary edema, and rapid weight gain with or without superficial edema. These reactions appear to be dose related, were more common in the blast crisis and accelerated phase studies (where the dose was 600 mg/day), and are more common in the elderly. These reactions were usually managed by interrupting imatinib mesylate treatment and using diuretics or other appropriate supportive care measures. These reactions may be serious or life threatening.

Adverse reactions, regardless of relationship to study drug, that were reported in at least 10% of the imatinib mesylate-treated patients are shown in Tables 2, 3, and 4.

**Table 2: Adverse Reactions Regardless of Relationship to Study Drug Reported in Newly Diagnosed CML Clinical Trial in the Imatinib Mesylate versus IFN+Ara-C Study (Greater Than or Equal to 10% of Imatinib Mesylate-Treated Patients)<sup>(1)</sup>**

| Preferred term                                 | All Grades<br>Imatinib<br>mesylate | IFN+Ara-C   | CTC* Grades 3/4      |             |
|------------------------------------------------|------------------------------------|-------------|----------------------|-------------|
|                                                | N = 551 (%)                        |             | Imatinib<br>mesylate | IFN+Ara-C   |
|                                                | N = 551 (%)                        | N = 533 (%) | N = 551 (%)          | N = 533 (%) |
| Fluid retention                                | 61.7                               | 11.1        | 2.5                  | 0.9         |
| – Superficial edema                            | 59.9                               | 9.6         | 1.5                  | 0.4         |
| – Other fluid retention reactions <sup>2</sup> | 6.9                                | 1.9         | 1.3                  | 0.6         |
| Nausea                                         | 49.5                               | 61.5        | 1.3                  | 5.1         |
| Muscle cramps                                  | 49.2                               | 11.8        | 2.2                  | 0.2         |
| Musculoskeletal pain                           | 47                                 | 44.8        | 5.4                  | 8.6         |
| Diarrhea                                       | 45.4                               | 43.3        | 3.3                  | 3.2         |
| Rash and related terms                         | 40.1                               | 26.1        | 2.9                  | 2.4         |
| Fatigue                                        | 38.8                               | 67          | 1.8                  | 25.1        |
| Headache                                       | 37                                 | 43.3        | 0.5                  | 3.8         |
| Joint pain                                     | 31.4                               | 38.1        | 2.5                  | 7.7         |
| Abdominal pain                                 | 36.5                               | 25.9        | 4.2                  | 3.9         |
| Nasopharyngitis                                | 30.5                               | 8.8         | 0                    | 0.4         |
| Hemorrhage                                     | 28.9                               | 21.2        | 1.8                  | 1.7         |
| - GI hemorrhage                                | 1.6                                | 1.1         | 0.5                  | 0.2         |
| - CNS hemorrhage                               | 0.2                                | 0.4         | 0                    | 0.4         |
| Myalgia                                        | 24.1                               | 38.8        | 1.5                  | 8.3         |
| Vomiting                                       | 22.5                               | 27.8        | 2                    | 3.4         |
| Dyspepsia                                      | 18.9                               | 8.3         | 0                    | 0.8         |
| Cough                                          | 20                                 | 23.1        | 0.2                  | 0.6         |
| Pharyngolaryngeal pain                         | 18.1                               | 11.4        | 0.2                  | 0           |
| Upper respiratory tract infection              | 21.2                               | 8.4         | 0.2                  | 0.4         |
| Dizziness                                      | 19.4                               | 24.4        | 0.9                  | 3.8         |
| Pyrexia                                        | 17.8                               | 42.6        | 0.9                  | 3           |
| Weight increased                               | 15.6                               | 2.6         | 2                    | 0.4         |
| Insomnia                                       | 14.7                               | 18.6        | 0                    | 2.3         |
| Depression                                     | 14.9                               | 35.8        | 0.5                  | 13.1        |
| Influenza                                      | 13.8                               | 6.2         | 0.2                  | 0.2         |
| Bone pain                                      | 11.3                               | 15.6        | 1.6                  | 3.4         |
| Constipation                                   | 11.4                               | 14.4        | 0.7                  | 0.2         |
| Sinusitis                                      | 11.4                               | 6           | 0.2                  | 0.2         |

Abbreviations: CML, chronic myeloid leukemia; CNS, central nervous system; CTC, common terminology criteria; GI, gastrointestinal; IFN, Interferon-alpha. \*NCI Common Terminology Criteria for Adverse Events, version 3.0.

<sup>(1)</sup> All adverse reactions occurring in greater than or equal to 10% of imatinib mesylate-treated patients are listed regardless of suspected relationship to treatment.

<sup>(2)</sup> Other fluid retention reactions include pleural effusion, ascites, pulmonary edema, pericardial effusion, anasarca, edema aggravated, and fluid retention not otherwise

specified.

**Table 3: Most Frequently Reported Non-Hematologic Adverse Reactions (regardless of relationship to study drug) in Patients With Newly Diagnosed Ph+ CML-CP in the Imatinib Mesylate Versus Nilotinib Study (Greater Than or Equal to 10% in Imatinib Mesylate 400 mg Once Daily or Nilotinib 300 mg Twice Daily groups) 60-Month Analysis<sup>a</sup>**

|                                                      |                                   | Patients With Newly Diagnosed Ph+ CML-CP    |                                      |                                             |                                      |
|------------------------------------------------------|-----------------------------------|---------------------------------------------|--------------------------------------|---------------------------------------------|--------------------------------------|
|                                                      |                                   | Imatinib mesylate 400 mg once-daily N = 280 | Nilotinib 300 mg twice-daily N = 279 | Imatinib mesylate 400 mg once-daily N = 280 | Nilotinib 300 mg twice-daily N = 279 |
| Body system and preferred term                       |                                   | All Grades (%)                              |                                      | CTC Grades <sup>b</sup> 3/4 (%)             |                                      |
| Skin and subcutaneous tissue disorders               | Rash                              | 19                                          | 38                                   | 2                                           | <1                                   |
|                                                      | Pruritus                          | 7                                           | 21                                   | 0                                           | <1                                   |
|                                                      | Alopecia                          | 7                                           | 13                                   | 0                                           | 0                                    |
|                                                      | Dry skin                          | 6                                           | 12                                   | 0                                           | 0                                    |
| Gastrointestinal disorders                           | Nausea                            | 41                                          | 22                                   | 2                                           | 2                                    |
|                                                      | Constipation                      | 8                                           | 20                                   | 0                                           | <1                                   |
|                                                      | Diarrhea                          | 46                                          | 19                                   | 4                                           | 1                                    |
|                                                      | Vomiting                          | 27                                          | 15                                   | <1                                          | <1                                   |
|                                                      | Abdominal pain upper              | 14                                          | 18                                   | <1                                          | 1                                    |
|                                                      | Abdominal pain                    | 12                                          | 15                                   | 0                                           | 2                                    |
|                                                      | Dyspepsia                         | 12                                          | 10                                   | 0                                           | 0                                    |
| Nervous system disorders                             | Headache                          | 23                                          | 32                                   | <1                                          | 3                                    |
|                                                      | Dizziness                         | 11                                          | 12                                   | <1                                          | <1                                   |
| General disorders and administration-site conditions | Fatigue                           | 20                                          | 23                                   | 1                                           | 1                                    |
|                                                      | Pyrexia                           | 13                                          | 14                                   | 0                                           | <1                                   |
|                                                      | Asthenia                          | 12                                          | 14                                   | 0                                           | <1                                   |
|                                                      | Peripheral edema                  | 20                                          | 9                                    | 0                                           | <1                                   |
|                                                      | Face edema                        | 14                                          | <1                                   | <1                                          | 0                                    |
| Musculoskeletal and connective tissue disorders      | Myalgia                           | 19                                          | 19                                   | <1                                          | <1                                   |
|                                                      | Arthralgia                        | 17                                          | 22                                   | <1                                          | <1                                   |
|                                                      | Muscle spasms                     | 34                                          | 12                                   | 1                                           | 0                                    |
|                                                      | Pain in extremity                 | 16                                          | 15                                   | <1                                          | <1                                   |
|                                                      | Back pain                         | 17                                          | 19                                   | 1                                           | 1                                    |
| Respiratory, thoracic and mediastinal disorders      | Cough                             | 13                                          | 17                                   | 0                                           | 0                                    |
|                                                      | Oropharyngeal pain                | 6                                           | 12                                   | 0                                           | 0                                    |
|                                                      | Dyspnea                           | 6                                           | 11                                   | <1                                          | 2                                    |
| Infections and infestations                          | Nasopharyngitis                   | 21                                          | 27                                   | 0                                           | 0                                    |
|                                                      | Upper respiratory tract infection | 14                                          | 17                                   | 0                                           | <1                                   |
|                                                      | Influenza                         | 9                                           | 13                                   | 0                                           | 0                                    |
|                                                      | Gastroenteritis                   | 10                                          | 7                                    | <1                                          | 0                                    |
| Eye disorders                                        | Eyelid edema                      | 19                                          | 1                                    | <1                                          | 0                                    |
|                                                      | Periorbital edema                 | 15                                          | <1                                   | 0                                           | 0                                    |

|                       |              |   |    |    |   |
|-----------------------|--------------|---|----|----|---|
| Psychiatric disorders | Insomnia     | 9 | 11 | 0  | 0 |
| Vascular disorder     | Hypertension | 4 | 10 | <1 | 1 |

Abbreviation: Ph+ CML-CP, Philadelphia chromosome positive chronic myeloid leukemia-chronic phase.

<sup>a</sup>Excluding laboratory abnormalities.

<sup>b</sup>NCI Common Terminology Criteria for Adverse Events, version 3.0.

**Table 4: Adverse Reactions Regardless of Relationship to Study Drug Reported in Other CML Clinical Trials (Greater Than or Equal to 10% of All Patients in Any Trial)<sup>(1)</sup>**

|                                                 | <b>Myeloid blast Crisis</b> |                  | <b>Accelerated phase</b> |                  | <b>Chronic phase, IFN</b> |                  |
|-------------------------------------------------|-----------------------------|------------------|--------------------------|------------------|---------------------------|------------------|
|                                                 | <b>(n = 260)</b>            |                  | <b>(n = 235)</b>         |                  | <b>failure</b>            |                  |
|                                                 |                             |                  |                          |                  | <b>(n = 532)</b>          |                  |
|                                                 | <b>%</b>                    |                  | <b>%</b>                 |                  | <b>%</b>                  |                  |
| <b>Preferred term</b>                           | <b>All Grades</b>           | <b>Grade 3/4</b> | <b>All Grades</b>        | <b>Grade 3/4</b> | <b>All Grades</b>         | <b>Grade 3/4</b> |
| Fluid retention                                 | 72                          | 11               | 76                       | 6                | 69                        | 4                |
| -Superficial edema                              | 66                          | 6                | 74                       | 3                | 67                        | 2                |
| -Other fluid retention reactions <sup>(2)</sup> | 22                          | 6                | 15                       | 4                | 7                         | 2                |
| Nausea                                          | 71                          | 5                | 73                       | 5                | 63                        | 3                |
| Muscle cramps                                   | 28                          | 1                | 47                       | 0.4              | 62                        | 2                |
| Vomiting                                        | 54                          | 4                | 58                       | 3                | 36                        | 2                |
| Diarrhea                                        | 43                          | 4                | 57                       | 5                | 48                        | 3                |
| Hemorrhage                                      | 53                          | 19               | 49                       | 11               | 30                        | 2                |
| - CNS hemorrhage                                | 9                           | 7                | 3                        | 3                | 2                         | 1                |
| - GI hemorrhage                                 | 8                           | 4                | 6                        | 5                | 2                         | 0.4              |
| Musculoskeletal pain                            | 42                          | 9                | 49                       | 9                | 38                        | 2                |
| Fatigue                                         | 30                          | 4                | 46                       | 4                | 48                        | 1                |
| Skin rash                                       | 36                          | 5                | 47                       | 5                | 47                        | 3                |
| Pyrexia                                         | 41                          | 7                | 41                       | 8                | 21                        | 2                |
| Arthralgia                                      | 25                          | 5                | 34                       | 6                | 40                        | 1                |
| Headache                                        | 27                          | 5                | 32                       | 2                | 36                        | 0.6              |
| Abdominal pain                                  | 30                          | 6                | 33                       | 4                | 32                        | 1                |
| Weight increased                                | 5                           | 1                | 17                       | 5                | 32                        | 7                |
| Cough                                           | 14                          | 0.8              | 27                       | 0.9              | 20                        | 0                |
| Dyspepsia                                       | 12                          | 0                | 22                       | 0                | 27                        | 0                |
| Myalgia                                         | 9                           | 0                | 24                       | 2                | 27                        | 0.2              |
| Nasopharyngitis                                 | 10                          | 0                | 17                       | 0                | 22                        | 0.2              |
| Asthenia                                        | 18                          | 5                | 21                       | 5                | 15                        | 0.2              |
| Dyspnea                                         | 15                          | 4                | 21                       | 7                | 12                        | 0.9              |
| Upper respiratory tract infection               | 3                           | 0                | 12                       | 0.4              | 19                        | 0                |
| Anorexia                                        | 14                          | 2                | 17                       | 2                | 7                         | 0                |
| Night sweats                                    | 13                          | 0.8              | 17                       | 1                | 14                        | 0.2              |
| Constipation                                    | 16                          | 2                | 16                       | 0.9              | 9                         | 0.4              |
| Dizziness                                       | 12                          | 0.4              | 13                       | 0                | 16                        | 0.2              |
| Pharyngitis                                     | 10                          | 0                | 12                       | 0                | 15                        | 0                |
| Insomnia                                        | 10                          | 0                | 14                       | 0                | 14                        | 0.2              |
| Pruritus                                        | 8                           | 1                | 14                       | 0.9              | 14                        | 0.8              |
| Hypokalemia                                     | 13                          | 4                | 9                        | 2                | 6                         | 0.8              |
| Pneumonia                                       | 13                          | 7                | 10                       | 7                | 4                         | 1                |
| Anxiety                                         | 8                           | 0.8              | 12                       | 0                | 8                         | 0.4              |
| Liver toxicity                                  | 10                          | 5                | 12                       | 6                | 6                         | 3                |
| Rigors                                          | 10                          | 0                | 12                       | 0.4              | 10                        | 0                |
| Chest pain                                      | 7                           | 2                | 10                       | 0.4              | 11                        | 0.8              |
| Influenza                                       | 0.8                         | 0.4              | 6                        | 0                | 11                        | 0.2              |
| Sinusitis                                       | 4                           | 0.4              | 11                       | 0.4              | 9                         | 0.4              |

Abbreviations: CML, chronic myeloid leukemia; IFN, Interferon-alpha.

<sup>(1)</sup>All adverse reactions occurring in greater than or equal to 10% of patients are listed

regardless of suspected relationship to treatment.

(2) Other fluid retention reactions include pleural effusion, ascites, pulmonary edema, pericardial effusion, anasarca, edema aggravated, and fluid retention not otherwise specified.

#### Hematologic and Biochemistry Laboratory Abnormalities

Cytopenias, and particularly neutropenia and thrombocytopenia, were a consistent finding in all studies, with a higher frequency at doses greater than or equal to 750 mg (Phase 1 study). The occurrence of cytopenias in CML patients was also dependent on the stage of the disease.

In patients with newly diagnosed CML, cytopenias were less frequent than in the other CML patients (see Tables 5, 6, and 7). The frequency of Grade 3 or 4 neutropenia and thrombocytopenia was between 2- and 3-fold higher in blast crisis and accelerated phase compared to chronic phase (see Tables 4 and 5). The median duration of the neutropenic and thrombocytopenic episodes varied from 2 to 3 weeks, and from 2 to 4 weeks, respectively.

These reactions can usually be managed with either a reduction of the dose or an interruption of treatment with imatinib mesylate but may require permanent discontinuation of treatment.

**Table 5: Laboratory Abnormalities in Newly Diagnosed CML Clinical Trial (Imatinib Mesylate Versus IFN+Ara-C)**

| CTC Grades                       | Imatinib Mesylate<br>N = 551 % |         | IFN+Ara-C<br>N = 533 % |         |
|----------------------------------|--------------------------------|---------|------------------------|---------|
|                                  | Grade 3                        | Grade 4 | Grade 3                | Grade 4 |
| <b>Hematology parameters*</b>    |                                |         |                        |         |
| – Neutropenia*                   | 13.1                           | 3.6     | 20.8                   | 4.5     |
| – Thrombocytopenia*              | 8.5                            | 0.4     | 15.9                   | 0.6     |
| – Anemia                         | 3.3                            | 1.1     | 4.1                    | 0.2     |
| <b>Biochemistry parameters</b>   |                                |         |                        |         |
| – Elevated creatinine            | 0                              | 0       | 0.4                    | 0       |
| – Elevated bilirubin             | 0.9                            | 0.2     | 0.2                    | 0       |
| – Elevated alkaline phosphatase  | 0.2                            | 0       | 0.8                    | 0       |
| – Elevated SGOT (AST)/SGPT (ALT) | 4.7                            | 0.5     | 7.1                    | 0.4     |

Abbreviations: CML, chronic myeloid leukemia; IFN, Interferon-alpha; SGOT, serum glutamic-oxaloacetic transaminase is now referred to as aspartate aminotransferase (AST); SGPT, serum glutamic-pyruvic transaminase is now referred to as alanine aminotransferase (ALT).

\*p less than 0.001 (difference in Grade 3 plus 4 abnormalities between the two treatment groups).

**Table 6: Percent Incidence of Clinically Relevant Grade 3/4\* Laboratory Abnormalities in the Newly Diagnosed CML Clinical Trial (Imatinib Mesylate Versus Nilotinib).**

|                                | Imatinib Mesylate<br>400 mg<br>once-daily<br>N = 280<br>(%) | Nilotinib<br>300 mg<br>twice-daily<br>N = 279<br>(%) |
|--------------------------------|-------------------------------------------------------------|------------------------------------------------------|
| <b>Hematologic parameters</b>  |                                                             |                                                      |
| Thrombocytopenia               | 9                                                           | 10                                                   |
| Neutropenia                    | 22                                                          | 12                                                   |
| Anemia                         | 6                                                           | 4                                                    |
| <b>Biochemistry parameters</b> |                                                             |                                                      |
| Elevated lipase                | 4                                                           | 9                                                    |
| Hyperglycemia                  | <1                                                          | 7                                                    |
| Hypophosphatemia               | 10                                                          | 8                                                    |
| Elevated bilirubin (total)     | <1                                                          | 4                                                    |
| Elevated SGPT (ALT)            | 3                                                           | 4                                                    |
| Hyperkalemia                   | 1                                                           | 2                                                    |
| Hyponatremia                   | <1                                                          | 1                                                    |
| Hypokalemia                    | 2                                                           | <1                                                   |
| Elevated SGOT (AST)            | 1                                                           | 1                                                    |
| Decreased albumin              | <1                                                          | 0                                                    |
| Hypocalcemia                   | <1                                                          | <1                                                   |
| Elevated alkaline phosphatase  | <1                                                          | 0                                                    |
| Elevated creatinine            | <1                                                          | 0                                                    |

Abbreviations: CML, chronic myeloid leukemia; SGOT, serum glutamic-oxaloacetic transaminase is now referred to as aspartate aminotransferase (AST); SGPT, serum glutamic-pyruvic transaminase is now referred to as alanine aminotransferase (ALT).

\*NCI Common Terminology Criteria for Adverse Events, version 3.0.

**Table 7: Laboratory Abnormalities in Other CML Clinical Trials**

|                                 | <b>Myeloid blast crisis</b>                                                   |                | <b>Accelerated phase</b>                                                      |                | <b>Chronic phase, IFN failure</b>             |                |
|---------------------------------|-------------------------------------------------------------------------------|----------------|-------------------------------------------------------------------------------|----------------|-----------------------------------------------|----------------|
|                                 | <b>(n = 260)</b><br><b>600 mg n = 223</b><br><b>400 mg n = 37</b><br><b>%</b> |                | <b>(n = 235)</b><br><b>600 mg n = 158</b><br><b>400 mg n = 77</b><br><b>%</b> |                | <b>(n = 532)</b><br><b>400 mg</b><br><b>%</b> |                |
| <b>CTC Grades<sup>1</sup></b>   | <b>Grade 3</b>                                                                | <b>Grade 4</b> | <b>Grade 3</b>                                                                | <b>Grade 4</b> | <b>Grade 3</b>                                | <b>Grade 4</b> |
| <b>Hematology parameters</b>    |                                                                               |                |                                                                               |                |                                               |                |
| – Neutropenia                   | 16                                                                            | 48             | 23                                                                            | 36             | 27                                            | 9              |
| – Thrombocytopenia              | 30                                                                            | 33             | 31                                                                            | 13             | 21                                            | <1             |
| – Anemia                        | 42                                                                            | 11             | 34                                                                            | 7              | 6                                             | 1              |
| <b>Biochemistry parameters</b>  |                                                                               |                |                                                                               |                |                                               |                |
| – Elevated creatinine           | 1.5                                                                           | 0              | 1.3                                                                           | 0              | 0.2                                           | 0              |
| – Elevated bilirubin            | 3.8                                                                           | 0              | 2.1                                                                           | 0              | 0.6                                           | 0              |
| – Elevated alkaline phosphatase | 4.6                                                                           | 0              | 5.5                                                                           | 0.4            | 0.2                                           | 0              |
| – Elevated SGOT (AST)           | 1.9                                                                           | 0              | 3.0                                                                           | 0              | 2.3                                           | 0              |
| – Elevated SGPT (ALT)           | 2.3                                                                           | 0.4            | 4.3                                                                           | 0              | 2.1                                           | 0              |

Abbreviations: CML, chronic myeloid leukemia; CTC, common terminology criteria; IFN, Interferon-alpha; SGOT, serum glutamic-oxaloacetic transaminase is now referred to as aspartate aminotransferase (AST); SGPT, serum glutamic-pyruvic transaminase is now referred to as alanine aminotransferase (ALT).

<sup>1</sup>CTC Grades: neutropenia (Grade 3 greater than or equal to  $0.5 \times 10^9/L$ , Grade 4 less than  $0.5 \times 10^9/L$ ), thrombocytopenia (Grade 3 greater than or equal to  $10 \times 10^9/L$ , Grade 4 less than  $10 \times 10^9/L$ ), anemia (hemoglobin greater than or equal to 65 to 80 g/L, Grade 4 less than 65 g/L), elevated creatinine (Grade 3 greater than 3 to 6 x upper limit normal range [ULN], Grade 4 greater than 6 x ULN), elevated bilirubin (Grade 3 greater than 3 to 10 x ULN, Grade 4 greater than 10 x ULN), elevated alkaline phosphatase (Grade 3 greater than 5 to 20 x ULN, Grade 4 greater than 20 x ULN), elevated SGOT or SGPT (Grade 3 greater than 5 to 20 x ULN, Grade 4 greater than 20 x ULN).

#### *Hepatotoxicity*

Severe elevation of transaminases or bilirubin occurred in approximately 5% of CML patients (see Tables 6 and 7) and were usually managed with dose reduction or interruption (the median duration of these episodes was approximately 1 week). Treatment was discontinued permanently because of liver laboratory abnormalities in less than 1.0% of CML patients. One patient, who was taking acetaminophen regularly for fever, died of acute liver failure. In the Phase 2 GIST trial, Grade 3 or 4 SGPT (ALT) elevations were observed in 6.8% of patients and Grade 3 or 4 SGOT (AST) elevations were observed in 4.8% of patients. Bilirubin elevation was observed in 2.7% of patients.

#### *Adverse Reactions in Pediatric Population*

##### Single-Agent Therapy

The overall safety profile of pediatric patients treated with imatinib mesylate in 93 children studied was similar to that found in studies with adult patients, except that musculoskeletal pain was less frequent (20.5%) and peripheral edema was not reported. Nausea and vomiting were the most commonly reported individual adverse reactions with an incidence similar to that seen in adult patients. Most patients experienced adverse reactions at some time during the study. The incidence of Grade 3/4 events across all types of adverse reactions was 75%; the events with the highest Grade 3/4 incidence in CML pediatric patients were mainly related to myelosuppression.

##### In Combination with Multi-Agent Chemotherapy

Pediatric and young adult patients with very high risk ALL, defined as those with an expected 5 year event-free survival (EFS) less than 45%, were enrolled after induction

therapy on a multicenter, non-randomized cooperative group pilot protocol. The study population included patients with a median age of 10 years (1 to 21 years), 61% of whom were male, 75% were white, 7% were black, and 6% were Asian/Pacific Islander. Patients with Ph+ ALL (n = 92) were assigned to receive imatinib mesylate tablets and treated in 5 successive cohorts. Imatinib mesylate tablets exposure was systematically increased in successive cohorts by earlier introduction and more prolonged duration.

The safety of imatinib mesylate tablets given in combination with intensive chemotherapy was evaluated by comparing the incidence of Grade 3 and 4 adverse events, neutropenia (less than 750/mcL) and thrombocytopenia (less than 75,000/mcL) in the 92 patients with Ph+ ALL compared to 65 patients with Ph-ALL enrolled on the trial who did not receive imatinib mesylate tablets. The safety was also evaluated comparing the incidence of adverse events in cycles of therapy administered with or without imatinib mesylate tablets. The protocol included up to 18 cycles of therapy. Patients were exposed to a cumulative total of 1,425 cycles of therapy, 778 with imatinib mesylate tablets, and 647 without imatinib mesylate tablets. The adverse events that were reported with a 5% or greater incidence in patients with Ph+ ALL compared to Ph-ALL or with a 1% or greater incidence in cycles of therapy that included imatinib mesylate tablets are presented in Table 8.

**Table 8: Adverse Reactions Reported More Frequently in Patients Treated With Study Drug (Greater Than 5%) or in Cycles With Study Drug (Greater Than 1%)**

| <b>Adverse event Grade 3 and 4 adverse events</b> | <b>Per patient incidence Ph+ ALL with Imatinib Mesylate Tablets N = 92n (%)</b> | <b>Per patient incidence Ph- ALL no Imatinib Mesylate Tablets N = 65n (%)</b> | <b>Per patient per cycle incidence with Imatinib Mesylate Tablets* N = 778n (%)</b> | <b>Per patient per cycle incidence no Imatinib Mesylate Tablets** N = 647n (%)</b> |
|---------------------------------------------------|---------------------------------------------------------------------------------|-------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|------------------------------------------------------------------------------------|
| Nausea and/or vomiting                            | 15 (16)                                                                         | 6 (9)                                                                         | 28 (4)                                                                              | 8 (1)                                                                              |
| Hypokalemia                                       | 31 (34)                                                                         | 16 (25)                                                                       | 72 (9)                                                                              | 32 (5)                                                                             |
| Pneumonitis                                       | 7 (8)                                                                           | 1 (1)                                                                         | 7 (1)                                                                               | 1 (< 1)                                                                            |
| Pleural effusion                                  | 6 (7)                                                                           | 0                                                                             | 6 (1)                                                                               | 0                                                                                  |
| Abdominal pain                                    | 8 (9)                                                                           | 2 (3)                                                                         | 9 (1)                                                                               | 3 (< 1)                                                                            |
| Anorexia                                          | 10 (11)                                                                         | 3 (5)                                                                         | 19 (2)                                                                              | 4 (1)                                                                              |
| Hemorrhage                                        | 11 (12)                                                                         | 4 (6)                                                                         | 17 (2)                                                                              | 8 (1)                                                                              |
| Hypoxia                                           | 8 (9)                                                                           | 2 (3)                                                                         | 12 (2)                                                                              | 2 (< 1)                                                                            |
| Myalgia                                           | 5 (5)                                                                           | 0                                                                             | 4 (1)                                                                               | 1 (< 1)                                                                            |
| Stomatitis                                        | 15 (16)                                                                         | 8 (12)                                                                        | 22 (3)                                                                              | 14 (2)                                                                             |
| Diarrhea                                          | 8 (9)                                                                           | 3 (5)                                                                         | 12 (2)                                                                              | 3 (< 1)                                                                            |
| Rash/Skin disorder                                | 4 (4)                                                                           | 0                                                                             | 5 (1)                                                                               | 0                                                                                  |
| Infection                                         | 49 (53)                                                                         | 32 (49)                                                                       | 131 (17)                                                                            | 92 (14)                                                                            |
| Hepatic (transaminase and/or bilirubin)           | 52 (57)                                                                         | 38 (58)                                                                       | 172 (22)                                                                            | 113 (17)                                                                           |
| Hypotension                                       | 10 (11)                                                                         | 5 (8)                                                                         | 16 (2)                                                                              | 6 (1)                                                                              |
| <b>Myelosuppression</b>                           |                                                                                 |                                                                               |                                                                                     |                                                                                    |
| Neutropenia (< 750/mcL)                           | 92 (100)                                                                        | 63 (97)                                                                       | 556 (71)                                                                            | 218 (34)                                                                           |
| Thrombocytopenia (< 75,000/mcL)                   | 90 (92)                                                                         | 63 (97)                                                                       | 431 (55)                                                                            | 329 (51)                                                                           |

Abbreviations: Ph+ ALL, Philadelphia chromosome positive acute lymphoblastic leukemia; Ph- ALL, Philadelphia chromosome negative acute lymphoblastic leukemia. \*Defined as the frequency of adverse events (AEs) per patient per treatment cycles that included imatinib mesylate tablets (includes patients with Ph+ ALL that received cycles with imatinib mesylate tablets).

\*\*Defined as the frequency of AEs per patient per treatment cycles that did not include imatinib mesylate tablets (includes patients with Ph+ ALL that received cycles without imatinib mesylate tablets as well as all patients with Ph- ALL who did not receive imatinib mesylate tablets in any treatment cycle).

#### *Adverse Reactions in Other Subpopulations*

In older patients (greater than or equal to 65 years old), with the exception of edema, where it was more frequent, there was no evidence of an increase in the incidence or severity of adverse reactions. In women there was an increase in the frequency of neutropenia, as well as Grade 1/2 superficial edema, headache, nausea, rigors, vomiting, rash, and fatigue. No differences were seen that were related to race but the subsets

were too small for proper evaluation.

#### *Acute Lymphoblastic Leukemia*

The adverse reactions were similar for Ph+ ALL as for Ph+ CML. The most frequently reported drug-related adverse reactions reported in the Ph+ ALL studies were mild nausea and vomiting, diarrhea, myalgia, muscle cramps, and rash. Superficial edema was a common finding in all studies and were described primarily as periorbital or lower limb edemas. These edemas were reported as Grade 3/4 events in 6.3% of the patients and may be managed with diuretics, other supportive measures, or in some patients by reducing the dose of imatinib mesylate.

#### *Myelodysplastic/Myeloproliferative Diseases*

Adverse reactions, regardless of relationship to study drug, that were reported in at least 10% of the patients treated with imatinib mesylate for MDS/MPD in the Phase 2 study, are shown in Table 9.

**Table 9: Adverse Reactions Regardless of Relationship to Study Drug Reported (More Than One Patient) in MPD Patients in the Phase 2 Study (Greater Than or Equal to 10% All Patients) All Grades**

| <b>Preferred Term</b> | <b>N = 7n (%)</b> |
|-----------------------|-------------------|
| Nausea                | 4 (57.1)          |
| Diarrhea              | 3 (42.9)          |
| Anemia                | 2 (28.6)          |
| Fatigue               | 2 (28.6)          |
| Muscle cramp          | 3 (42.9)          |
| Arthralgia            | 2 (28.6)          |
| Periorbital edema     | 2 (28.6)          |

Abbreviation: MPD, myeloproliferative disease.

#### *Aggressive Systemic Mastocytosis*

All aggressive systemic mastocytosis (ASM) patients experienced at least one adverse reaction at some time. The most frequently reported adverse reactions were diarrhea, nausea, ascites, muscle cramps, dyspnea, fatigue, peripheral edema, anemia, pruritus, rash, and lower respiratory tract infection. None of the 5 patients in the Phase 2 study with ASM discontinued imatinib mesylate due to drug-related adverse reactions or abnormal laboratory values.

#### *Hypereosinophilic Syndrome and Chronic Eosinophilic Leukemia*

The safety profile in the HES/CEL patient population does not appear to be different from the safety profile of imatinib mesylate observed in other hematologic malignancy populations, such as Ph+ CML. All patients experienced at least one adverse reaction, the most common being GI, cutaneous and musculoskeletal disorders. Hematological abnormalities were also frequent, with instances of CTC Grade 3 leukopenia, neutropenia, lymphopenia, and anemia.

#### *Dermatofibrosarcoma Protuberans*

Adverse reactions, regardless of relationship to study drug, that were reported in at least 10% of the 12 patients treated with imatinib mesylate for DFSP in the Phase 2 study are shown in Table 10.

**Table 10: Adverse Reactions Regardless of Relationship to Study Drug Reported in DFSP Patients in the Phase 2 Study (Greater Than or Equal to 10% All Patients) All Grades**

| <b>Preferred term</b> | <b>N = 12n (%)</b> |
|-----------------------|--------------------|
| Nausea                | 5 (41.7)           |
| Diarrhea              | 3 (25)             |
| Vomiting              | 3 (25)             |
| Periorbital edema     | 4 (33.3)           |
| Face edema            | 2 (16.7)           |
| Rash                  | 3 (25)             |
| Fatigue               | 5 (41.7)           |
| Edema peripheral      | 4 (33.3)           |
| Pyrexia               | 2 (16.7)           |
| Eye edema             | 4 (33.3)           |
| Lacrimation increased | 3 (25)             |
| Dyspnea exertional    | 2 (16.7)           |
| Anemia                | 3 (25)             |
| Rhinitis              | 2 (16.7)           |

Abbreviation: DFSP, dermatofibrosarcoma protuberans.

Clinically relevant or severe laboratory abnormalities in the 12 patients treated with imatinib mesylate for DFSP in the Phase 2 study are presented in Table 11.

**Table 11: Laboratory Abnormalities Reported in DFSP Patients in the Phase 2 Study**

| CTC Grades <sup>1</sup>        | N = 12   |   |
|--------------------------------|----------|---|
| Grade 3%                       | Grade 4% |   |
| <b>Hematology parameters</b>   |          |   |
| - Anemia                       | 17       | 0 |
| - Thrombocytopenia             | 17       | 0 |
| - Neutropenia                  | 0        | 8 |
| <b>Biochemistry parameters</b> |          |   |
| - Elevated creatinine          | 0        | 8 |

Abbreviation: CTC, common terminology criteria. <sup>1</sup>CTC Grades: neutropenia (Grade 3 greater than or equal to  $0.5 \times 10^9/L$ , Grade 4 less than  $0.5 \times 10^9/L$ ), thrombocytopenia (Grade 3 greater than or equal to  $10 \times 10^9/L$ , Grade 4 less than  $10 \times 10^9/L$ ), anemia (Grade 3 greater than or equal to 65 to 80 g/L, Grade 4 less than 65 g/L), elevated creatinine (Grade 3 greater than 3 to 6 x upper limit normal range [ULN], Grade 4 greater than 6 x ULN).

#### *Gastrointestinal Stromal Tumors*

##### Unresectable and/or Malignant Metastatic GIST

In the Phase 3 trials, the majority of imatinib mesylate-treated patients experienced adverse reactions at some time. The most frequently reported adverse reactions were edema, fatigue, nausea, abdominal pain, diarrhea, rash, vomiting, myalgia, anemia, and anorexia. Drug was discontinued for adverse reactions in a total of 89 patients (5.4%). Superficial edema, most frequently periorbital or lower extremity edema was managed with diuretics, other supportive measures, or by reducing the dose of imatinib mesylate [see *Dosage and Administration (2.13)*]. Severe (CTC Grade 3/4) edema was observed in 182 patients (11.1%).

Adverse reactions, regardless of relationship to study drug, that were reported in at least 10% of the patients treated with imatinib mesylate are shown in Table 12.

Overall the incidence of all grades of adverse reactions and the incidence of severe adverse reactions (CTC Grade 3 and above) were similar between the two treatment arms except for edema, which was reported more frequently in the 800 mg group.

**Table 12: Number (%) of Patients With Adverse Reactions Regardless of Relationship to Study Drug Where Frequency is Greater Than or Equal to 10% in any One Group (Full Analysis Set) in the Phase 3 Unresectable and/or Malignant Metastatic GIST Clinical Trials**

|                                           | Imatinib 400 mg N=818 |              | Imatinib 800 mg N=822 |              |
|-------------------------------------------|-----------------------|--------------|-----------------------|--------------|
|                                           | All Grades            | Grades 3/4/5 | All Grades            | Grades 3/4/5 |
| <b>Reported or specified term</b>         | %                     | %            | %                     | %            |
| Edema                                     | 76.7                  | 9            | 86.1                  | 13.1         |
| Fatigue/lethargy, malaise, asthenia       | 69.3                  | 11.7         | 74.9                  | 12.2         |
| Nausea                                    | 58.1                  | 9            | 64.5                  | 7.8          |
| Abdominal pain/cramping                   | 57.2                  | 13.8         | 55.2                  | 11.8         |
| Diarrhea                                  | 56.2                  | 8.1          | 58.2                  | 8.6          |
| Rash/desquamation                         | 38.1                  | 7.6          | 49.8                  | 8.9          |
| Vomiting                                  | 37.4                  | 9.2          | 40.6                  | 7.5          |
| Myalgia                                   | 32.2                  | 5.6          | 30.2                  | 3.8          |
| Anemia                                    | 32                    | 4.9          | 34.8                  | 6.4          |
| Anorexia                                  | 31.1                  | 6.6          | 35.8                  | 4.7          |
| Other GI toxicity                         | 25.2                  | 8.1          | 28.1                  | 6.6          |
| Headache                                  | 22                    | 5.7          | 19.7                  | 3.6          |
| Other pain (excluding tumor related pain) | 20.4                  | 5.9          | 20.8                  | 5            |
| Other dermatology/skin toxicity           | 17.6                  | 5.9          | 20.1                  | 5.7          |
| Leukopenia                                | 17                    | 0.7          | 19.6                  | 1.6          |
| Other constitutional symptoms             | 16.7                  | 6.4          | 15.2                  | 4.4          |

|                                                                  |      |     |      |     |
|------------------------------------------------------------------|------|-----|------|-----|
| Cough                                                            | 16.1 | 4.5 | 14.5 | 3.2 |
| Infection (without neutropenia)                                  | 15.5 | 6.6 | 16.5 | 5.6 |
| Pruritus                                                         | 15.4 | 5.4 | 18.9 | 4.3 |
| Other neurological toxicity                                      | 15   | 6.4 | 15.2 | 4.9 |
| Constipation                                                     | 14.8 | 5.1 | 14.4 | 4.1 |
| Other renal/genitourinary toxicity                               | 14.2 | 6.5 | 13.6 | 5.2 |
| Arthralgia (joint pain)                                          | 13.6 | 4.8 | 12.3 | 3   |
| Dyspnea (shortness of breath)                                    | 13.6 | 6.8 | 14.2 | 5.6 |
| Fever in absence of neutropenia (ANC < 1.0 x 10 <sup>9</sup> /L) | 13.2 | 4.9 | 12.9 | 3.4 |
| Sweating                                                         | 12.7 | 4.6 | 8.5  | 2.8 |
| Other hemorrhage                                                 | 12.3 | 6.7 | 13.3 | 6.1 |
| Weight gain                                                      | 12   | 1   | 10.6 | 0.6 |
| Alopecia                                                         | 11.9 | 4.3 | 14.8 | 3.2 |
| Dyspepsia/heartburn                                              | 11.5 | 0.6 | 10.9 | 0.5 |
| Neutropenia/granulocytopenia                                     | 11.5 | 3.1 | 16.1 | 4.1 |
| Rigors/chills                                                    | 11   | 4.6 | 10.2 | 3   |
| Dizziness/lightheadedness                                        | 11   | 4.8 | 10   | 2.8 |
| Creatinine increase                                              | 10.8 | 0.4 | 10.1 | 0.6 |
| Flatulence                                                       | 10   | 0.2 | 10.1 | 0.1 |
| Stomatitis/pharyngitis (oral/pharyngeal mucositis)               | 9.2  | 5.4 | 10   | 4.3 |
| Lymphopenia                                                      | 6    | 0.7 | 10.1 | 1.9 |

Abbreviations: ANC, absolute neutrophil count; GI, gastrointestinal; GIST, gastrointestinal stromal tumors.

Clinically relevant or severe abnormalities of routine hematologic or biochemistry laboratory values were not reported or evaluated in the Phase 3 GIST trials. Severe abnormal laboratory values reported in the Phase 2 GIST trial are presented in Table 13.

**Table 13: Laboratory Abnormalities in the Phase 2 Unresectable and/or Malignant Metastatic GIST Trial**

|                                | 400 mg(n = 73)% |         | 600 mg(n = 74)% |         |
|--------------------------------|-----------------|---------|-----------------|---------|
| CTC Grades <sup>1</sup>        | Grade 3         | Grade 4 | Grade 3         | Grade 4 |
| <b>Hematology parameters</b>   |                 |         |                 |         |
| -Anemia                        | 3               | 0       | 8               | 1       |
| -Thrombocytopenia              | 0               | 0       | 1               | 0       |
| -Neutropenia                   | 7               | 3       | 8               | 3       |
| <b>Biochemistry parameters</b> |                 |         |                 |         |
| -Elevated creatinine           | 0               | 0       | 3               | 0       |
| -Reduced albumin               | 3               | 0       | 4               | 0       |
| -Elevated bilirubin            | 1               | 0       | 1               | 3       |
| -Elevated alkaline phosphatase | 0               | 0       | 3               | 0       |
| -Elevated SGOT (AST)           | 4               | 0       | 3               | 3       |
| -Elevated SGPT (ALT)           | 6               | 0       | 7               | 1       |

Abbreviations: CTC, common terminology criteria; GIST, gastrointestinal stromal tumors; SGOT, serum glutamic-oxaloacetic transaminase is now referred to as aspartate aminotransferase (AST); SGPT, serum glutamic-pyruvic transaminase is now referred to as alanine aminotransferase (ALT). <sup>1</sup>CTC Grades: neutropenia (Grade 3 greater than or equal to 0.5–1.0 x 10<sup>9</sup>/L, Grade 4 less than 0.5 x 10<sup>9</sup>/L), thrombocytopenia (Grade 3 greater than or equal to 10 to 50 x 10<sup>9</sup>/L, Grade 4 less than 10 x 10<sup>9</sup>/L), anemia (Grade 3 greater than or equal to 65 to 80 g/L, Grade 4 less than 65 g/L), elevated creatinine (Grade 3 greater than 3 to 6 x upper limit normal range [ULN], Grade 4 greater than 6 x ULN), elevated bilirubin (Grade 3 greater than 3 to 10 x ULN, Grade 4 greater than 10 x ULN), elevated alkaline phosphatase, SGOT or SGPT (Grade 3 greater than 5 to 20 x ULN, Grade 4 greater than 20 x ULN), albumin (Grade 3 less than 20 g/L).

#### *Adjuvant Treatment of GIST*

In Study 1, the majority of both imatinib mesylate and placebo-treated patients experienced at least one adverse reaction at some time. The most frequently reported adverse reactions were similar to those reported in other clinical studies in other patient populations and include diarrhea, fatigue, nausea, edema, decreased hemoglobin, rash, vomiting, and abdominal pain. No new adverse reactions were reported in the adjuvant GIST-treatment setting that had not been previously reported in other patient populations, including patients with unresectable and/or malignant metastatic GIST. Drug

was discontinued for adverse reactions in 57 patients (17%) and 11 patients (3%) of the imatinib mesylate and placebo-treated patients, respectively. Edema, GI disturbances (nausea, vomiting, abdominal distention, and diarrhea), fatigue, low hemoglobin, and rash were the most frequently reported adverse reactions at the time of discontinuation.

In Study 2, discontinuation of therapy due to adverse reactions occurred in 15 patients (8%) and 27 patients (14%) of the imatinib mesylate 12-month, and 36-month treatment arms, respectively. As in previous trials the most common adverse reactions were diarrhea, fatigue, nausea, edema, decreased hemoglobin, rash, vomiting, and abdominal pain.

Adverse reactions, regardless of relationship to study drug, that were reported in at least 5% of the patients treated with imatinib mesylate are shown in Table 14 (Study 1) and Table 15 (Study 2). There were no deaths attributable to imatinib mesylate treatment in either trial.

**Table 14: Adverse Reactions Regardless of Relationship to Study Drug Reported in Study 1 (Greater Than or Equal to 5% of Imatinib Mesylate-Treated Patients)<sup>(1)</sup>**

|                                      | All CTC Grades    |         | CTC Grade 3* and Above |         |
|--------------------------------------|-------------------|---------|------------------------|---------|
|                                      | Imatinib Mesylate | Placebo | Imatinib Mesylate      | Placebo |
|                                      | (n=337)           | (n=345) | (n=337)                | (n=345) |
| Preferred term                       | %                 | %       | %                      | %       |
| Diarrhea                             | 59.3              | 29.3    | 3                      | 1.4     |
| Fatigue                              | 57                | 40.9    | 2.1                    | 1.2     |
| Nausea                               | 53.1              | 27.8    | 2.4                    | 1.2     |
| Periorbital edema                    | 47.2              | 14.5    | 1.2                    | 0       |
| Hemoglobin decreased                 | 46.9              | 27      | 0.6                    | 0       |
| Peripheral edema                     | 26.7              | 14.8    | 0.3                    | 0       |
| Rash (Exfoliative)                   | 26.1              | 12.8    | 2.7                    | 0       |
| Vomiting                             | 25.5              | 13.9    | 2.4                    | 0.6     |
| Abdominal pain                       | 21.1              | 22.3    | 3                      | 1.4     |
| Headache                             | 19.3              | 20.3    | 0.6                    | 0       |
| Dyspepsia                            | 17.2              | 13      | 0.9                    | 0       |
| Anorexia                             | 16.9              | 8.7     | 0.3                    | 0       |
| Weight increased                     | 16.9              | 11.6    | 0.3                    | 0       |
| Liver enzymes (ALT) increased        | 16.6              | 13      | 2.7                    | 0       |
| Muscle spasms                        | 16.3              | 3.3     | 0                      | 0       |
| Neutrophil count decreased           | 16                | 6.1     | 3.3                    | 0.9     |
| Arthralgia                           | 15.1              | 14.5    | 0                      | 0.3     |
| White blood cell count decreased     | 14.5              | 4.3     | 0.6                    | 0.3     |
| Constipation                         | 12.8              | 17.7    | 0                      | 0.3     |
| Dizziness                            | 12.5              | 10.7    | 0                      | 0.3     |
| Liver enzymes (AST) increased        | 12.2              | 7.5     | 2.1                    | 0       |
| Myalgia                              | 12.2              | 11.6    | 0                      | 0.3     |
| Blood creatinine increased           | 11.6              | 5.8     | 0                      | 0.3     |
| Cough                                | 11                | 11.3    | 0                      | 0       |
| Pruritus                             | 11                | 7.8     | 0.9                    | 0       |
| Weight decreased                     | 10.1              | 5.2     | 0                      | 0       |
| Hyperglycemia                        | 9.8               | 11.3    | 0.6                    | 1.7     |
| Insomnia                             | 9.8               | 7.2     | 0.9                    | 0       |
| Lacrimation increased                | 9.8               | 3.8     | 0                      | 0       |
| Alopecia                             | 9.5               | 6.7     | 0                      | 0       |
| Flatulence                           | 8.9               | 9.6     | 0                      | 0       |
| Rash                                 | 8.9               | 5.2     | 0.9                    | 0       |
| Abdominal distension                 | 7.4               | 6.4     | 0.3                    | 0.3     |
| Back pain                            | 7.4               | 8.1     | 0.6                    | 0       |
| Pain in extremity                    | 7.4               | 7.2     | 0.3                    | 0       |
| Hypokalemia                          | 7.1               | 2       | 0.9                    | 0.6     |
| Depression                           | 6.8               | 6.4     | 0.9                    | 0.6     |
| Facial edema                         | 6.8               | 1.2     | 0.3                    | 0       |
| Blood alkaline phosphatase increased | 6.5               | 7.5     | 0                      | 0       |
| Dry skin                             | 6.5               | 5.2     | 0                      | 0       |

|                                   |     |     |     |   |
|-----------------------------------|-----|-----|-----|---|
| Dysgeusia                         | 6.5 | 2.9 | 0   | 0 |
| Abdominal pain upper              | 6.2 | 6.4 | 0.3 | 0 |
| Neuropathy peripheral             | 5.9 | 6.4 | 0   | 0 |
| Hypocalcemia                      | 5.6 | 1.7 | 0.3 | 0 |
| Leukopenia                        | 5   | 2.6 | 0.3 | 0 |
| Platelet count decreased          | 5   | 3.5 | 0   | 0 |
| Stomatitis                        | 5   | 1.7 | 0.6 | 0 |
| Upper respiratory tract infection | 5   | 3.5 | 0   | 0 |
| Vision blurred                    | 5   | 2.3 | 0   | 0 |

Abbreviations: CTC, common terminology criteria; GIST, gastrointestinal stromal tumors; SGOT, serum glutamic-oxaloacetic transaminase is now referred to as aspartate aminotransferase (AST); SGPT, serum glutamic-pyruvic transaminase is now referred to as alanine aminotransferase (ALT). \*NCI Common Terminology Criteria for Adverse Events, version 3.0. <sup>(1)</sup> All adverse reactions occurring in greater than or equal to 5% of patients are listed regardless of suspected relationship to treatment. A patient with multiple occurrences of an adverse reaction is counted only once in the adverse reaction category.

**Table 15: Adverse Reactions Regardless of Relationship to Study Drug by Preferred Term All Grades and 3/4 Grades (Greater Than or Equal to 5% of Imatinib Mesylate-Treated Patients) Study 2<sup>(1)</sup>**

| Preferred term                        | All CTC Grades    |                   | CTC Grades 3 and Above |                   |
|---------------------------------------|-------------------|-------------------|------------------------|-------------------|
|                                       | Imatinib Mesylate | Imatinib Mesylate | Imatinib Mesylate      | Imatinib Mesylate |
|                                       | 12 Months         | 36 Months         | 12 Months              | 36 Months         |
|                                       | (N = 194)         | (N = 198)         | (N = 194)              | (N = 198)         |
|                                       | %                 | %                 | %                      | %                 |
| Patients with at least one AE         | 99                | 100               | 20.1                   | 32.8              |
| Hemoglobin decreased                  | 72.2              | 80.3              | 0.5                    | 0.5               |
| Periorbital edema                     | 59.3              | 74.2              | 0.5                    | 1                 |
| Blood lactate dehydrogenase increased | 43.3              | 60.1              | 0                      | 0                 |
| Diarrhea                              | 43.8              | 54                | 0.5                    | 2                 |
| Nausea                                | 44.8              | 51                | 1.5                    | 0.5               |
| Muscle spasms                         | 30.9              | 49                | 0.5                    | 1                 |
| Fatigue                               | 48.5              | 48.5              | 1                      | 0.5               |
| White blood cell count decreased      | 34.5              | 47                | 2.1                    | 3                 |
| Pain                                  | 25.8              | 45.5              | 1                      | 3                 |
| Blood creatinine increased            | 30.4              | 44.4              | 0                      | 0                 |
| Peripheral edema                      | 33                | 40.9              | 0.5                    | 1                 |
| Dermatitis                            | 29.4              | 38.9              | 2.1                    | 1.5               |
| Aspartate aminotransferase increased  | 30.9              | 37.9              | 1.5                    | 3                 |
| Alanine aminotransferase increased    | 28.9              | 34.3              | 2.1                    | 3                 |
| Neutrophil count decreased            | 24.2              | 33.3              | 4.6                    | 5.1               |
| Hypoproteinemia                       | 23.7              | 31.8              | 0                      | 0                 |
| Infection                             | 13.9              | 27.8              | 1.5                    | 2.5               |
| Weight increased                      | 13.4              | 26.8              | 0                      | 0.5               |
| Pruritus                              | 12.9              | 25.8              | 0                      | 0                 |
| Flatulence                            | 19.1              | 24.7              | 1                      | 0.5               |
| Vomiting                              | 10.8              | 22.2              | 0.5                    | 1                 |
| Dyspepsia                             | 17.5              | 21.7              | 0.5                    | 1                 |
| Hypoalbuminemia                       | 11.9              | 21.2              | 0                      | 0                 |
| Edema                                 | 10.8              | 19.7              | 0                      | 0.5               |
| Abdominal distension                  | 11.9              | 19.2              | 0.5                    | 0                 |
| Headache                              | 8.2               | 18.2              | 0                      | 0                 |
| Lacrimation increased                 | 18                | 17.7              | 0                      | 0                 |
| Arthralgia                            | 8.8               | 17.2              | 0                      | 1                 |
| Blood alkaline phosphatase increased  | 10.8              | 16.7              | 0                      | 0.5               |
| Dyspnea                               | 6.2               | 16.2              | 0.5                    | 1.5               |
| Myalgia                               | 9.3               | 15.2              | 0                      | 1                 |

|                           |      |      |     |     |
|---------------------------|------|------|-----|-----|
| Platelet count decreased  | 11.3 | 14.1 | 0   | 0   |
| Blood bilirubin increased | 11.3 | 13.1 | 0   | 0   |
| Dysgeusia                 | 9.3  | 12.6 | 0   | 0   |
| Paresthesia               | 5.2  | 12.1 | 0   | 0.5 |
| Vision blurred            | 10.8 | 11.1 | 1.0 | 0.5 |
| Alopecia                  | 11.3 | 10.6 | 0   | 0   |
| Decreased appetite        | 9.8  | 10.1 | 0   | 0   |
| Constipation              | 8.8  | 9.6  | 0   | 0   |
| Pyrexia                   | 6.2  | 9.6  | 0   | 0   |
| Depression                | 3.1  | 8.1  | 0   | 0   |
| Abdominal pain            | 2.6  | 7.6  | 0   | 0   |
| Conjunctivitis            | 5.2  | 7.6  | 0   | 0   |
| Photosensitivity reaction | 3.6  | 7.1  | 0   | 0   |
| Dizziness                 | 4.6  | 6.6  | 0.5 | 0   |
| Hemorrhage                | 3.1  | 6.6  | 0   | 0   |
| Dry skin                  | 6.7  | 6.1  | 0.5 | 0   |
| Nasopharyngitis           | 1    | 6.1  | 0   | 0.5 |
| Palpitations              | 5.2  | 5.1  | 0   | 0   |

Abbreviations: AE, adverse event; CTC, common terminology criteria.

<sup>(1)</sup> All adverse reactions occurring in greater than or equal to 5% of patients are listed regardless of suspected relationship to treatment. A patient with multiple occurrences of an adverse reaction is counted only once in the adverse reaction category.

#### *Adverse Reactions from Multiple Clinical Trials*

##### Cardiac Disorders:

Estimated 1% to 10%: palpitations, pericardial effusion

Estimated 0.1% to 1%: congestive cardiac failure, tachycardia, pulmonary edema

Estimated 0.01% to 0.1%: arrhythmia, atrial fibrillation, cardiac arrest, myocardial infarction, angina pectoris

##### Vascular Disorders:

Estimated 1% to 10%: flushing, hemorrhage

Estimated 0.1% to 1%: hypertension, hypotension, peripheral coldness, Raynaud's phenomenon, hematoma, subdural hematoma

##### Investigations:

Estimated 1% to 10%: blood creatine phosphokinase (CPK) increased, blood amylase increased

Estimated 0.1% to 1%: blood lactate dehydrogenase (LDH) increased

##### Skin and Subcutaneous Tissue Disorders:

Estimated 1% to 10%: dry skin, alopecia, face edema, erythema, photosensitivity reaction, nail disorder, purpura

Estimated 0.1% to 1%: exfoliative dermatitis, bullous eruption, psoriasis, rash pustular, contusion, sweating increased, urticaria, ecchymosis, increased tendency to bruise, hypotrichosis, skin hypopigmentation, skin hyperpigmentation, onychoclasia, folliculitis, petechiae, erythema multiforme, panniculitis (including erythema nodosum)

Estimated 0.01% to 0.1%: vesicular rash, Stevens-Johnson syndrome, acute generalized exanthematous pustulosis, acute febrile neutrophilic dermatosis (Sweet's syndrome), nail discoloration, angioneurotic edema, leucocytoclastic vasculitis

##### Gastrointestinal Disorders:

Estimated 1% to 10%: abdominal distention, gastroesophageal reflux, dry mouth, gastritis

Estimated 0.1% to 1%: gastric ulcer, stomatitis, mouth ulceration, eructation, melena, esophagitis, ascites, hematemesis, chelitis, dysphagia, pancreatitis

Estimated 0.01% to 0.1%: colitis, ileus, inflammatory bowel disease

##### General Disorders and Administration-Site Conditions:

Estimated 1% to 10%: weakness, anasarca, chills

Estimated 0.1% to 1%: malaise

##### Blood and Lymphatic System Disorders:

Estimated 1% to 10%: pancytopenia, febrile neutropenia, lymphopenia, eosinophilia

Estimated 0.1% to 1%: thrombocythemia, bone marrow depression, lymphadenopathy

Estimated 0.01% to 0.1%: hemolytic anemia, aplastic anemia

#### Hepatobiliary Disorders:

Estimated 0.1% to 1%: hepatitis, jaundice

Estimated 0.01% to 0.1%: hepatic failure and hepatic necrosis<sup>1</sup>

#### Immune System Disorders:

Estimated 0.01% to 0.1%: angioedema

#### Infections and Infestations:

Estimated 0.1% to 1%: sepsis, herpes simplex, herpes zoster, cellulitis, urinary tract infection, gastroenteritis

Estimated 0.01% to 0.1%: fungal infection

#### Metabolism and Nutrition Disorders:

Estimated 1% to 10%: weight decreased, decreased appetite

Estimated 0.1% to 1%: dehydration, gout, increased appetite, hyperuricemia, hypercalcemia, hyperglycemia, hyponatremia, hyperkalemia, hypomagnesemia

#### Musculoskeletal and Connective Tissue Disorders:

Estimated 1% to 10%: joint swelling

Estimated 0.1% to 1%: joint and muscle stiffness, muscular weakness, arthritis

#### Nervous System/Psychiatric Disorders:

Estimated 1% to 10%: paresthesia, hypesthesia

Estimated 0.1% to 1%: syncope, peripheral neuropathy, somnolence, migraine, memory impairment, libido decreased, sciatica, restless leg syndrome, tremor

Estimated 0.01% to 0.1%: increased intracranial pressure<sup>1</sup>, confusional state, convulsions, optic neuritis

#### Renal and Urinary Disorders:

Estimated 0.1% to 1%: renal failure acute, urinary frequency increased, hematuria, renal pain

#### Reproductive System and Breast Disorders:

Estimated 0.1% to 1%: breast enlargement, menorrhagia, sexual dysfunction, gynecomastia, erectile dysfunction, menstruation irregular, nipple pain, scrotal edema

#### Respiratory, Thoracic and Mediastinal Disorders:

Estimated 1% to 10%: epistaxis

Estimated 0.1% to 1%: pleural effusion

Estimated 0.01% to 0.1%: interstitial pneumonitis, pulmonary fibrosis, pleuritic pain, pulmonary hypertension, pulmonary hemorrhage

#### Endocrine Disorders:

Estimated 0.1% to 1%: hypothyroidism, hyperthyroidism

#### Eye, Ear, and Labyrinth Disorders:

Estimated 1% to 10%: conjunctivitis, vision blurred, orbital edema, conjunctival hemorrhage, dry eye

Estimated 0.1% to 1%: vertigo, tinnitus, eye irritation, eye pain, scleral hemorrhage, retinal hemorrhage, blepharitis, macular edema, hearing loss, cataract

Estimated 0.01% to 0.1%: papilledema<sup>1</sup>, glaucoma

<sup>1</sup>Including some fatalities.

## **6.2 Postmarketing Experience**

The following additional adverse reactions have been identified during post approval use of imatinib mesylate tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Blood and Lymphatic System Disorders:* thrombotic microangiopathy

*Cardiac Disorders:* pericarditis, cardiac tamponade<sup>1</sup>

*Eye Disorders:* vitreous hemorrhage

*Gastrointestinal Disorders:* ileus/intestinal obstruction, tumor hemorrhage/tumor necrosis, GI perforation<sup>1</sup> [see *Warnings and Precautions* (5.6)], diverticulitis, gastric antral vascular ectasia

*Infections:* hepatitis B virus reactivation<sup>1</sup>

*Musculoskeletal and Connective Tissue Disorders:* osteonecrosis, rhabdomyolysis/myopathy, growth retardation in children, musculoskeletal pain upon treatment discontinuation (including myalgia, pain in extremity, arthralgia, bone pain)

*Nervous System Disorders:* cerebral edema<sup>1</sup>

*Reproduction Disorders:* hemorrhagic corpus luteum/hemorrhagic ovarian cyst

*Respiratory, Thoracic and Mediastinal Disorders:* acute respiratory failure<sup>1</sup>, interstitial lung disease

*Skin and Subcutaneous Tissue Disorders:* lichenoid keratosis, lichen planus, toxic epidermal necrolysis, palmar-plantar erythrodysesthesia syndrome, drug rash with eosinophilia and systemic symptoms (DRESS), pseudoporphyria, pemphigus

*Vascular Disorders:* thrombosis/embolism, anaphylactic shock

<sup>1</sup>Including some fatalities.

## **7 DRUG INTERACTIONS**

### **7.1 Agents Inducing CYP3A Metabolism**

Concomitant administration of imatinib mesylate tablets and strong CYP3A4 inducers may reduce total exposure of imatinib; consider alternative agents [see *Clinical Pharmacology* (12.3)].

### **7.2 Agents Inhibiting CYP3A Metabolism**

Concomitant administration of imatinib mesylate tablets and strong CYP3A4 inhibitors may result in a significant imatinib exposure increase. Grapefruit juice may also increase plasma concentrations of imatinib; avoid grapefruit juice [see *Clinical Pharmacology* (12.3)].

### **7.3 Interactions With Drugs Metabolized by CYP3A4**

Imatinib mesylate will increase plasma concentration of CYP3A4 metabolized drugs (e.g., triazolo-benzodiazepines, dihydropyridine calcium channel blockers, certain HMG-CoA reductase inhibitors, etc.). Use caution when administering imatinib mesylate tablets with CYP3A4 substrates that have a narrow therapeutic window.

Because warfarin is metabolized by CYP2C9 and CYP3A4, use low-molecular weight or standard heparin instead of warfarin in patients who require anticoagulation [see *Clinical Pharmacology* (12.3)].

### **7.4 Interactions With Drugs Metabolized by CYP2D6**

Use caution when administering imatinib mesylate with CYP2D6 substrates that have a narrow therapeutic window.

### **7.5 Interactions With Methotrexate**

Imatinib mesylate tablets delay the clearance of methotrexate and result in sustained methotrexate concentrations when methotrexate is used at high doses (>500 mg/m<sup>2</sup>). Refer to the prescribing information for methotrexate injection for recommendations on management of methotrexate concentrations.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### *Risk Summary*

Imatinib mesylate tablets can cause fetal harm when administered to a pregnant woman based on human and animal data. There are no clinical studies regarding use of imatinib mesylate tablets in pregnant women. There have been postmarket reports of spontaneous abortions and congenital anomalies from women who have been exposed to imatinib mesylate tablets during pregnancy. Reproductive studies in rats have demonstrated that imatinib mesylate induced teratogenicity and increased incidence of congenital abnormalities following prenatal exposure to imatinib mesylate at doses equal to the highest recommended human dose of 800 mg/day based on BSA. Advise women to avoid pregnancy when taking imatinib mesylate tablets. If this drug is used during pregnancy, or if the patient becomes pregnant while taking this drug, apprise the patient of the potential hazard to the fetus.

The background risk of major birth defects and miscarriage for the indicated population is not known; however, in the U.S. general population, the estimated background risk of major birth defects of clinically recognized pregnancies is 2% to 4% and of miscarriage is 15% to 20%.

#### *Data*

##### Animal Data

In embryo-fetal development studies in rats and rabbits, pregnant animals received oral doses of imatinib mesylate up to 100 mg/kg/day and 60 mg/kg/day, respectively, during the period of organogenesis.

In rats, imatinib mesylate was teratogenic at 100 mg/kg/day (approximately equal to the maximum human dose of 800 mg/day based on BSA), the number of fetuses with encephalocoele and exencephaly was higher than historical control values and these findings were associated with missing or underdeveloped cranial bones. Lower mean fetal body weights were associated with retarded skeletal ossifications.

In rabbits, at doses 1.5 times higher than the maximum human dose of 800 mg/day based on BSA, no effects on the reproductive parameters with respect to implantation sites, number of live fetuses, sex ratio or fetal weight were observed. The examinations of the fetuses did not reveal any drug related morphological changes.

In a pre- and postnatal development study in rats, pregnant rats received oral doses of imatinib mesylate during gestation (organogenesis) and lactation up to 45 mg/kg/day. Five animals developed a red vaginal discharge in the 45 mg/kg/day group on Days 14 or 15 of gestation, the significance of which is unknown since all females produced viable litters and none had increased post-implantation loss. Other maternal effects noted only at the dose of 45 mg/kg/day (approximately one-half the maximum human dose of 800 mg/day based on BSA) included an increased number of stillborn pups and pups dying between postpartum Days 0 and 4. In the F1 offspring at this same dose level, mean body weights were reduced from birth until terminal sacrifice and the number of litters achieving criterion for preputial separation was slightly decreased. There were no other significant effects in developmental parameters or behavioral testing. F1 fertility was not affected but reproductive effects were noted at 45 mg/kg/day, including an increased number of resorptions and a decreased number of viable fetuses. The no-observed-effect level (NOEL) for both maternal animals and the F1 generation was 15 mg/kg/day.

## **8.2 Lactation**

#### *Risk Summary*

Imatinib and its active metabolite are excreted into human milk. Because of the potential for serious adverse reactions in breastfed infants from imatinib mesylate, advise a lactating woman not to breastfeed during treatment and for 1 month after the last dose.

#### *Human Data*

Based on data from three breastfeeding women taking imatinib mesylate tablets, the milk: plasma ratio is about 0.5 for imatinib and about 0.9 for the active metabolite. Considering the combined concentration of imatinib and active metabolite, a breastfed infant could receive up to 10% of the maternal therapeutic dose based on body weight.

## **8.3 Females and Males of Reproductive Potential**

Human postmarketing reports and animal studies have shown imatinib mesylate to be harmful to the developing fetus [see *Use in Specific Populations* (8.1)]. <sup>1</sup>

#### *Pregnancy Testing*

Test pregnancy status in females with reproductive potential prior to the initiation of treatment with imatinib mesylate tablets.

#### *Contraception*

##### Females

Advise female patients of reproductive potential to use effective contraception (methods that result in less than 1% pregnancy rates) when using imatinib mesylate tablets during treatment and for fourteen days after stopping treatment with imatinib mesylate tablets [see *Use in Specific Populations* (8.1)].

#### *Infertility*

The risk of infertility in females or males of reproductive potential has not been studied in humans. In a rat study, the fertility in males and females was not affected [see *Nonclinical Toxicology* (13)].

## **8.4 Pediatric Use**

The safety and effectiveness of imatinib mesylate tablets have been demonstrated in pediatric patients with newly diagnosed Ph+ chronic phase CML and Ph+ ALL [see *Clinical Studies* (14.2, 14.4)]. There are no data in children under 1 year of age.

## 8.5 Geriatric Use

In the CML clinical studies, approximately 20% of patients were older than 65 years. In the study of patients with newly diagnosed CML, 6% of patients were older than 65 years. The frequency of edema was higher in patients older than 65 years as compared to younger patients; no other difference in the safety profile was observed [see *Warnings and Precautions* (5.1)]. The efficacy of imatinib mesylate was similar in older and younger patients.

In the unresectable or metastatic GIST study, 16% of patients were older than 65 years. No obvious differences in the safety or efficacy profile were noted in patients older than 65 years as compared to younger patients, but the small number of patients does not allow a formal analysis.

In the adjuvant GIST study, 221 patients (31%) were older than 65 years. No difference was observed in the safety profile in patients older than 65 years as compared to younger patients, with the exception of a higher frequency of edema. The efficacy of imatinib mesylate was similar in patients older than 65 years and younger patients.

## 8.6 Hepatic Impairment

The effect of hepatic impairment on the pharmacokinetics of both imatinib and its major metabolite, CGP74588, was assessed in 84 patients with cancer with varying degrees of hepatic impairment at imatinib doses ranging from 100 mg to 800 mg.

Mild and moderate hepatic impairment do not influence exposure to imatinib and CGP74588. In patients with severe hepatic impairment, the imatinib  $C_{max}$  and area under curve (AUC) increased by 63% and 45% and the CGP74588  $C_{max}$  and AUC increased by 56% and 55%, relative to patients with normal hepatic function [see *Clinical Pharmacology* (12.3)]. Reduce the dose by 25% for patients with severe hepatic impairment [see *Dosage and Administration* (2.12)].

**Table 16: Liver Function Classification**

| Liver function test    | Normal (n = 14)           | Mild (n = 30)                                                           | Moderate (n = 20)                   | Severe (n = 20)                    |
|------------------------|---------------------------|-------------------------------------------------------------------------|-------------------------------------|------------------------------------|
| <b>Total bilirubin</b> | less than or equal to ULN | greater than 1.0 to 1.5 times the ULN                                   | greater than 1.5 to 3 times the ULN | greater than 3 to 10 times the ULN |
| <b>SGOT</b>            | less than or equal to ULN | greater than ULN (can be normal if Total Bilirubin is greater than ULN) | Any                                 | Any                                |

Abbreviation: SGOT, serum glutamic-oxaloacetic transaminase is now referred to as aspartate aminotransferase (AST); ULN, upper limit of normal for the institution.

## 8.7 Renal Impairment

The effect of renal impairment on the pharmacokinetics of imatinib was assessed in 59 patients with cancer and varying degrees of renal impairment at single and steady state imatinib doses ranging from 100 to 800 mg/day. The mean exposure to imatinib (dose normalized AUC) in patients with mild and moderate renal impairment increased 1.5- to 2-fold compared to patients with normal renal function. There are not sufficient data in patients with severe renal impairment [see *Clinical Pharmacology* (12.3)]. Dose reductions are necessary for patients with moderate and severe renal impairment [see *Dosage and Administration* (2.12)].

**Table 17: Renal Function Classification**

| Renal dysfunction | Renal function tests       |
|-------------------|----------------------------|
| <b>Mild</b>       | CrCL = 40 to 59 mL/min     |
| <b>Moderate</b>   | CrCL = 20 to 39 mL/min     |
| <b>Severe</b>     | CrCL = less than 20 mL/min |

Abbreviation: CrCL, creatinine clearance.

## 10 OVERDOSAGE

Experience with doses greater than 800 mg is limited. Isolated cases of imatinib mesylate overdose have been reported. In the event of overdosage, observe the patient and give appropriate supportive treatment.

### Adult Overdose

**1,200 to 1,600 mg (duration varying between 1 to 10 days):** Nausea, vomiting, diarrhea, rash erythema, edema, swelling, fatigue, muscle spasms, thrombocytopenia, pancytopenia, abdominal pain, headache, decreased appetite.

**1,800 to 3,200 mg (as high as 3,200 mg daily for 6 days):** Weakness, myalgia, increased CPK, increased bilirubin, GI pain.

**6,400 mg (single dose):** One case in the literature reported one patient who experienced nausea, vomiting, abdominal pain, pyrexia, facial swelling, neutrophil count decreased, increase transaminases.

**8 to 10 g (single dose):** Vomiting and GI pain have been reported.


A patient with myeloid blast crisis experienced Grade 1 elevations of serum creatinine, Grade 2 ascites and elevated liver transaminase levels, and Grade 3 elevations of bilirubin after inadvertently taking 1,200 mg of imatinib mesylate (imatinib as free base) daily for 6 days. Therapy was temporarily interrupted and complete reversal of all abnormalities occurred within 1 week. Treatment was resumed at a dose of 400 mg daily (imatinib as free base) without recurrence of adverse reactions. Another patient developed severe muscle cramps after taking 1,600 mg of imatinib mesylate (imatinib as free base) daily for 6 days. Complete resolution of muscle cramps occurred following interruption of therapy and treatment was subsequently resumed. Another patient that was prescribed 400 mg daily (imatinib as free base), took 800 mg of imatinib mesylate (imatinib as free base) on Day 1 and 1,200 mg (imatinib as free base) on Day 2. Therapy was interrupted, no adverse reactions occurred and the patient resumed therapy.

#### *Pediatric Overdose*

One 3 year old male exposed to a single dose of 400 mg experienced vomiting, diarrhea, and anorexia and another 3 year old male exposed to a single dose of 980 mg experienced decreased white blood cell (WBC) count and diarrhea.

## 11 DESCRIPTION

Imatinib is a small molecule kinase inhibitor. Imatinib mesylate film-coated tablets are supplied as 100 mg and 400 mg tablets for oral administration. Each 100 mg tablet contains 119.5 mg of imatinib mesylate equivalent to 100 mg of imatinib free base. Each 400 mg tablet contains 478 mg of imatinib mesylate equivalent to 400 mg of imatinib free base. Imatinib mesylate is designated chemically as 4-[(4-Methyl-1-piperazinyl)methyl]-N-[4-methyl-3-[[4-(3-pyridinyl)-2-pyrimidinyl]amino]-phenyl]benzamide methanesulfonate and its structural formula is:



Imatinib mesylate is a white to off-white crystalline powder. Its molecular formula is  $C_{29}H_{31}N_7O \cdot CH_4SO_3$  and its molecular weight is 589.7 g/mol. Imatinib mesylate is freely soluble in water and freely to sparingly soluble in methanol.

Inactive Ingredients: colloidal silicon dioxide (NF); croscopovidone (NF); and magnesium stearate (NF). Tablet coating: ferric oxide red (NF); ferric oxide yellow (NF); hypromellose (USP); hydroxypropyl cellulose, and polyethylene glycol (NF).

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Imatinib mesylate is a protein-tyrosine kinase inhibitor that inhibits the BCR-ABL tyrosine kinase, the constitutive abnormal tyrosine kinase created by the Philadelphia chromosome abnormality in CML. Imatinib inhibits proliferation and induces apoptosis in BCR-ABL positive cell lines as well as fresh leukemic cells from Philadelphia chromosome positive chronic myeloid leukemia. Imatinib inhibits colony formation in assays using *ex vivo* peripheral blood and bone marrow samples from CML patients.

*In vivo*, imatinib inhibits tumor growth of BCR-ABL transfected murine myeloid cells as well as BCR-ABL positive leukemia lines derived from CML patients in blast crisis.

Imatinib is also an inhibitor of the receptor tyrosine kinases for platelet-derived growth factor (PDGF) and stem cell factor (SCF), c-Kit, and inhibits PDGF- and SCF-mediated cellular events. *In vitro*, imatinib inhibits proliferation and induces apoptosis in GIST cells, which express an activating c-Kit mutation.

## 12.3 Pharmacokinetics

The pharmacokinetics of imatinib mesylate have been evaluated in studies in healthy subjects and in population pharmacokinetic studies in over 900 patients. The pharmacokinetics of imatinib mesylate are similar in CML and GIST patients.

### *Absorption and Distribution*

Imatinib is well absorbed after oral administration with  $C_{max}$  achieved within 2 to 4 hours post-dose. Mean absolute bioavailability is 98%. Mean imatinib AUC increases proportionally with increasing doses ranging from 25 mg to 1,000 mg. There is no significant change in the pharmacokinetics of imatinib on repeated dosing, and accumulation is 1.5- to 2.5-fold at steady state when imatinib mesylate is dosed once daily. At clinically relevant concentrations of imatinib, binding to plasma proteins in *in vitro* experiments is approximately 95%, mostly to albumin and  $\alpha$ 1-acid glycoprotein.

### *Elimination*

#### Metabolism

CYP3A4 is the major enzyme responsible for metabolism of imatinib. Other cytochrome P450 enzymes, such as CYP1A2, CYP2D6, CYP2C9, and CYP2C19, play a minor role in its metabolism. The main circulating active metabolite in humans is the N-demethylated piperazine derivative, formed predominantly by CYP3A4. It shows *in vitro* potency similar to the parent imatinib. The plasma AUC for this metabolite is about 15% of the AUC for imatinib. The plasma protein binding of N-demethylated metabolite CGP74588 is similar to that of the parent compound.

#### Excretion

Imatinib elimination is predominately in the feces, mostly as metabolites. Based on the recovery of compound(s) after an oral  $^{14}\text{C}$ -labeled dose of imatinib, approximately 81% of the dose was eliminated within 7 days, in feces (68% of dose) and urine (13% of dose). Unchanged imatinib accounted for 25% of the dose (5% urine, 20% feces), the remainder being metabolites.

Following oral administration in healthy volunteers, the elimination half-lives of imatinib and its major active metabolite, the N-demethyl derivative (CGP74588), are approximately 18 and 40 hours, respectively.

Typically, clearance of imatinib in a 50-year-old patient weighing 50 kg is expected to be 8 L/h, while for a 50-year-old patient weighing 100 kg the clearance will increase to 14 L/h. The inter-patient variability of 40% in clearance does not warrant initial dose adjustment based on body weight and/or age but indicates the need for close monitoring for treatment-related toxicity.

### *Specific Populations*

#### Hepatic Impairment

The effect of hepatic impairment on the pharmacokinetics of both imatinib and its major metabolite, CGP74588, was assessed in 84 patients with cancer and varying degrees of hepatic impairment [see *Use in Specific Populations* (8.6)] at imatinib doses ranging from 100 mg to 800 mg. Exposure to both imatinib and CGP74588 was comparable between each of the mildly and moderately hepatically-impaired groups and the normal group. Patients with severe hepatic impairment tend to have higher exposure to both imatinib and its metabolite than patients with normal hepatic function. At steady state, the mean  $C_{max}$ /dose and AUC/dose for imatinib increased by about 63% and 45%, respectively, in patients with severe hepatic impairment compared to patients with normal hepatic function. The mean  $C_{max}$ /dose and AUC/dose for CGP74588 increased by about 56% and 55%, respectively, in patients with severe hepatic impairment compared to patients with normal hepatic function. Dose reductions are necessary for patients with severe hepatic impairment [see *Dosage and Administration* (2.12)].

#### Renal Impairment

The effect of renal impairment on the pharmacokinetics of imatinib was assessed in 59 cancer patients with varying degrees of renal impairment [see *Use in Specific Populations* (8.7)] at single and steady state imatinib doses ranging from 100 to 800 mg/day. The mean exposure to imatinib (dose normalized AUC) in patients with mild and moderate renal impairment increased 1.5- to 2-fold compared to patients with normal renal function. The AUCs did not increase for doses greater than 600 mg in patients with mild renal impairment. The AUCs did not increase for doses greater than 400 mg in patients with moderate renal impairment. Two patients with severe renal impairment were dosed with 100 mg/day and their exposures were similar to those seen in patients with normal renal function receiving 400 mg/day. Dose reductions are necessary for patients with moderate and severe renal impairment [see *Dosage and Administration* (2.12)].

#### Pediatric Use

As in adult patients, imatinib was rapidly absorbed after oral administration in pediatric patients, with a  $C_{max}$  of 2 to 4 hours. Apparent oral clearance was similar to adult values (11.0 L/hr/m<sup>2</sup> in children vs 10.0 L/hr/m<sup>2</sup> in adults), as was the half-life (14.8 hours in children vs 17.1 hours in adults). Dosing in children at both 260 mg/m<sup>2</sup> and 340 mg/m<sup>2</sup> achieved an AUC similar to the 400 mg dose in adults. The comparison of AUC on Day 8

vs Day 1 at 260 mg/m<sup>2</sup> and 340 mg/m<sup>2</sup> dose levels revealed a 1.5- and 2.2-fold drug accumulation, respectively, after repeated once-daily dosing. Mean imatinib AUC did not increase proportionally with increasing dose.

Based on pooled population pharmacokinetic analysis in pediatric patients with hematological disorders (CML, Ph+ ALL, or other hematological disorders treated with imatinib), clearance of imatinib increases with increasing BSA. After correcting for the BSA effect, other demographics, such as age, body weight, and body mass index did not have clinically significant effects on the exposure of imatinib. The analysis confirmed that exposure of imatinib in pediatric patients receiving 260 mg/m<sup>2</sup> once daily (not exceeding 400 mg once daily) or 340 mg/m<sup>2</sup> once daily (not exceeding 600 mg once daily) were similar to those in adult patients who received imatinib 400 mg or 600 mg once daily.

#### *Drug Interaction Studies*

##### Clinical Studies

##### Agents Inducing CYP3A Metabolism

Pretreatment of healthy volunteers with multiple doses of rifampin followed by a single dose of imatinib mesylate tablets, increased imatinib mesylate oral-dose clearance by 3.8-fold, which significantly (p less than 0.05) decreased mean C<sub>max</sub> and AUC.

Similar findings were observed in patients receiving 400 to 1200 mg/day imatinib mesylate tablets concomitantly with enzyme-inducing anti-epileptic drugs (EIAED) (e.g., carbamazepine, oxcarbamazepine, phenytoin, fosphenytoin, phenobarbital, and primidone). The mean dose normalized AUC for imatinib in the patients receiving EIAED's decreased by 73% compared to patients not receiving EIAED.

Concomitant administration of imatinib mesylate tablets and St. John's Wort led to a 30% reduction in the AUC of imatinib.

Consider alternative therapeutic agents with less enzyme induction potential in patients when rifampin or other CYP3A4 inducers are indicated. Imatinib mesylate tablets doses up to 1200 mg/day (600 mg twice daily) have been given to patients receiving concomitant strong CYP3A4 inducers [see *Dosage and Administration* (2.12)].

##### Agents Inhibiting CYP3A Metabolism

There was a significant increase in exposure to imatinib (mean C<sub>max</sub> and AUC increased by 26% and 40%, respectively) in healthy subjects when imatinib mesylate tablets were coadministered with a single dose of ketoconazole (a CYP3A4 inhibitor). Caution is recommended when administering imatinib mesylate tablets with strong CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, clarithromycin, atazanavir, indinavir, nefazodone, nelfinavir, ritonavir, saquinavir, telithromycin, and voriconazole). Grapefruit juice may also increase plasma concentrations of imatinib and should be avoided.

##### Interactions with Drugs Metabolized by CYP3A4

Imatinib mesylate tablets increase the mean C<sub>max</sub> and AUC of simvastatin (CYP3A4 substrate) 2- and 3.5-fold, respectively, suggesting an inhibition of the CYP3A4 by imatinib mesylate. Particular caution is recommended when administering imatinib mesylate tablets with CYP3A4 substrates that have a narrow therapeutic window (e.g., alfentanil, cyclosporine, diergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus, or tacrolimus).

Imatinib mesylate tablets will increase plasma concentration of other CYP3A4 metabolized drugs (e.g., triazolo-benzodiazepines, dihydropyridine calcium channel blockers, certain HMG-CoA reductase inhibitors, etc.).

Because warfarin is metabolized by CYP2C9 and CYP3A4, patients who require anticoagulation should receive low-molecular weight or standard heparin instead of warfarin.

##### Interactions with Drugs Metabolized by CYP2D6

Imatinib mesylate tablets increased the mean C<sub>max</sub> and AUC of metoprolol by approximately 23% suggesting that imatinib mesylate has a weak inhibitory effect on CYP2D6-mediated metabolism. No dose adjustment is necessary, however, caution is recommended when administering imatinib mesylate tablets with CYP2D6 substrates that have a narrow therapeutic window.

##### Interactions with Acetaminophen

Coadministration of imatinib mesylate tablets (400 mg/day for eight days) with acetaminophen (1,000 mg single dose on Day 8) in patients with CML did not result in any changes in the pharmacokinetics of acetaminophen. Imatinib mesylate pharmacokinetics were not altered in the presence of single-dose acetaminophen. There is no pharmacokinetic or safety data on the concomitant use of imatinib mesylate tablets at doses greater than 400 mg/day or the chronic use of concomitant acetaminophen and imatinib mesylate tablets.

##### In Vitro Studies

Cytochrome P450 (CYP450) Enzymes: Imatinib inhibits CYP2C9 and CYP2D6.

Transporter Systems: Imatinib is a substrate of BCRP, P-gp, OATP1A2, OATP1B3, and OCT1. Imatinib inhibits BCRP, BSEP, P-gp, MATE1, MATE2-K, OCT1 and OCT2.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In the 2-year rat carcinogenicity study administration of imatinib at 15, 30, and 60 mg/kg/day resulted in a statistically significant reduction in the longevity of males at 60 mg/kg/day and females at greater than or equal to 30 mg/kg/day. Target organs for neoplastic changes were the kidneys (renal tubule and renal pelvis), urinary bladder, urethra, preputial and clitoral gland, small intestine, parathyroid glands, adrenal glands and non-glandular stomach. Neoplastic lesions were not seen at: 30 mg/kg/day for the kidneys, urinary bladder, urethra, small intestine, parathyroid glands, adrenal glands and non-glandular stomach, and 15 mg/kg/day for the preputial and clitoral gland. The papilloma/carcinoma of the preputial/clitoral gland were noted at 30 and 60 mg/kg/day, representing approximately 0.5 to 4 or 0.3 to 2.4 times the human daily exposure (based on AUC) at 400 mg/day or 800 mg/day, respectively, and 0.4 to 3.0 times the daily exposure in children (based on AUC) at 340 mg/m<sup>2</sup>. The renal tubule adenoma/carcinoma, renal pelvis transitional cell neoplasms, the urinary bladder and urethra transitional cell papillomas, the small intestine adenocarcinomas, the parathyroid glands adenomas, the benign and malignant medullary tumors of the adrenal glands and the non-glandular stomach papillomas/carcinomas were noted at 60 mg/kg/day. The relevance of these findings in the rat carcinogenicity study for humans is not known.

Positive genotoxic effects were obtained for imatinib in an *in vitro* mammalian cell assay (Chinese hamster ovary) for clastogenicity (chromosome aberrations) in the presence of metabolic activation. Two intermediates of the manufacturing process, which are also present in the final product, are positive for mutagenesis in the Ames assay. One of these intermediates was also positive in the mouse lymphoma assay. Imatinib was not genotoxic when tested in an *in vitro* bacterial cell assay (Ames test), an *in vitro* mammalian cell assay (mouse lymphoma) and an *in vivo* rat micronucleus assay.

In a study of fertility, male rats were dosed for 70 days prior to mating and female rats were dosed 14 days prior to mating and through to gestational Day 6. Testicular and epididymal weights and percent motile sperm were decreased at 60 mg/kg, approximately three-fourths the maximum clinical dose of 800 mg/day based on BSA. This was not seen at doses less than or equal to 20 mg/kg (one-fourth the maximum human dose of 800 mg). The fertility of male and female rats was not affected.

Fertility was not affected in the preclinical fertility and early embryonic development study although lower testes and epididymal weights as well as a reduced number of motile sperm were observed in the high dose male rats. In the preclinical pre- and postnatal study in rats, fertility in the first generation offspring was also not affected by imatinib mesylate.

### 13.2 Animal Toxicology and/or Pharmacology

#### Toxicities from Long-Term Use

It is important to consider potential toxicities suggested by animal studies, specifically, *liver, kidney, and cardiac toxicity and immunosuppression*. Severe liver toxicity was observed in dogs treated for 2 weeks, with elevated liver enzymes, hepatocellular necrosis, bile duct necrosis, and bile duct hyperplasia. Renal toxicity was observed in monkeys treated for 2 weeks, with focal mineralization and dilation of the renal tubules and tubular nephrosis. Increased blood urea nitrogen (BUN) and creatinine were observed in several of these animals. An increased rate of opportunistic infections was observed with chronic imatinib treatment in laboratory animal studies. In a 39 week monkey study, treatment with imatinib resulted in worsening of normally suppressed malarial infections in these animals. Lymphopenia was observed in animals (as in humans). Additional long-term toxicities were identified in a 2-year rat study. Histopathological examination of the treated rats that died on study revealed cardiomyopathy (both sexes), chronic progressive nephropathy (females) and preputial gland papilloma as principal causes of death or reasons for sacrifice. Non-neoplastic lesions seen in this 2-year study which were not identified in earlier preclinical studies were the cardiovascular system, pancreas, endocrine organs, and teeth. The most important changes included cardiac hypertrophy and dilatation, leading to signs of cardiac insufficiency in some animals.

## 14 CLINICAL STUDIES

### 14.1 Chronic Myeloid Leukemia

#### Chronic Phase, Newly Diagnosed:


An open-label, multicenter, international randomized Phase 3 study (imatinib mesylate versus IFN+ Ara-C) has been conducted in patients with newly diagnosed Philadelphia

chromosome positive (Ph+) chronic myeloid leukemia (CML) in chronic phase. This study compared treatment with either single-agent imatinib mesylate or a combination of interferon-alpha (IFN) plus cytarabine (Ara-C). Patients were allowed to cross over to the alternative treatment arm if they failed to show a complete hematologic response (CHR) at 6 months, a major cytogenetic response (MCyR) at 12 months, or if they lost a CHR or MCyR. Patients with increasing WBC or severe intolerance to treatment were also allowed to cross over to the alternative treatment arm with the permission of the study monitoring committee (SMC). In the imatinib mesylate arm, patients were treated initially with 400 mg daily. Dose escalations were allowed from 400 mg daily to 600 mg daily, then from 600 mg daily to 800 mg daily. In the IFN arm, patients were treated with a target dose of IFN of 5 MIU/m<sup>2</sup>/day subcutaneously in combination with subcutaneous Ara-C 20 mg/m<sup>2</sup>/day for 10 days/month.


A total of 1,106 patients were randomized from 177 centers in 16 countries, 553 to each arm. Baseline characteristics were well balanced between the two arms. Median age was 51 years (range, 18 to 70 years), with 21.9% of patients greater than or equal to 60 years of age. There were 59% males and 41% females; 89.9% Caucasian and 4.7% black patients. At the cut-off for this analysis (7 years after last patient had been recruited), the median duration of first-line treatment was 82 and 8 months in the imatinib mesylate and IFN arm, respectively. The median duration of second-line treatment with imatinib mesylate was 64 months. Sixty percent of patients randomized to imatinib mesylate are still receiving first-line treatment. In these patients, the average dose of imatinib mesylate was 403 mg  $\pm$  57 mg. Overall, in patients receiving first line imatinib mesylate, the average daily dose delivered was 406 mg  $\pm$  76 mg. Due to discontinuations and cross-overs, only 2% of patients randomized to IFN were still on first-line treatment. In the IFN arm, withdrawal of consent (14%) was the most frequent reason for discontinuation of first-line therapy, and the most frequent reason for cross over to the imatinib mesylate arm was severe intolerance to treatment (26%) and progression (14%).

The primary efficacy endpoint of the study was progression-free survival (PFS). Progression was defined as any of the following events: progression to accelerated phase or blast crisis (AP/BC), death, loss of CHR or MCyR, or in patients not achieving a CHR an increasing WBC despite appropriate therapeutic management. The protocol specified that the progression analysis would compare the intent to treat (ITT) population: patients randomized to receive imatinib mesylate were compared with patients randomized to receive IFN. Patients that crossed over prior to progression were not censored at the time of cross-over, and events that occurred in these patients following cross-over were attributed to the original randomized treatment. The estimated rate of progression-free survival at 84 months in the ITT population was 81.2% [95% CI: 78, 85] in the imatinib mesylate arm and 60.6% [56, 65] in the IFN arm (p less than 0.0001, log-rank test), (Figure 1). With 7 years follow up there were 93 (16.8%) progression events in the imatinib mesylate arm: 37 (6.7%) progression to AP/BC, 31 (5.6%) loss of MCyR, 15 (2.7%) loss of CHR or increase in WBC and 10 (1.8%) CML unrelated deaths. In contrast, there were 165 (29.8%) events in the IFN+Ara-C arm of which 130 occurred during first-line treatment with IFN-Ara-C. The estimated rate of patients free of progression to accelerated phase (AP) or blast crisis (BC) at 84 months was 92.5% [90, 95] in the imatinib mesylate arm compared to the 85.1%, [82, 89] (p less than or equal to 0.001) in the IFN arm, (Figure 2). The annual rates of any progression events have decreased with time on therapy. The probability of remaining progression free at 60 months was 95% for patients who were in complete cytogenetic response (CCyR) with molecular response (greater than or equal to 3 log reduction in BCR-ABL transcripts as measured by quantitative reverse transcriptase polymerase chain reaction) at 12 months, compared to 89% for patients in CCyR but without a major molecular response and 70% in patients who were not in CCyR at this time point (p less than 0.001).

**Figure 1: Progression Free Survival (ITT Principle)**



**Figure 2: Time to Progression to AP or BC (ITT Principle)**



A total of 71 (12.8%) and 85 (15.4%) patients died in the imatinib mesylate and IFN+Ara-C group, respectively. At 84 months the estimated overall survival is 86.4% (83, 90) vs 83.3% (80, 87) in the randomized imatinib mesylate and the IFN+Ara-C group, respectively ( $p = 0.073$  log-rank test). The hazard ratio is 0.750 with 95% CI 0.547 to 1.028. This time-to-event endpoint may be affected by the high crossover rate from IFN+Ara-C to imatinib mesylate. Major cytogenetic response, hematologic response, evaluation of minimal residual disease (molecular response), time to accelerated phase or blast crisis and survival were main secondary endpoints. Response data are shown in Table 18. Complete hematologic response, major cytogenetic response and CCyR were also statistically significantly higher in the imatinib mesylate arm compared to the IFN + Ara-C arm (no cross-over data considered for evaluation of responses). Median time to CCyR in the 454 responders was 6 months (range, 2 to 64 months, 25<sup>th</sup> to 75<sup>th</sup> percentiles = 3 to 11 months) with 10% of responses seen only after 22 months of therapy.

**Table 18: Response in Newly Diagnosed CML Study (84-Month Data)**

|                                         | Imatinib Mesylate<br>n = 553 | IFN+Ara-C<br>n = 553 |
|-----------------------------------------|------------------------------|----------------------|
| <b>Best response rate</b>               |                              |                      |
| <b>Hematologic response<sup>1</sup></b> |                              |                      |
| CHR Rate n (%)                          | 534 (96.6%)*                 | 313 (56.6%)*         |
| [95% CI]                                | [94.7%, 97.9%]               | [52.4%, 60.8%]       |
| <b>Cytogenetic response<sup>2</sup></b> |                              |                      |
| Major cytogenetic response n (%)        | 472 (85.4%)*                 | 93 (16.8%)*          |
| [95% CI]                                | [82.1%, 88.2%]               | [13.8%, 20.2%]       |
| Unconfirmed <sup>3</sup>                | 88.6%*                       | 23.3%*               |
| Complete cytogenetic response n (%)     | 413 (74.7%)*                 | 36 (6.5%)*           |

|                          |              |            |
|--------------------------|--------------|------------|
| [95% CI]                 | [70.8, 78.3] | [4.6, 8.9] |
| Unconfirmed <sup>3</sup> | 82.5%*       | 11.6%*     |

\*p less than 0.001, Fischer's exact test.

**<sup>1</sup>Hematologic response criteria** (all responses to be confirmed after greater than or equal to 4 weeks): WBC less than  $10 \times 10^9/L$ , platelet less than  $450 \times 10^9/L$ , myelocyte + metamyelocyte less than 5% in blood, no blasts and promyelocytes in blood, no extramedullary involvement.

**<sup>2</sup>Cytogenetic response criteria** (confirmed after greater than or equal to 4 weeks): complete (0% Ph+ metaphases) or partial (1% to 35%). A major response (0% to 35%) combines both complete and partial responses.

**<sup>3</sup>Unconfirmed cytogenetic response** is based on a single bone marrow cytogenetic evaluation, therefore unconfirmed complete or partial cytogenetic responses might have had a lesser cytogenetic response on a subsequent bone marrow evaluation.

**Molecular response was defined as follows:** in the peripheral blood, after 12 months of therapy, reduction of greater than or equal to 3 logarithms in the amount of BCR-ABL transcripts (measured by real-time quantitative reverse transcriptase PCR assay) over a standardized baseline. Molecular response was only evaluated in a subset of patients who had a CCyR by 12 months or later (N = 333). The molecular response rate in patients who had a CCyR in the imatinib mesylate arm was 59% at 12 months and 72% at 24 months.

Physical, functional, and treatment-specific biologic response modifier scales from the FACT-BRM (Functional Assessment of Cancer Therapy - Biologic Response Modifier) instrument were used to assess patient-reported general effects of interferon toxicity in 1,067 patients with CML in chronic phase. After one month of therapy to six months of therapy, there was a 13% to 21% decrease in median index from baseline in patients treated with IFN, consistent with increased symptoms of IFN toxicity. There was no apparent change from baseline in median index for patients treated with imatinib mesylate.

An open-label, multicenter, randomized trial (imatinib mesylate versus nilotinib) was conducted to determine the efficacy of imatinib mesylate versus nilotinib in adult patients with cytogenetically confirmed, newly diagnosed Ph+ CML-CP. Patients were within 6 months of diagnosis and were previously untreated for CML-CP, except for hydroxyurea and/or anagrelide. Efficacy was based on a total of 846 patients: 283 patients in the imatinib mesylate 400 mg once daily group, 282 patients in the nilotinib 300 mg twice daily group, 281 patients in the nilotinib 400 mg twice daily group.

Median age was 46 years in the imatinib mesylate group and 47 years in both nilotinib groups, with 12%, 13%, and 10% of patients greater than or equal to 65 years of age in imatinib mesylate 400 mg once daily, nilotinib 300 mg twice daily and nilotinib 400 mg twice daily treatment groups, respectively. There were slightly more male than female patients in all groups (56%, 56%, and 62% in imatinib mesylate 400 mg once daily, nilotinib 300 mg twice daily and nilotinib 400 mg twice daily treatment groups, respectively). More than 60% of all patients were Caucasian, and 25% were Asian.

The primary data analysis was performed when all 846 patients completed 12 months of treatment or discontinued earlier. Subsequent analyses were done when patients completed 24, 36, 48, and 60 months of treatment or discontinued earlier. The median time on treatment was approximately 61 months in all three treatment groups.

The primary efficacy endpoint was major molecular response (MMR) at 12 months after the start of study medication. MMR was defined as less than or equal to 0.1% BCR-ABL/ABL % by international scale measured by RQ-PCR, which corresponds to a greater than or equal to 3 log reduction of BCR-ABL transcript from standardized baseline. Efficacy endpoints are summarized in Table 19.

Twelve patients in the imatinib mesylate arm progressed to either accelerated phase or blast crises (7 patients within first 6 months, 2 patients within 6 to 12 months, 2 patients within 12 to 18 months and 1 patient within 18 to 24 months) while two patients on the nilotinib arm progressed to either accelerated phase or blast crisis (both within the first 6 months of treatment).

**Table 19: Efficacy (MMR and CCyR) of Imatinib Mesylate Compared to Nilotinib in Newly Diagnosed Ph+ CML-CP**

| Imatinib Mesylate<br>400 mg<br>once daily | Nilotinib<br>300 mg<br>twice daily |
|-------------------------------------------|------------------------------------|
|-------------------------------------------|------------------------------------|

|                                         | N = 283          | N = 282          |
|-----------------------------------------|------------------|------------------|
| MMR at 12 months (95% CI)               | 22% (17.6, 27.6) | 44% (38.4, 50.3) |
| P-Value <sup>a</sup>                    | <0.0001          |                  |
| CCyR <sup>b</sup> by 12 months (95% CI) | 65% (59.2, 70.6) | 80% (75.0, 84.6) |
| MMR at 24 months (95% CI)               | 38% (31.8, 43.4) | 62% (55.8, 67.4) |
| CCyR <sup>b</sup> by 24 months (95% CI) | 77% (71.7, 81.8) | 87% (82.4, 90.6) |

Abbreviations: CCyR, complete cytogenetic response; MMR, major molecular response; Ph+ CML-CP, Philadelphia chromosome positive chronic myeloid leukemia-chronic phase.

<sup>a</sup> CMH test stratified by Sokal risk group.

<sup>b</sup> CCyR: 0% Ph+ metaphases. Cytogenetic responses were based on the percentage of Ph-positive metaphases among greater than or equal to 20 metaphase cells in each bone marrow sample.

By 60 months, MMR was achieved by 60% of patients on imatinib mesylate and 77% of patients on nilotinib.

Median overall survival was not reached in either arm. At the time of the 60-month final analysis, the estimated survival rate was 91.7% for patients on imatinib mesylate and 93.7% for patients on nilotinib.

**Late Chronic Phase CML and Advanced Stage CML:** Three international, open-label, single-arm Phase 2 studies were conducted to determine the safety and efficacy of imatinib mesylate in patients with Ph+ CML: 1) in the chronic phase after failure of IFN therapy, 2) in accelerated phase disease, or 3) in myeloid blast crisis. About 45% of patients were women and 6% were black. In clinical studies, 38% to 40% of patients were greater than or equal to 60 years of age and 10% to 12% of patients were greater than or equal to 70 years of age.

**Chronic Phase, Prior Interferon-Alpha Treatment:** 532 patients were treated at a starting dose of 400 mg; dose escalation to 600 mg was allowed. The patients were distributed in three main categories according to their response to prior interferon: failure to achieve (within 6 months), or loss of a complete hematologic response (29%), failure to achieve (within 1 year) or loss of a major cytogenetic response (35%), or intolerance to interferon (36%). Patients had received a median of 14 months of prior IFN therapy at doses greater than or equal to  $25 \times 10^6$  units/week and were all in late chronic phase, with a median time from diagnosis of 32 months. Effectiveness was evaluated on the basis of the rate of hematologic response and by bone marrow exams to assess the rate of major cytogenetic response (up to 35% Ph+ metaphases) or CCyR (0% Ph+ metaphases). Median duration of treatment was 29 months with 81% of patients treated for greater than or equal to 24 months (maximum = 31.5 months). Efficacy results are reported in Table 20. Confirmed major cytogenetic response rates were higher in patients with IFN intolerance (66%) and cytogenetic failure (64%), than in patients with hematologic failure (47%). Hematologic response was achieved in 98% of patients with cytogenetic failure, 94% of patients with hematologic failure, and 92% of IFN-intolerant patients.

**Accelerated Phase:** 235 patients with accelerated phase disease were enrolled. These patients met one or more of the following criteria: greater than or equal to 15% to less than 30% blasts in PB or BM; greater than or equal to 30% blasts + promyelocytes in PB or BM; greater than or equal to 20% basophils in PB; and less than  $100 \times 10^9$ /L platelets. The first 77 patients were started at 400 mg, with the remaining 158 patients starting at 600 mg.

Effectiveness was evaluated primarily on the basis of the rate of hematologic response, reported as either complete hematologic response, no evidence of leukemia (i.e., clearance of blasts from the marrow and the blood, but without a full peripheral blood recovery as for complete responses), or return to chronic phase CML. Cytogenetic responses were also evaluated. Median duration of treatment was 18 months with 45% of patients treated for greater than or equal to 24 months (maximum = 35 months). Efficacy results are reported in Table 20. Response rates in accelerated phase CML were higher for the 600 mg dose group than for the 400 mg group: hematologic response (75% vs 64%), confirmed and unconfirmed major cytogenetic response (31% vs 19%).

**Myeloid Blast Crisis:** 260 patients with myeloid blast crisis were enrolled. These patients had greater than or equal to 30% blasts in PB or BM and/or extramedullary involvement other than spleen or liver; 95 (37%) had received prior chemotherapy for treatment of either accelerated phase or blast crisis ("pretreated patients") whereas 165 (63%) had not ("untreated patients"). The first 37 patients were started at 400 mg; the remaining 223 patients were started at 600 mg.

Effectiveness was evaluated primarily on the basis of rate of hematologic response, reported as either complete hematologic response, no evidence of leukemia, or return to chronic phase CML using the same criteria as for the study in accelerated phase. Cytogenetic responses were also assessed. Median duration of treatment was 4 months with 21% of patients treated for greater than or equal to 12 months and 10% for

greater than or equal to 24 months (maximum = 35 months). Efficacy results are reported in Table 20. The hematologic response rate was higher in untreated patients than in treated patients (36% vs 22%, respectively) and in the group receiving an initial dose of 600 mg rather than 400-mg (33% vs 16%). The confirmed and unconfirmed major cytogenetic response rate was also higher for the 600 mg dose group than for the 400-mg dose group (17% vs 8%).

**Table 20: Response in Chronic Myeloid Leukemia Studies**

|                                               | <b>Chronic phase IFN failure(n = 532)</b> | <b>Accelerated phase(n = 235)</b>  | <b>Myeloid blast crisis(n = 260)</b> |
|-----------------------------------------------|-------------------------------------------|------------------------------------|--------------------------------------|
|                                               | <b>400 mg</b>                             | <b>600 mg n = 158400 mg n = 77</b> | <b>600 mg n = 223400 mg n = 37</b>   |
|                                               |                                           | <b>% of patients [CI 95%]</b>      |                                      |
| <b>Hematologic response<sup>1</sup></b>       | 95% [92.3–96.3]                           | 71% [64.8-76.8]                    | 31% [25.2–36.8]                      |
| Complete hematologic response (CHR)           | 95%                                       | 38%                                | 7%                                   |
| No evidence of leukemia (NEL)                 | Not applicable                            | 13%                                | 5%                                   |
| Return to chronic phase (RTC)                 | Not applicable                            | 20%                                | 18%                                  |
| <b>Major cytogenetic response<sup>2</sup></b> | 60% [55.3–63.8]                           | 21% [16.2–27.1]                    | 7% [4.5–11.2]                        |
| (Unconfirmed <sup>3</sup> )                   | (65%)                                     | (27%)                              | (15%)                                |
| Complete <sup>4</sup>                         | 39% (47%)                                 | 16% (20%)                          | 2% (7%)                              |
| (Unconfirmed <sup>3</sup> )                   |                                           |                                    |                                      |

Abbreviations: BM, bone marrow; PB, peripheral blood.

<sup>1</sup>**Hematologic response criteria** (all responses to be confirmed after greater than or equal to 4 weeks): CHR: Chronic phase study [WBC less than  $10 \times 10^9/L$ , platelet less than  $450 \times 10^9/L$ , myelocytes + metamyelocytes less than 5% in blood, no blasts and promyelocytes in blood, basophils less than 20%, no extramedullary involvement] and in the accelerated and blast crisis studies [absolute neutrophil count (ANC) greater than or equal to  $1.5 \times 10^9/L$ , platelets greater than or equal to  $100 \times 10^9/L$ , no blood blasts, BM blasts less than 5% and no extramedullary disease].

NEL: Same criteria as for CHR but ANC greater than or equal to  $1 \times 10^9/L$  and platelets greater than or equal to  $20 \times 10^9/L$  (accelerated and blast crisis studies).

RTC: less than 15% blasts BM and PB, less than 30% blasts + promyelocytes in BM and PB, less than 20% basophils in PB, no extramedullary disease other than spleen and liver (accelerated and blast crisis studies).

<sup>2</sup>**Cytogenetic response criteria** (confirmed after greater than or equal to 4 weeks): complete (0% Ph+ metaphases) or partial (1% to 35%). A major response (0% to 35%) combines both complete and partial responses.

<sup>3</sup>**Unconfirmed cytogenetic response** is based on a single bone marrow cytogenetic evaluation, therefore unconfirmed complete or partial cytogenetic responses might have had a lesser cytogenetic response on a subsequent bone marrow evaluation.

<sup>4</sup>**Complete cytogenetic response** confirmed by a second bone marrow cytogenetic evaluation performed at least 1 month after the initial bone marrow study.

The median time to hematologic response was 1 month. In late chronic phase CML, with a median time from diagnosis of 32 months, an estimated 87.8% of patients who achieved MCyR maintained their response 2 years after achieving their initial response. After 2 years of treatment, an estimated 85.4% of patients were free of progression to AP or BC, and estimated overall survival was 90.8% [88.3, 93.2]. In accelerated phase, median duration of hematologic response was 28.8 months for patients with an initial dose of 600 mg (16.5 months for 400 mg). An estimated 63.8% of patients who achieved MCyR were still in response 2 years after achieving initial response. The median survival was 20.9 [13.1, 34.4] months for the 400 mg group and was not yet reached for the 600 mg group ( $p = 0.0097$ ). An estimated 46.2% [34.7, 57.7] vs 65.8% [58.4, 73.3] of patients were still alive after 2 years of treatment in the 400 mg vs 600 mg dose groups, respectively. In blast crisis, the estimated median duration of hematologic response is 10 months. An estimated 27.2% [16.8, 37.7] of hematologic responders maintained their response 2 years after achieving their initial response. Median survival was 6.9 [5.8, 8.6] months, and an estimated 18.3% [13.4, 23.3] of all patients with blast crisis were alive 2 years after start of study.

Efficacy results were similar in men and women and in patients younger and older than age 65. Responses were seen in black patients, but there were too few black patients to

allow a quantitative comparison.

## 14.2 Pediatric CML

A total of 51 pediatric patients with newly diagnosed and untreated CML in chronic phase were enrolled in an open-label, multicenter, single-arm Phase 2 trial. Patients were treated with imatinib mesylate 340 mg/m<sup>2</sup>/day, with no interruptions in the absence of dose limiting toxicity. Complete hematologic response (CHR) was observed in 78% of patients after 8 weeks of therapy. The complete cytogenetic response rate (CCyR) was 65%, comparable to the results observed in adults. Additionally, partial cytogenetic response (PCyR) was observed in 16%. The majority of patients who achieved a CCyR developed the CCyR between Months 3 and 10 with a median time to response based on the Kaplan-Meier estimate of 6.74 months. Patients were allowed to be removed from protocol therapy to undergo alternative therapy, including hematopoietic stem cell transplantation. Thirty-one children received stem cell transplantation. Of the 31 children, 5 were transplanted after disease progression on study and 1 withdrew from study during first week treatment and received transplant approximately 4 months after withdrawal. Twenty-five children withdrew from protocol therapy to undergo stem cell transplant after receiving a median of 9 twenty-eight day courses (range, 4 to 24). Of the 25 patients 13 (52%) had CCyR and 5 (20%) had PCyR at the end of protocol therapy.

One open-label, single-arm study enrolled 14 pediatric patients with Ph+ chronic phase CML recurrent after stem cell transplant or resistant to interferon-alpha therapy. These patients had not previously received imatinib mesylate and ranged in age from 3 to 20 years old; 3 were 3 to 11 years old, 9 were 12 to 18 years old, and 2 were greater than 18 years old. Patients were treated at doses of 260 mg/m<sup>2</sup>/day (n = 3), 340 mg/m<sup>2</sup>/day (n = 4), 440 mg/m<sup>2</sup>/day (n = 5) and 570 mg/m<sup>2</sup>/day (n = 2). In the 13 patients for whom cytogenetic data are available, 4 achieved a major cytogenetic response, 7 achieved a CCyR, and 2 had a minimal cytogenetic response.

In a second study, 2 of 3 patients with Ph+ chronic phase CML resistant to interferon-alpha therapy achieved a CCyR at doses of 242 and 257 mg/m<sup>2</sup>/day.

## 14.3 Acute Lymphoblastic Leukemia

A total of 48 Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) patients with relapsed/refractory disease were studied, 43 of whom received the recommended imatinib mesylate dose of 600 mg/day. In addition 2 patients with relapsed/refractory Ph+ ALL received imatinib mesylate 600 mg/day in a Phase 1 study.

Confirmed and unconfirmed hematologic and cytogenetic response rates for the 43 relapsed/refractory Ph+ALL Phase 2 study patients and for the 2 Phase 1 patients are shown in Table 21. The median duration of hematologic response was 3.4 months and the median duration of MCyR was 2.3 months.

**Table 21: Effect of Imatinib Mesylate on Relapsed/Refractory Ph+ ALL**

|                | Phase 2 study<br>(N = 43)<br>n (%) | Phase 1 study<br>(N = 2)<br>n(%) |
|----------------|------------------------------------|----------------------------------|
| <b>CHR</b>     | 8 (19)                             | 2 (100)                          |
| <b>NEL</b>     | 5 (12)                             |                                  |
| <b>RTC/PHR</b> | 11 (26)                            |                                  |
| <b>MCyR</b>    | 15 (35)                            |                                  |
| <b>CCyR</b>    | 9 (21)                             |                                  |
| <b>PCyR</b>    | 6 (14)                             |                                  |

Abbreviations: CCyR, complete cytogenetic response; CHR, complete hematologic response; MCyR, major cytogenetic response; NEL, no evidence of leukemia; PCyR, partial cytogenetic response; Ph+ ALL, Philadelphia chromosome positive acute lymphoblastic leukemia; PHR, partial hematologic response; RTC, return to chronic phase.

## 14.4 Pediatric ALL

Pediatric and young adult patients with very high risk ALL, defined as those with an expected 5-year event-free survival (EFS) less than 45%, were enrolled after induction therapy on a multicenter, non-randomized cooperative group pilot protocol.

The safety and effectiveness of imatinib mesylate tablets (340 mg/m<sup>2</sup> /day) in combination with intensive chemotherapy was evaluated in a subgroup of patients with Ph+ ALL. The protocol included intensive chemotherapy and hematopoietic stem cell transplant after 2 courses of chemotherapy for patients with an appropriate HLA-matched family donor. There were 92 eligible patients with Ph+ ALL enrolled. The median age was 9.5 years (1 to 21 years: 2.2% between 1 and less than 2 years, 56.5%

between 2 and less than 12 years, 34.8% between 12 and less than 18 years, and 6.5% between 18 and 21 years). Sixty-four percent were male, 75% were white, 9% were Asian/Pacific Islander, and 5% were black. In 5 successive cohorts of patients, imatinib mesylate tablets exposure was systematically increased by earlier introduction and prolonged duration. Cohort 1 received the lowest intensity and cohort 5 received the highest intensity of imatinib mesylate tablets exposure.

There were 50 patients with Ph+ ALL assigned to cohort 5 all of whom received imatinib mesylate tablets plus chemotherapy; 30 were treated exclusively with chemotherapy and imatinib mesylate tablets and 20 received chemotherapy plus imatinib mesylate tablets and then underwent hematopoietic stem cell transplant, followed by further imatinib mesylate tablets treatment. Patients in cohort 5 treated with chemotherapy received continuous daily exposure to imatinib mesylate tablets beginning in the first course of post induction chemotherapy continuing through maintenance cycles 1 through 4 chemotherapy. During maintenance cycles 5 through 12, imatinib mesylate tablets was administered 28 days out of the 56 day cycle. Patients who underwent hematopoietic stem cell transplant received 42 days of imatinib mesylate tablets prior to HSCT, and 28 weeks (196 days) of imatinib mesylate tablets after the immediate post-transplant period. The estimated 4-year EFS of patients in cohort 5 was 70% (95% CI: 54, 81). The median follow-up time for EFS at data cutoff in cohort 5 was 40.5 months.

#### 14.5 Myelodysplastic/Myeloproliferative Diseases

An open-label, multicenter, Phase 2 clinical trial was conducted testing imatinib mesylate in diverse populations of patients suffering from life-threatening diseases associated with Abl, Kit or PDGFR protein tyrosine kinases. This study included 7 patients with MDS/MPD. These patients were treated with imatinib mesylate 400 mg daily (imatinib as free base). The ages of the enrolled patients ranged from 20 to 86 years. A further 24 patients with MDS/MPD aged 2 to 79 years were reported in 12 published case reports and a clinical study. These patients also received imatinib mesylate at a dose of 400 mg (imatinib as free base) daily with the exception of three patients who received lower doses. Of the total population of 31 patients treated for MDS/MPD, 14 (45%) achieved a complete hematological response and 12 (39%) a major cytogenetic response (including 10 with a CCyR). Sixteen patients had a translocation, involving chromosome 5q33 or 4q12, resulting in a PDGFR gene re-arrangement. All of these patients responded hematologically (13 completely). Cytogenetic response was evaluated in 12 out of 14 patients, all of whom responded (10 patients completely). Only 1 (7%) out of the 14 patients without a translocation associated with PDGFR gene re-arrangement achieved a complete hematological response and none achieved a major cytogenetic response. A further patient with a PDGFR gene re-arrangement in molecular relapse after bone marrow transplant responded molecularly. Median duration of therapy was 12.9 months (0.8 to 26.7) in the 7 patients treated within the Phase 2 study and ranged between 1 week and more than 18 months in responding patients in the published literature. Results are provided in Table 22. Response durations of Phase 2 study patients ranged from 141+ days to 457+ days.

**Table 22: Response in MDS/MPD**

|                                  | <b>Number<br/>of<br/>patients<br/>N</b> | <b>Complete<br/>hematologic<br/>response<br/>N (%)</b> | <b>Major<br/>cytogenetic<br/>response<br/>N (%)</b> |
|----------------------------------|-----------------------------------------|--------------------------------------------------------|-----------------------------------------------------|
| <b>Overall<br/>population</b>    | 31                                      | 14 (45)                                                | 12 (39)                                             |
| Chromosome<br>5<br>translocation | 14                                      | 11 (79)                                                | 11 (79)                                             |
| Chromosome<br>4<br>translocation | 2                                       | 2 (100)                                                | 1 (50)                                              |
| Others/no<br>translocation       | 14                                      | 1 (7)                                                  | 0                                                   |
| Molecular<br>relapse             | 1                                       | NE <sup>1</sup>                                        | NE <sup>1</sup>                                     |

Abbreviations: NE, not evaluable; MDS/MPD, myelodysplastic/myeloproliferative disease.

#### 14.6 Aggressive Systemic Mastocytosis

One open-label, multicenter, Phase 2 study was conducted testing imatinib mesylate in diverse populations of patients with life-threatening diseases associated with Abl, Kit or PDGFR protein tyrosine kinases. This study included 5 patients with ASM treated with 100 mg to 400 mg of imatinib mesylate (imatinib as free base) daily. These 5 patients ranged from 49 to 74 years of age. In addition to these 5 patients, 10 published case

reports and case series describe the use of imatinib mesylate in 23 additional patients with ASM aged 26 to 85 years who also received 100 mg to 400 mg of imatinib mesylate (imatinib as free base) daily.

Cytogenetic abnormalities were evaluated in 20 of the 28 ASM patients treated with imatinib mesylate from the published reports and in the Phase 2 study. Seven of these 20 patients had the FIP1L1-PDGFR $\alpha$  fusion kinase (or CHIC2 deletion). Patients with this cytogenetic abnormality were predominantly males and had eosinophilia associated with their systemic mast cell disease. Two patients had a Kit mutation in the juxtamembrane region (one Phe522Cys and one K509I) and four patients had a D816V c-Kit mutation (not considered sensitive to imatinib mesylate), one with concomitant CML.

Of the 28 patients treated for ASM, 8 (29%) achieved a complete hematologic response and 9 (32%) a partial hematologic response (PHR) (61% overall response rate). Median duration of imatinib mesylate therapy for the 5 ASM patients in the Phase 2 study was 13 months (range, 1.4 to 22.3 months) and between 1 month and more than 30 months in the responding patients described in the published medical literature. A summary of the response rates to imatinib mesylate in ASM is provided in Table 23. Response durations of literature patients ranged from 1+ to 30+ months.

**Table 23: Response in ASM**

| <b>Cytogenetic abnormality</b>                          | <b>Number of patients<br/>N</b> | <b>Complete hematologic response<br/>N (%)</b> | <b>Partial hematologic response<br/>N (%)</b> |
|---------------------------------------------------------|---------------------------------|------------------------------------------------|-----------------------------------------------|
| FIP1L1-PDGFR $\alpha$ fusion kinase (or CHIC2 deletion) | 7                               | 7 (100)                                        | 0                                             |
| Juxtamembrane mutation                                  | 2                               | 0                                              | 2 (100)                                       |
| Unknown or no cytogenetic abnormality detected          | 15                              | 0                                              | 7 (44)                                        |
| D816V mutation                                          | 4                               | 1* (25)                                        | 0                                             |
| <b>Total</b>                                            | <b>28</b>                       | <b>8 (29)</b>                                  | <b>9 (32)</b>                                 |

Abbreviations: ASM, aggressive systemic mastocytosis; PDGFR, platelet-derived growth factor receptor. \*Patient had concomitant chronic myeloid leukemia CML and ASM.

Imatinib mesylate has not been shown to be effective in patients with less aggressive forms of systemic mastocytosis (SM). Imatinib mesylate is therefore not recommended for use in patients with cutaneous mastocytosis, indolent systemic mastocytosis (smoldering SM or isolated bone marrow mastocytosis), SM with an associated clonal hematological non-mast cell lineage disease, mast cell leukemia, mast cell sarcoma or extracutaneous mastocytoma. Patients that harbor the D816V mutation of c-Kit are not sensitive to imatinib mesylate and should not receive imatinib mesylate.

#### **14.7 Hypereosinophilic Syndrome/Chronic Eosinophilic Leukemia**

One open-label, multicenter, Phase 2 study was conducted testing imatinib mesylate in diverse populations of patients with life-threatening diseases associated with Abl, Kit or PDGFR protein tyrosine kinases. This study included 14 patients with Hypereosinophilic Syndrome/Chronic Eosinophilic Leukemia (HES/CEL). HES patients were treated with 100 mg to 1000 mg of imatinib mesylate daily. The ages of these patients ranged from 16 to 64 years. A further 162 patients with HES/CEL aged 11 to 78 years were reported in 35 published case reports and case series. These patients received imatinib mesylate at doses of 75 mg to 800 mg (imatinib as free base) daily. Hematologic response rates are summarized in Table 24. Response durations for literature patients ranged from 6+ weeks to 44 months.

**Table 24: Response in HES/CEL**

| <b>Cytogenetic abnormality</b>               | <b>Number of patients</b> | <b>Complete hematological response<br/>N (%)</b> | <b>Partial hematological response<br/>N (%)</b> |
|----------------------------------------------|---------------------------|--------------------------------------------------|-------------------------------------------------|
| Positive FIP1L1-PDGFR $\alpha$ fusion kinase | 61                        | 61 (100)                                         | 0                                               |
| Negative FIP1L1-PDGFR $\alpha$ fusion kinase | 56                        | 12 (21)                                          | 9 (16)                                          |
| Unknown                                      |                           |                                                  |                                                 |

|                         |     |          |         |
|-------------------------|-----|----------|---------|
| cytogenetic abnormality | 59  | 34 (58)  | 7 (12)  |
| Total                   | 176 | 107 (61) | 23 (13) |

Abbreviations: CEL, chronic eosinophilic leukemia; HES, hypereosinophilic syndrome; PDGFR, platelet-derived growth factor receptor.

#### 14.8 Dermatofibrosarcoma Protuberans

Dermatofibrosarcoma Protuberans (DFSP) is a cutaneous soft tissue sarcoma. It is characterized by a translocation of chromosomes 17 and 22 that results in the fusion of the collagen type 1 alpha 1 gene and the PDGF B gene.

An open-label, multicenter, Phase 2 study was conducted testing imatinib mesylate in a diverse population of patients with life-threatening diseases associated with Abl, Kit or PDGFR protein tyrosine kinases. This study included 12 patients with DFSP who were treated with imatinib mesylate 800 mg daily (imatinib as free base) (age range, 23 to 75 years). DFSP was metastatic, locally recurrent following initial surgical resection and not considered amenable to further surgery at the time of study entry. A further 6 DFSP patients treated with imatinib mesylate are reported in 5 published case reports, their ages ranging from 18 months to 49 years. The total population treated for DFSP therefore comprises 18 patients, 8 of them with metastatic disease. The adult patients reported in the published literature were treated with either 400 mg (4 cases) or 800 mg (1 case) imatinib mesylate (imatinib as free base) daily. A single pediatric patient received 400 mg/m<sup>2</sup>/daily, subsequently increased to 520 mg/m<sup>2</sup>/daily. Ten patients had the PDGF B gene rearrangement, 5 had no available cytogenetics and 3 had complex cytogenetic abnormalities. Responses to treatment are described in Table 25.

**Table 25: Response in DFSP**

|                    | Number of patients (n = 18) | %  |
|--------------------|-----------------------------|----|
| Complete response  | 7                           | 39 |
| Partial response * | 8                           | 44 |
| Total responders   | 15                          | 83 |

\* 5 patients made disease free by surgery.

Twelve of these 18 patients either achieved a complete response (7 patients) or were made disease free by surgery after a partial response (5 patients, including one child) for a total complete response rate of 67%. A further 3 patients achieved a partial response, for an overall response rate of 83%. Of the 8 patients with metastatic disease, five responded (62%), three of them completely (37%). For the 10 study patients with the PDGF B gene rearrangement, there were 4 complete and 6 partial responses. The median duration of response in the Phase 2 study was 6.2 months, with a maximum duration of 24.3 months, while in the published literature it ranged between 4 weeks and more than 20 months.

#### 14.9 Gastrointestinal Stromal Tumors

##### *Unresectable and/or Malignant Metastatic GIST*

Two open-label, randomized, multinational Phase 3 studies were conducted in patients with unresectable or metastatic malignant GIST. The two study designs were similar allowing a predefined combined analysis of safety and efficacy. A total of 1,640 patients were enrolled into the two studies and randomized 1:1 to receive either 400 mg or 800 mg orally daily continuously until disease progression or unacceptable toxicity. Patients in the 400 mg daily treatment group who experienced disease progression were permitted to crossover to receive treatment with 800 mg daily. The studies were designed to compare response rates, progression-free survival and overall survival between the dose groups. Median age at patient entry was 60 years. Males comprised 58% of the patients enrolled. All patients had a pathologic diagnosis of CD117 positive unresectable and/or metastatic malignant GIST.

The primary objective of the two studies was to evaluate either progression-free survival (PFS) with a secondary objective of overall survival (OS) in one study or overall survival with a secondary objective of PFS in the other study. A planned analysis of both OS and PFS from the combined datasets from these two studies was conducted. Results from this combined analysis are shown in Table 26.

**Table 26: Overall Survival, Progression-Free Survival and Tumor Response Rates in the Phase 3 GIST Trials**

|                           | Imatinib Mesylate 400 mg<br>N = 818 | Imatinib Mesylate 800 mg<br>N = 822 |
|---------------------------|-------------------------------------|-------------------------------------|
| Progression-free survival | 19.0                                | 22.2                                |

|                                    |             |             |
|------------------------------------|-------------|-------------|
| <b>(months) Median</b>             | 18.7        | 23.2        |
| 95% CI                             | 17.4-21.2   | 20.8-24.9   |
| <b>Overall survival (months)</b>   | 49          | 48.7        |
| 95% CI                             | 45.3-60     | 45.3-51.6   |
| <b>Best overall tumor response</b> | 43 (5.3%)   | 41 (5%)     |
| Complete response                  |             |             |
| Partial response                   | 377 (46.1%) | 402 (48.9%) |

Abbreviation: GIST, gastrointestinal stromal tumors.

Median follow up for the combined studies was 37.5 months. There were no observed differences in overall survival between the treatment groups ( $p = 0.98$ ). Patients who crossed over following disease progression from the 400 mg/day treatment group to the 800 mg/day treatment group ( $n = 347$ ) had a 3.4 month median and a 7.7 month mean exposure to imatinib mesylate following crossover.


One open-label, multinational Phase 2 study was conducted in patients with Kit (CD117) positive unresectable or metastatic malignant GIST. In this study, 147 patients were enrolled and randomized to receive either 400 mg or 600 mg orally every day for up to 36 months. The primary outcome of the study was objective response rate. Tumors were required to be measurable at entry in at least one site of disease, and response characterization was based on Southwestern Oncology Group (SWOG) criteria. There were no differences in response rates between the 2 dose groups. The response rate was 68.5% for the 400 mg group and 67.6% for the 600 mg group. The median time to response was 12 weeks (range was 3 to 98 weeks) and the estimated median duration of response is 118 weeks (95% CI: 86, not reached).

#### *Adjuvant Treatment of GIST*

In the adjuvant setting, imatinib mesylate was investigated in a multicenter, double-blind, placebo-controlled, randomized trial involving 713 patients (Study 1). Patients were randomized one to one to imatinib mesylate (imatinib as free base) at 400 mg/day or matching placebo for 12 months. The ages of these patients ranged from 18 to 91 years. Patients were included who had a histologic diagnosis of primary GIST, expressing KIT protein by immunohistochemistry and a tumor size greater than or equal to 3 cm in maximum dimension with complete gross resection of primary GIST within 14 to 70 days prior to registration.

Recurrence-free survival (RFS) was defined as the time from date of randomization to the date of recurrence or death from any cause. In a planned interim analysis, the median follow up was 15 months in patients without a RFS event; there were 30 RFS events in the 12-month imatinib mesylate arm compared to 70 RFS events in the placebo arm with a hazard ratio of 0.398 (95% CI: 0.259, 0.610),  $p$  less than 0.0001. After the interim analysis of RFS, 79 of the 354 patients initially randomized to the placebo arm were eligible to cross over to the 12-month imatinib mesylate arm. Seventy-two of these 79 patients subsequently crossed over to imatinib mesylate therapy. In an updated analysis, the median follow-up for patients without a RFS event was 50 months. There were 74 (21%) RFS events in the 12-month imatinib mesylate arm compared to 98 (28%) events in the placebo arm with a hazard ratio of 0.718 (95% CI: 0.531 to 0.971) (Figure 3). The median follow-up for OS in patients still living was 61 months. There were 26 (7%) and 33 (9%) deaths in the 12-month imatinib mesylate and placebo arms, respectively with a hazard ratio of 0.816 (95% CI: 0.488 to 1.365).

#### **Figure 3: Study 1 Recurrence-Free Survival (ITT Population)**




A second randomized, multicenter, open-label, Phase 3 trial in the adjuvant setting (Study 2) compared 12 months of imatinib mesylate treatment to 36 months of imatinib mesylate (imatinib as free base) treatment at 400 mg/day in adult patients with KIT (CD117) positive GIST after surgical resection with one of the following: tumor diameter greater than 5 cm and mitotic count greater than 5/50 high power fields (HPF), or tumor diameter greater than 10 cm and any mitotic count, or tumor of any size with mitotic count greater than 10/50 HPF, or tumors ruptured into the peritoneal cavity. There were a total of 397 patients randomized in the trial with 199 patients on the 12-month treatment arm and 198 patients on the 36-month treatment arm. The median age was 61 years (range, 22 to 84 years).


RFS was defined as the time from date of randomization to the date of recurrence or death from any cause. The median follow-up for patients without a RFS event was 42 months. There were 84 (42%) RFS events in the 12-month treatment arm and 50 (25%) RFS events in the 36-month treatment arm. Thirty-six months of imatinib mesylate treatment significantly prolonged RFS compared to 12 months of imatinib mesylate treatment with a hazard ratio of 0.46 (95% CI: 0.32, 0.65),  $p$  less than 0.0001 (Figure 4).

The median follow-up for overall survival (OS) in patients still living was 48 months. There were 25 (13%) deaths in the 12-month treatment arm and 12 (6%) deaths in the 36-month treatment arm. Thirty-six months of imatinib mesylate treatment significantly prolonged OS compared to 12 months of imatinib mesylate treatment with a hazard ratio of 0.45 (95% CI: 0.22, 0.89),  $p=0.0187$  (Figure 5).

**Figure 4: Study 2 Recurrence-Free Survival (ITT Population)**



**Figure 5: Study 2 Overall Survival (ITT Population)**



## 15 REFERENCES

OSHA Hazardous Drugs. *OSHA*. [Accessed on 20-September-2013, from <http://www.osha.gov/SLTC/hazardousdrugs/index.html>]

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Each imatinib mesylate film-coated tablet contains 100 mg or 400 mg of imatinib free base.

### • 100 mg Tablets

Brownish orange, slightly biconvex, round film-coated tablets with functional scoring, engraved "IMA" over score "100" on one side, "APO" on the other side

Bottles of 30 tablets.....NDC 60505-2900-3  
 Bottles of 90 tablets.....NDC 60505-2900-9  
 Bottles of 100 tablets.....NDC 60505-2900-1  
 Bottles of 1,000 tablets.....NDC 60505-2900-8  
 Blisters of 100 tablets (10 x 10).....NDC 60505-2900-0

### • 400 mg Tablets

Brownish orange, capsule shaped, biconvex film-coated tablets with functional scoring, engraved "IMA" score "400" on one side, "APO" on the other side

Bottles of 30 tablets.....NDC 60505-2901-3  
 Bottles of 90 tablets.....NDC 60505-2901-9  
 Bottles of 100 tablets.....NDC 60505-2901-1  
 Bottles of 500 tablets.....NDC 60505-2901-5  
 Blisters of 100 tablets (10 x 10).....NDC 60505-2901-0

## Storage and Handling

Store at 20°C to 25°C (68°F to 77°F); excursions permitted from 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from moisture.

Dispense in a tight container, USP.

Do not crush imatinib mesylate tablets. Avoid direct contact of crushed tablets with the skin or mucous membranes. If such contact occurs, wash thoroughly as outlined in the references. Avoid exposure to crushed tablets.

## 17 PATIENT COUNSELING INFORMATION

### Dosing and Administration

Advise patients to take imatinib mesylate tablets exactly as prescribed, not to change their dose or to stop taking imatinib mesylate unless they are told to do so by their doctor. If the patient missed a dose of imatinib mesylate tablets, the patient should take the next scheduled dose at its regular time. The patient should not take two doses at the same time. Advise patients to take imatinib mesylate tablets with a meal and a large glass of water [see *Dosage and Administration* (2.1)].

### Fluid Retention and Edema

Inform patients of the possibility of developing edema and fluid retention. Advise patients to contact their health care provider if unexpected rapid weight gain occurs [see *Warnings and Precautions* (5.1)].

#### Hepatotoxicity

Inform patients of the possibility of developing liver function abnormalities and serious hepatic toxicity. Advise patients to immediately contact their health care provider if signs of liver failure occur, including jaundice, anorexia, bleeding, or bruising [see *Warnings and Precautions* (5.4)].

#### Pregnancy and Breastfeeding

Advise patients to inform their doctor if they are or think they may be pregnant. Advise women of reproductive potential to avoid becoming pregnant while taking imatinib mesylate tablets. Female patients of reproductive potential taking imatinib mesylate tablets should use highly effective contraception during treatment and for fourteen days after stopping treatment with imatinib mesylate tablets [see *Use in Specific Populations* (8.3)]. Avoid breastfeeding during treatment and for 1 month after the last dose [see *Use in Specific Populations* (8.2)].

#### Drug Interactions

Imatinib mesylate tablets and certain other medicines, such as warfarin, erythromycin, and phenytoin, including over-the-counter medications, such as herbal products, can interact with each other. Advise patients to tell their doctor if they are taking or plan to take iron supplements. Avoid grapefruit juice and other foods known to inhibit CYP3A4 while taking imatinib mesylate tablets [see *Drug Interactions* (7)].

#### Pediatric

Advise patients that growth retardation has been reported in children and pre-adolescents receiving imatinib mesylate. The long-term effects of prolonged treatment with imatinib mesylate on growth in children are unknown. Therefore, closely monitor growth in children under imatinib mesylate treatment [see *Warnings and Precautions* (5.11)].

#### Driving and Using Machines

Advise patients that they may experience side effects, such as dizziness, blurred vision, or somnolence during treatment with imatinib mesylate. Therefore, caution patients about driving a car or operating machinery [see *Warnings and Precautions* (5.13)].

#### **APOTEX INC.**

#### **Imatinib Mesylate Tablets 100 mg and 400 mg**

---

#### **Manufactured by      Manufactured for**

|                  |                |
|------------------|----------------|
| Apotex Inc.      | Apotex Corp.   |
| Toronto, Ontario | Weston Florida |
| Canada M9L 1T9   | USA 33326      |

---

Revision: 25

#### **PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - 100 MG LABEL**

**APOTEX CORP.** NDC 60505-2900-9


#### **Imatinib Mesylate Tablets**

**Equivalent to 100 mg of imatinib**

100 mg

**Rx only**

90 Tablets



PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - 400 MG LABEL

APOTEX CORP. NDC 60505-2901-3


Imatinib Mesylate Tablets

Equivalent to 400 mg of imatinib

400 mg

Rx only

30 Tablets



IMATINIB MESYLATE

imatinib mesylate tablet, film coated

| Product Information                                               |                          |                    |                |
|-------------------------------------------------------------------|--------------------------|--------------------|----------------|
| Product Type                                                      | HUMAN PRESCRIPTION DRUG  | Item Code (Source) | NDC:60505-2900 |
| Route of Administration                                           | ORAL                     |                    |                |
|                                                                   |                          |                    |                |
| Active Ingredient/Active Moiety                                   |                          |                    |                |
| Ingredient Name                                                   |                          | Basis of Strength  | Strength       |
| IMATINIB MESYLATE (UNII: 8A1O1M485B) (Imatinib - UNII:BKJ8M8G5HI) |                          | Imatinib           | 100 mg         |
|                                                                   |                          |                    |                |
| Inactive Ingredients                                              |                          |                    |                |
| Ingredient Name                                                   |                          |                    | Strength       |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)                                |                          |                    |                |
| CROSPVIDONE (120 .MU.M) (UNII: 68401960MK)                        |                          |                    |                |
| MAGNESIUM STEARATE (UNII: 70097M6I30)                             |                          |                    |                |
| HYPPROMELLOSE 2910 (5 MPA.S) (UNII: R75537T0T4)                   |                          |                    |                |
| HYDROXYPROPYL CELLULOSE (90000 WAMW) (UNII: UKE75GEA7F)           |                          |                    |                |
| FERRIC OXIDE RED (UNII: 1K09F3G675)                               |                          |                    |                |
| FERRIC OXIDE YELLOW (UNII: EX438O2MRT)                            |                          |                    |                |
| POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)                       |                          |                    |                |
|                                                                   |                          |                    |                |
| Product Characteristics                                           |                          |                    |                |
| Color                                                             | ORANGE (Brownish Orange) | Score              | 2 pieces       |
| Shape                                                             | ROUND                    | Size               | 9mm            |
| Flavor                                                            |                          | Imprint Code       | IMA;100;APO    |
| Contains                                                          |                          |                    |                |

Packaging

| # | Item Code        | Package Description                                     | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:60505-2900-3 | 30 in 1 BOTTLE; Type 0: Not a Combination Product       | 08/05/2016           |                    |
| 2 | NDC:60505-2900-9 | 90 in 1 BOTTLE; Type 0: Not a Combination Product       | 08/05/2016           |                    |
| 3 | NDC:60505-2900-1 | 100 in 1 BOTTLE; Type 0: Not a Combination Product      | 08/05/2016           |                    |
| 4 | NDC:60505-2900-8 | 1000 in 1 BOTTLE; Type 0: Not a Combination Product     | 08/05/2016           |                    |
| 5 | NDC:60505-2900-0 | 10 in 1 CARTON                                          | 08/05/2016           |                    |
| 5 |                  | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|------------------------------------------|----------------------|--------------------|
| ANDA               | ANDA079179                               | 08/05/2016           |                    |

IMATINIB MESYLATE

imatinib mesylate tablet, film coated

Product Information

|                         |                         |                    |                |
|-------------------------|-------------------------|--------------------|----------------|
| Product Type            | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:60505-2901 |
| Route of Administration | ORAL                    |                    |                |

Active Ingredient/Active Moiety

| Ingredient Name                                                   | Basis of Strength | Strength |
|-------------------------------------------------------------------|-------------------|----------|
| IMATINIB MESYLATE (UNII: 8A1O1M485B) (Imatinib - UNII:BKJ8M8G5HI) | Imatinib          | 400 mg   |

Inactive Ingredients

| Ingredient Name                                         | Strength |
|---------------------------------------------------------|----------|
| SILICON DIOXIDE (UNII: ETJ7Z 6XBU4)                     |          |
| CROSPVIDONE (120 .MU.M) (UNII: 68401960MK)              |          |
| MAGNESIUM STEARATE (UNII: 70097M6I30)                   |          |
| HYPROMELLOSE 2910 (5 MPA.S) (UNII: R75537T0T4)          |          |
| HYDROXYPROPYL CELLULOSE (90000 WAMW) (UNII: UKE75GEA7F) |          |
| FERRIC OXIDE RED (UNII: 1K09F3G675)                     |          |
| FERRIC OXIDE YELLOW (UNII: EX438O2MRT)                  |          |
| POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)             |          |

Product Characteristics

|          |                          |              |             |
|----------|--------------------------|--------------|-------------|
| Color    | ORANGE (Brownish Orange) | Score        | 2 pieces    |
| Shape    | CAPSULE                  | Size         | 17mm        |
| Flavor   |                          | Imprint Code | IMA;400;APO |
| Contains |                          |              |             |

Packaging

| # | Item Code        | Package Description                                     | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:60505-2901-3 | 30 in 1 BOTTLE; Type 0: Not a Combination Product       | 08/05/2016           |                    |
| 2 | NDC:60505-2901-9 | 90 in 1 BOTTLE; Type 0: Not a Combination Product       | 08/05/2016           |                    |
| 3 | NDC:60505-2901-1 | 100 in 1 BOTTLE; Type 0: Not a Combination Product      | 08/05/2016           |                    |
| 4 | NDC:60505-2901-5 | 500 in 1 BOTTLE; Type 0: Not a Combination Product      | 08/05/2016           |                    |
| 5 | NDC:60505-2901-0 | 10 in 1 CARTON                                          | 08/05/2016           |                    |
| 5 |                  | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |

Marketing Information

| Marketing | Application Number or Monograph | Marketing Start | Marketing End |
|-----------|---------------------------------|-----------------|---------------|
|-----------|---------------------------------|-----------------|---------------|

| Category | Citation   | Date       | Date |
|----------|------------|------------|------|
| ANDA     | ANDA079179 | 08/05/2016 |      |

**Labeler -** Apotex Corp (845263701)

Revised: 1/2026

Apotex Corp