

RAPIDOX BASE- sodium chlorite solution, concentrate
DeLaval

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

RapidOx™ Base

Concentrated Disinfecting Teat Dip

Aids in reducing the spread of mastitis-causing organisms

ACTIVE INGREDIENTS:

Sodium chlorite 7.1%

KEEP OUT OF REACH OF CHILDREN

! DANGER

READ AND UNDERSTAND ALL INSTRUCTIONS BEFORE USING THIS PRODUCT

RapidOx is a three-part Chlorine Dioxide teat dip that must be used with the InstaMix G Emollient Concentrate. Recommended for pre/post milking applications and can be used to reduce the spread of mastitis causing organisms.

HAZARD STATEMENTS

Harmful in contact with skin. Causes severe skin burns and eye damage. May cause damage to organs through prolonged or repeated exposure: (Respiratory, Kidney, and Blood Systems). May cause damage to renal system (Kidneys), and Blood system through prolonged or repeated exposure. May be corrosive to metals. Wear protective gloves/protective clothing/eye protections/face protection. Do not breathe dust/fume/gas/mist/vapors/spray. Wash face, hands and any exposed skin thoroughly after handling. Keep only in original container.

FIRST AID

Immediately call a POISON CENTER or doctor/physician

If in Eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a **POISON CENTER** or doctor/physician.

If on Skin: Remove/Take off immediately all contaminated clothing. Rinse skin with water/shower. Wash contaminated clothing before reuse. Call a **POISON CENTER** or doctor/physician if you feel unwell.

If Inhaled: Remove victim to fresh air and keep at rest in a position comfortable for breathing. Immediately call a **POISON CENTER** or doctor/physician.

If Swallowed: Rinse mouth. Do NOT induce vomiting.

STORAGE AND DISPOSAL

THOROUGHLY RINSE ALL MEASURING DEVICES AND STORE OUT OF REACH OF CHILDREN. Avoid contamination of food or feed. Avoid freezing. Keep out of direct

sunlight. Keep away from strong bases. Keep out of the reach of children. Store in suitable labelled containers. Storage temperatures: 32°F to 86°F. Store locked up. **FOR PRODUCT SPILLS, ONLY WATER SHOULD BE USED FOR CLEANUP. IMMEDIATELY DILUTE THE MATERIAL WITH PLENTY OF WATER. THIS PRODUCT BECOMES A FIRE HAZARD IF ALLOWED TO DRY.** Ensure adequate ventilation, especially in confined areas. Wash hands thoroughly after handling. Do not mix with bleach or other chlorinated products - will cause chlorine gas. Dispose of contents/container to an approved waste disposal plant.

CHEMTREC EMERGENCY NUMBER 1-800-424-9300

Not for Human Use

Additional Information on Safety Data Sheet

DIRECTIONS

IMPORTANT THIS IS A THREE-PART CONCENTRATE - MUST BE MIXED PRIOR TO USE WITH INSTAMIX™ G EMOLLIENT CONCENTRATE.

CREATES CHLORINE DIOXIDE See Dilution Chart for mix ratios.

General Use

- Concentrated formula must be diluted into a ready-to-use product.
- Using a manual pump or automatic system, mix into a clean teat cup or container at least 15 minutes prior to use. Prepare only the quantity needed for one milking.
- If cow's teats are sore or chapped, consult a veterinarian before starting to use teat dip.
- If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.
- If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Pre-Dipping Use

- Perform forestripping and inspect teats for clinical mastitis.
- Pre-wash teats thoroughly just prior to milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels.
- Just before each cow is milked, dip, or spray each teat full-length into the teat dip cup containing product.
- Allow product to stand for 30 seconds.
- Wipe teats dry after dipping, using single-service towels to avoid contamination of milk

Post-Dipping Use

- Immediately after each milking, use product.
- Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.
- If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Undiluted product may undergo an increase in viscosity when exposed to cold temperatures. Pumps must be recalibrated in those conditions to maintain accurate

dosing. If product freezes, thaw it completely and mix properly before use. Consult your DeLaval representative for questions on specific applications.

DILUTION CHART

% Emollient	RapidOx Activator (gal/100)	RapidOx Base (gal/100)	Instamix™ G Conc (gal/100)	Emollient (gal/100)	Water (gal/100)
1%	2	2		2	94
2%	2	2		4	92
3%	2	2		5	91
4%	2	2		7	89
5%	2	2		9	87
6%	2	2		11	85
7%	2	2		12	84
8%	2	2		14	82
9%	2	2		16	80
10%	2	2		18	78
20%	2	2		36	60

Manufactured for: **DeLaval Inc.**

11100 N. Congress Avenue

Kansas City, MO 6153

Phone: 816-891-7700

Manufactured by: West Agro, Inc.

© 2025, Made in UA

Labels

Label 55 Gallons

DeLaval

4203-B1225

RapidOx™

Base

Concentrated Disinfecting Teat Dip
Aids in reducing the spread of mastitis-causing organisms

Sellador desinfectante concentrado
Ayuda a reducir la transmisión de microorganismos que causan mastitis

ACTIVE INGREDIENTS:
Sodium chlorite.....7.1%

4203
NDC NUMBER: 55756-203

KEEP OUT OF REACH OF CHILDREN
⚠ DANGER
MANTENER FUERA DEL ALCANCE DE LOS NIÑOS
⚠ PELIGRO

READ AND UNDERSTAND ALL INSTRUCTIONS BEFORE USING THIS PRODUCT
SI USTED NO PUEDE LEER ESTA ETIQUETA, POR FAVOR PIDA A ALGUIEN MAS QUE SE LA LEA ANTES DE USAR ESTE PRODUCTO

RapidOx is a three-part Chlorine Dioxide teat dip that must be used with the InstaMix™ G Emollient Concentrate. Recommended for pre/post milking applications and can be used to reduce the spread of mastitis causing organisms.

HAZARD STATEMENTS
Harmful in contact with skin. Causes severe skin burns and eye damage. May cause damage to organs through prolonged or repeated exposure. (Respiratory, Kidney, and Blood Systems). May cause damage to renal system (Kidney), and Blood system through prolonged or repeated exposure. May be corrosive to metals. Wear protective gloves/protective clothing/eye protection/face protection. Do not breathe dust/fume/gas/mist/vapors/spray. Wash face, hands and any exposed skin thoroughly after handling. Keep only in original container.

FIRST AID
If in **Eyes:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a **POISON CONTROL CENTER** or doctor/physician.
If on **Skin:** Remove/Take off immediately all contaminated clothing. Rinse skin with water/shower. Wash contaminated clothing before reuse. Call a **POISON CONTROL CENTER** or doctor/physician if you feel unwell.
If **Inhaled:** Remove victim to fresh air and keep at rest in a position comfortable for breathing. Immediately call a **POISON CONTROL CENTER** or doctor/physician.
If **Swallowed:** Rinse mouth. Do NOT induce vomiting.

STORAGE AND DISPOSAL
THOROUGHLY RINSE ALL MEASURING DEVICES AND STORE OUT OF REACH OF CHILDREN. Avoid contamination of food or feed. Avoid freezing. Keep out of direct sunlight. Keep away from strong bases. Keep out of the reach of children. Store in suitable labeled containers. Storage temperature: 32°F to 86°F. Store locked up. **FOR PRODUCT SPILLS, ONLY WATER SHOULD BE USED FOR CLEANUP. IMMEDIATELY DILUTE THE MATERIAL WITH PLENTY OF WATER. THIS PRODUCT BECOMES A FIRE HAZARD IF ALLOWED TO DRY.** Ensure adequate ventilation, especially in confined areas. Wash hands thoroughly after handling. Do not mix with bleach or other chlorinated products - will cause chlorine gas. Dispose of contents/container to an approved waste disposal plant.

CHEMTREC EMERGENCY NUMBER 1-800-424-9300
Not For Human Use
Additional Information on Safety Data Sheet

DILUTION CHART

% Emollient	RapidOx Activator (gal/100 gal)	RapidOx Base (gal/100)	InstaMix™ G Emollient Conc (gal/100)	Water (gal/100)
1%	2	2	2	94
2%	2	2	4	88
3%	2	2	5	89
4%	2	2	7	89
5%	2	2	9	87
6%	2	2	11	85
7%	2	2	12	84
8%	2	2	14	82
9%	2	2	16	80
10%	2	2	18	78
20%	2	2	36	60

DIRECTIONS
IMPORTANT: THIS IS A THREE-PART CONCENTRATE - MUST BE MIXED PRIOR TO USE WITH INSTAMIX™ G EMOLLIENT CONCENTRATE. CREATES CHLORINE DIOXIDE. See Dilution Chart for mix ratios.

General Use

- Concentrated formula must be diluted into a ready-to-use product.
- Using a manual pump or automatic system, mix into a clean teat cup or container at least 15 minutes prior to use. Prepare only the quantity needed for one milking.
- If cow's teats are sore or chapped, consult a veterinarian before starting to use teat dip.
- If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.
- If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Pre-Clipping Use

- Perform footstripping and inspect teats for clinical mastitis.
- Pre-wash teats thoroughly just prior to milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels.
- Just before each cow is milked, dip, or spray each teat full-length into the teat dip cup containing product.
- Allow product to stand for 30 seconds.
- Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post-Clipping Use

- Immediately after each milking, use product.
- Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.
- If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Undiluted product may undergo an increase in viscosity when exposed to cold temperatures. Pumps must be recalibrated in those conditions to maintain accurate dosing. If product freezes, thaw it completely and mix properly before use. Consult your DeLaval representative for questions on specific applications.

Manufactured by: **DeLaval Inc.**
1100 N. Congress avenue
Kansas City, MO 64153
Phone: 816-891-7700
Manufactured by: West Agro, Inc.
© 2025, Made in USA

NET CONTENTS: 55 US GAL. (208.2 L)
LOT NO.:
EXP. DATE:

UN1908
Chlorite solution

Label 275 Gallons

DeLaval

4203-B1225

RapidOx™

Base

Concentrated Disinfecting Teat Dip
Aids in reducing the spread of mastitis-causing organisms

Sellador desinfectante concentrado
Ayuda a reducir la transmisión de microorganismos que causan mastitis

ACTIVE INGREDIENTS:
Sodium chlorite.....7.1%

4203
NDC NUMBER: 55756-203

KEEP OUT OF REACH OF CHILDREN
⚠ DANGER
MANTENER FUERA DEL ALCANCE DE LOS NIÑOS
⚠ PELIGRO

READ AND UNDERSTAND ALL INSTRUCTIONS BEFORE USING THIS PRODUCT
SI USTED NO PUEDE LEER ESTA ETIQUETA, POR FAVOR PIDA A ALGUIEN MAS QUE SE LA LEA ANTES DE USAR ESTE PRODUCTO

RapidOx is a three-part Chlorine Dioxide teat dip that must be used with the InstaMix™ G Emollient Concentrate. Recommended for pre/post milking applications and can be used to reduce the spread of mastitis causing organisms.

HAZARD STATEMENTS
Harmful in contact with skin. Causes severe skin burns and eye damage. May cause damage to organs through prolonged or repeated exposure. (Respiratory, Kidney, and Blood Systems). May cause damage to renal system (Kidney), and Blood system through prolonged or repeated exposure. May be corrosive to metals. Wear protective gloves/protective clothing/eye protection/face protection. Do not breathe dust/fume/gas/mist/vapors/spray. Wash face, hands and any exposed skin thoroughly after handling. Keep only in original container.

FIRST AID
If in **Eyes:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a **POISON CONTROL CENTER** or doctor/physician.
If on **Skin:** Remove/Take off immediately all contaminated clothing. Rinse skin with water/shower. Wash contaminated clothing before reuse. Call a **POISON CONTROL CENTER** or doctor/physician if you feel unwell.
If **Inhaled:** Remove victim to fresh air and keep at rest in a position comfortable for breathing. Immediately call a **POISON CONTROL CENTER** or doctor/physician.
If **Swallowed:** Rinse mouth. Do NOT induce vomiting.

STORAGE AND DISPOSAL
THOROUGHLY RINSE ALL MEASURING DEVICES AND STORE OUT OF REACH OF CHILDREN. Avoid contamination of food or feed. Avoid freezing. Keep out of direct sunlight. Keep away from strong bases. Keep out of the reach of children. Store in suitable labeled containers. Storage temperature: 32°F to 86°F. Store locked up. **FOR PRODUCT SPILLS, ONLY WATER SHOULD BE USED FOR CLEANUP. IMMEDIATELY DILUTE THE MATERIAL WITH PLENTY OF WATER. THIS PRODUCT BECOMES A FIRE HAZARD IF ALLOWED TO DRY.** Ensure adequate ventilation, especially in confined areas. Wash hands thoroughly after handling. Do not mix with bleach or other chlorinated products - will cause chlorine gas. Dispose of contents/container to an approved waste disposal plant.

CHEMTREC EMERGENCY NUMBER 1-800-424-9300
Not For Human Use
Additional Information on Safety Data Sheet

DILUTION CHART

% Emollient	RapidOx Activator (gal/100 gal)	RapidOx Base (gal/100)	InstaMix™ G Emollient Conc (gal/100)	Water (gal/100)
1%	2	2	2	94
2%	2	2	4	88
3%	2	2	5	89
4%	2	2	7	89
5%	2	2	9	87
6%	2	2	11	85
7%	2	2	12	84
8%	2	2	14	82
9%	2	2	16	80
10%	2	2	18	78
20%	2	2	36	60

DIRECTIONS
IMPORTANT: THIS IS A THREE-PART CONCENTRATE - MUST BE MIXED PRIOR TO USE WITH INSTAMIX™ G EMOLLIENT CONCENTRATE. CREATES CHLORINE DIOXIDE. See Dilution Chart for mix ratios.

General Use

- Concentrated formula must be diluted into a ready-to-use product.
- Using a manual pump or automatic system, mix into a clean teat cup or container at least 15 minutes prior to use. Prepare only the quantity needed for one milking.
- If cow's teats are sore or chapped, consult a veterinarian before starting to use teat dip.
- If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.
- If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Pre-Clipping Use

- Perform footstripping and inspect teats for clinical mastitis.
- Pre-wash teats thoroughly just prior to milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels.
- Just before each cow is milked, dip, or spray each teat full-length into the teat dip cup containing product.
- Allow product to stand for 30 seconds.
- Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post-Clipping Use

- Immediately after each milking, use product.
- Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.
- If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Undiluted product may undergo an increase in viscosity when exposed to cold temperatures. Pumps must be recalibrated in those conditions to maintain accurate dosing. If product freezes, thaw it completely and mix properly before use. Consult your DeLaval representative for questions on specific applications.

Manufactured by: **DeLaval Inc.**
1100 N. Congress avenue
Kansas City, MO 64153
Phone: 816-891-7700
Manufactured by: West Agro, Inc.
© 2025, Made in USA

NET CONTENTS: 55 US GAL. (208.2 L)
LOT NO.:
EXP. DATE:

UN1908
Chlorite solution

RAPIDOX BASE
sodium chlorite solution, concentrate

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:55756-203
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORITE (UNII: G538EBV4VF) (CHLORITE ION - UNII:Z63H374SB6)	SODIUM CHLORITE	71 g in 1 L

Inactive Ingredients

Ingredient Name	Strength
TETRASODIUM EDTA (UNII: MP1J8420LU)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55756-203-01	208 L in 1 DRUM		
2	NDC:55756-203-27	1041 L in 1 TANK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/19/2025	

Labeler - DeLaval (173704867)

Establishment

Name	Address	ID/FEI	Business Operations
West Agro, Inc.		147528723	api manufacture, manufacture, pack, label, analysis

Establishment

Name	Address	ID/FEI	Business Operations
West Agro, Inc.		095906480	api manufacture, manufacture, pack, label, analysis