

GOODYS MAX- acetaminophen, aspirin, and caffeine powder
Medtech Products Inc.

Goodys Max Relief Tropical Citrus

GOODY'S

Drug Facts

Active ingredients (in each powder)

Acetaminophen 500 mg

Aspirin (NSAID*) 500 mg

Caffeine 130 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Pain reliever

Pain reliever aid

Uses

Temporarily relieves minor aches and pains due to:

- headache
- minor arthritis pain
- a cold
- backaches
- toothache
- muscular aches
- menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause severe allergic reaction which may include:

- hives

- facial swelling
- asthma (wheezing)
- shock.

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 packs in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: This product contains caffeine. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, gout, or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - any new symptoms appear
 - ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed** (see overdose warning)
- **adults and children 12 years and over:** open 1 pack, place powder on tongue every 6 hours, while symptoms persist. Drink a full glass of water with each dose or may stir powder into a glass of water or other liquid.
- do not take more than 4 packs in 24 hours unless directed by a doctor.
- **children under 12 years of age:** ask a doctor.

Other information

Store below 25°C (77°F)

Inactive ingredients

colloidal silicon dioxide, flavors, guar gum, mannitol, microcrystalline cellulose, sucralose

Questions?

1-866-255-5197 weekdays or goodyspowder.com

Principal Display Panel

Goody's® for adults

MAX PAIN RELIEF *

TRIPLE ACTION

POWDER

ACETAMINOPHEN

ASPIRIN (NSAID) / CAFFEINE

PAIN RELIEVER / PAIN RELIEVER AID

TROPICAL CITRUS

16 STICK PACKS

1 PACK = 1 DOSE



Drug Facts		1	Drug Facts (continued)	2
Active ingredients	Purposes			
Acetaminophen 325 mg	Relieves minor aches and pains due to: * headache * minor arthritis pain * cold * backache * toothache * muscular aches * menstrual cramps			
Aspirin (NSAID) 300 mg	Relieves minor aches and pains due to: * headache * minor arthritis pain * cold * backache * toothache * muscular aches * menstrual cramps			
Caffeine 100 mg				
Warnings: Children and teenagers who have or may have had anemia or Reye's syndrome should not use this product. When using this product, if you have a fever and a rash occurs, consult a doctor because these symptoms can be early signs of Reye's syndrome, a rare but serious disease.				
Allergy alert: Aspirin may cause severe allergic reactions which may include: hives, facial swelling, asthma (breathing困难) and/or Acetaminophen may cause severe skin reactions. Symptoms may include: * skin reddening and/or rash. If skin reaction occurs, stop use and seek medical help right away.				
TAMPER EVIDENT: DO NOT USE IF PACK IS DAMAGED OR OPEN.				
KEEP CARTON FOR FUTURE REFERENCE.				
* MAX PAIN RELIEVERS WITHOUT A PRESCRIPTION				
OPEN FLAP FOR DRUG FACTS IN NUMERICAL ORDER.				
TO OPEN CHILD-RESISTANT PACKAGE				
1. SHAKE PACK as powder surface on bottom.				
2. Push and turn at either arrow or tear scores.				

Drug Facts (continued)	3	Drug Facts (continued)	4
<p>Ask a doctor before use if:</p> <ul style="list-style-type: none"> you have liver disease you are taking aspirin or any other medicine that may affect your liver you have high blood pressure, heart disease, liver disease, or kidney disease you are taking a diuretic you have asthma <p>Ask a doctor or pharmacist before use if you are taking:</p> <ul style="list-style-type: none"> a prescription drug for diabetes, gout, or asthma any medicine that may affect your liver any medicine that may affect your heart any medicine or supplement more than 12 days any cold medicine or cough medicine any new symptom appears any ringing in the ears or a loss of hearing occurs <p>If pregnant or breast feeding, ask a health professional before use. It is especially important not to use aspirin or any medicine that may affect your liver or heart. If you are pregnant or nursing, it is extremely important to do as the doctor directs. It is important to tell your doctor if you are pregnant or nursing, or if you have any of the above conditions or complications during delivery.</p>		<p>Keep out of reach of children.</p> <p>Do not take more than the recommended dose. Taking more than the recommended dose can cause you to feel sick with problems. In case of overdose, get medical help or contact a Poison Control Center 1-800-222-1222 right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions:</p> <ul style="list-style-type: none"> do not take more than directed (see overuse warning) adults and children 12 years and over: open 1 pack, pour powder on tongue every 8 hours, while on resting period. Drink a full glass of water with each dose or with the next dose if it is greater than 4 packs in 24 hours unless directed by a doctor children under 12 years of age: use a doctor <p>Other Information: Storage below 20°C (68°F)</p> <p>Active Ingredients: acetaminophen, aspirin, and caffeine, m</p> <p>Questions? 1-866-255-5187 weekdays or goodeysonline.com</p>	
Drug Facts (continued)	3	Drug Facts (continued)	4
<p>Ask a doctor before use if:</p> <ul style="list-style-type: none"> you have liver disease you are taking aspirin or any other medicine that may affect your liver you have high blood pressure, heart disease, liver disease, or kidney disease you are taking a diuretic you have asthma <p>Ask a doctor or pharmacist before use if you are taking:</p> <ul style="list-style-type: none"> a prescription drug for diabetes, gout, or asthma any medicine that may affect your liver any medicine that may affect your heart any medicine or supplement more than 12 days any cold medicine or cough medicine any new symptom appears any ringing in the ears or a loss of hearing occurs <p>If pregnant or breast feeding, ask a health professional before use. It is especially important not to use aspirin or any medicine that may affect your liver or heart. If you are pregnant or nursing, it is extremely important to do as the doctor directs. It is important to tell your doctor if you are pregnant or nursing, or if you have any of the above conditions or complications during delivery.</p>		<p>Keep out of reach of children.</p> <p>Do not take more than the recommended dose. Taking more than the recommended dose can cause you to feel sick with problems. In case of overdose, get medical help or contact a Poison Control Center 1-800-222-1222 right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions:</p> <ul style="list-style-type: none"> do not take more than directed (see overuse warning) adults and children 12 years and over: open 1 pack, pour powder on tongue every 8 hours, while on resting period. Drink a full glass of water with each dose or with the next dose if it is greater than 4 packs in 24 hours unless directed by a doctor children under 12 years of age: use a doctor <p>Other Information: Storage below 20°C (68°F)</p> <p>Active Ingredients: acetaminophen, aspirin, and caffeine, m</p> <p>Questions? 1-866-255-5187 weekdays or goodeysonline.com</p>	
Drug Facts (continued)	3	Drug Facts (continued)	4

Dist. by Meditch Products Inc.
Tarrytown, NY 10591
A Prestige Consumer Healthcare company
© 2015 Trade dress is owned by Meditch
Products Inc. All rights reserved.
GO/5187101



GOODYS MAX

acetaminophen, aspirin, and caffeine powder

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63029-668
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362091TL9D) (ACETAMINOPHEN - UNII:362091TL9D)	ACETAMINOPHEN	500 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	130 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
GUAR GUM (UNII: E89I1637KE)	
MANNITOL (UNII: 30WL53L36A)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor	CITRUS (Tropical Citrus)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63029-668-16	16 in 1 CARTON; Type 0: Not a Combination Product	02/02/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/02/2026	

Labeler - Medtech Products Inc. (122715688)

Revised: 2/2026

Medtech Products Inc.