

PAIN RELIEVING PATCH- lidocaine 4% patch simple diagnostics

Pain Relieving Patch

Drug Facts

LIDOCAINE PATCH 4%

Active Ingredient

Lidocaine 4%

Purpose

Topical anesthetic

USE

For the temporary relief of pain

Warnings

For external use only

Do not use more than 1 patch on your body at a time on cut, irritated, or swollen skin on puncture wounds for more than one week without consulting a doctor

When using this product

Use only as directed. Read and follow all directions and warnings on this carton. do not allow contact with the eyes do not bandage tightly or apply local heat (such as heating pads) to the area of use do not use at the same time as other topical analgesics dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

DO NOT USE:

- More than 1 patch at a time
- On wounds or damaged skin
- For more than one week without consulting a doctor
- If you are allergic to any ingredients of this product

STOP USE AND ASK A DOCTOR IF:

- Localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- Condition worsen
- Symptoms persist for more than 7 days
- Symptoms clear up and occur again within a few days

IF PREGNANT OR BREAST-FEEDING:

Ask a health professional before use

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away(1-800-222-1222).

Directions

Adults and children 12 years of age and older:

Wash and dry the affected area.

Peel off the backing film from one side of the patch carefully

Apply the exposed portion of the patch to the affected area.

Gently press the patch with your hand to fit the affected area completely.

Do not exercise vigorously during use to prevent the patch from falling off.

Use only 1 patch for a maximum of 8 hours.

Children under 12 years of age: Ask a doctor before use.

QUESTIONS OR COMMENTS:

- Customerservice@simplediagnosics.com
- 877-342-2385

Other information

- Store at room temperature 15°-30°C (59°-86°F)
- Store in a cool dry place away from direct sunlight

Inactive ingredient

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylic Acid, Povidone K90, Propylparaben, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Water.

PACKAGE LABEL. PRINCIPAL DISPLAY PANEL:

MAXIMUM STRENGTH NDC 62379-201-02

UNSCENTED

MEDICATED

Pharmacist Choice

Pain Relief Patch

LIDOCAINE 4%
Pain Relieving Ointment on a Breathable Adhesive Pad

- ✓ **Fast Acting**
Temporarily relieves minor pain & desensitize aggravated nerves
- ✓ **Flexible Fabric**
Flexible fabric patch, easy to apply and remove
- ✓ **No Mess**
No-Mess, single-use application which stays put

NUMBS AWAY PAIN

5 Patches
3.94" x 5.5" (10cm x 14cm) each

LIDOCAINE 4%
Pain Relieving Ointment on a Breathable Adhesive Pad

Pharmacist Choice

Drug Facts

Active ingredient Lidocaine 4%	Purpose Topical anesthetic
--	--------------------------------------

Use
For the temporary relief of pain

Warnings
For external use only

Do not use more than 1 patch on your body at a time on cut, irritated, or swollen skin on puncture wounds for more than one week without consulting a doctor

When using this product

- use only as directed. Read and follow all directions and warnings on this carton. do not allow contact with the eyes
- do not bandage tightly or apply local heat (such as heating pads) to the area of use do not use at the same time as other topical analgesics
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests.

Stop use and ask a doctor if

- condition worsens
- symptoms persist for more than 7 days or clear up and occur again within a few days
- redness or irritation develops
- you experience signs of skin injury, such as pain, swelling, or blistering where applied

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
Adults and children 12 years of age and older:

- Wash and dry the affected area.
- Peel off the backing film from one side of the patch carefully.
- Apply the exposed portion of the patch to the affected area.
- Gently press the patch with your hand to fit the affected area completely.
- Do not exercise vigorously during use to prevent the patch from falling off.
- Use only 1 patch for a maximum of 8 hours.

Children under 12 years of age: Ask a doctor before use.

Other information

- store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

Inactive ingredients
Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Potassium Hydroxide, Polyacrylate, Tartaric Acid, Titanium Dioxide, Water.

Questions or Comments? Contact customerservice@simplediagnostics.com

Child Resistant Packaging NDC 62379-201-02

LOT: [] BATCH: [] USE BY: []

Distributed by: **Simple Diagnostics**
PO Box 128
Williston Park, NY 11596, USA
Tel: 877-342-2385
www.SimpleDiagnostics.com
Made in China

8 50002 86069 6

Child Resistant Packaging NDC 62379-201-03

Pharmacist Choice

Pain Relieving Patch

LIDOCAINE 4%
Pain Relieving Ointment on a Breathable Adhesive Pad

UNSCENTED **MAXIMUM STRENGTH** **MEDICATED**

30 Patches
3.94" x 5.5" (10cm x 14cm) each

Fast Acting
Temporarily relieves minor pain

Flexible Fabric
Flexible fabric patch, easy to apply and remove

No Mess
No-Mess, single-use application which stays put

Distributed by: **Simple Diagnostics**
PO Box 128
Williston Park, NY 11596, USA
Tel: 877-342-2385
www.SimpleDiagnostics.com
Made in China

NDC 62379-201-03

LOT: [] BATCH: [] USE BY: []

Drug Facts

Active ingredient Lidocaine 4%	Purpose Topical anesthetic
--	--------------------------------------

Use
For the temporary relief of pain

Warnings
For external use only

Do not use more than 1 patch on your body at a time on cut, irritated, or swollen skin on puncture wounds for more than one week without consulting a doctor

When using this product

- use only as directed. Read and follow all directions and warnings on this carton. do not allow contact with the eyes
- do not bandage tightly or apply local heat (such as heating pads) to the area of use do not use at the same time as other topical analgesics
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests.

Stop use and ask a doctor if

- condition worsens
- symptoms persist for more than 7 days or clear up and occur again within a few days
- redness or irritation develops
- you experience signs of skin injury, such as pain, swelling, or blistering where applied

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Child Resistant Packaging

Pharmacist Choice

Pain Relief Patch

30 Patches
3.94" x 5.5" (10cm x 14cm) each

Drug Facts (continued)

Directions
Adults and children 12 years of age and older:

- Wash and dry the affected area.
- Peel off the backing film from one side of the patch carefully.
- Apply the exposed portion of the patch to the affected area.
- Gently press the patch with your hand to fit the affected area completely.
- Do not exercise vigorously during use to prevent the patch from falling off.
- Use only 1 patch for a maximum of 8 hours.

Children under 12 years of age: Ask a doctor before use.

Other information

- store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

Inactive ingredients
Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylate, Potassium Hydroxide, Propylparaben, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide.

Questions or Comments? Contact customerservice@simplediagnostics.com

CUT HERE

NDC 62379-201-01

Pharmacist Choice™

Pain Relief Patch

APPLY FOR UP TO 8 HOURS

LIDOCAINE 4%

Pain Relieving Ointment on a Breathable Adhesive Pad

MAXIMUM STRENGTH

NUMBS AWAY PAIN

- ✓ Fast Acting
- ✓ Flexible Fabric
- ✓ Fragrance-Free

Contains **1 Patch**
3.94" x 5.5"
(10cm x 14cm) each

Drug Facts

Active Ingredient	Purpose
Lidocaine 4%	Topical anesthetic

Use
For the temporary relief of pain

Warnings
For external use only

Do not use ■ more than 1 patch on your body at a time ■ on cut, irritated, or swollen skin ■ on puncture wounds
■ for more than one week without consulting a doctor

When using this product
■ use only as directed. Read and follow all directions and warnings on this carton. ■ do not allow contact with the eyes
■ do not bandage tightly or apply local heat (such as heating pads) to the area of use ■ do not use at the same time as other topical analgesics ■ dispose of used patch in manner that always keeps product away from children and pets.
Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests.

Stop use and ask a doctor if
■ condition worsens ■ symptoms persist for more than 7 days or clear up and occur again within a few days
■ redness or irritation develops ■ you experience signs of skin injury, such as pain, swelling, or blistering where applied

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
Adults and children 12 years of age and older:
■ Wash and dry the affected area.
■ Peel off the backing film from one side of the patch carefully.
■ Apply the exposed portion of the patch to the affected area.
■ Gently press the patch with your hand to fit the affected area completely.
■ Do not exercise vigorously during use to prevent the patch from falling off.
■ Use only 1 patch for a maximum of 8 hours.
Children under 12 years of age: Ask a doctor before use.

Other information
■ store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

Inactive Ingredients
Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylic Acid, Povidone K90, Propylparaben, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Water.

Questions or Comments? Contact customerservice@simplediagnostics.com

Child-Resistant Packaging Lot: Exp:

Distributed by:
Simple Diagnostics
PO Box 128
Williston Park, NY 11596, USA
Tel: 877-342-2385
www.SimpleDiagnostics.com
Made in China



20850002860690

PAIN RELIEVING PATCH

lidocaine 4% patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62379-201
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.4 g

Inactive Ingredients

Ingredient Name	Strength
KAOLIN (UNII: 24H4NWX5CO)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JNI2J)	
POVIDONE K90 (UNII: RDH86HJV5Z)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
DIHYDROXYALUMINUM AMINOACETATE ANHYDROUS (UNII: 1K713C615K)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)

TARTARIC ACID (UNII: W4888I119H)

WATER (UNII: 059QF0KO0R)

GLYCERIN (UNII: PDC6A3C0OX)

POLYSORBATE 80 (UNII: 6OZP39ZG8H)

Product Characteristics

Color		Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62379-201-03	30 in 1 CASE	06/01/2023	
1	NDC:62379-201-02	5 in 1 BOX		
1	NDC:62379-201-01	1 in 1 BAG; Type 0: Not a Combination Product		

NUMBS AWAY PAIN NDC 62379-0201-03

Pharmacist Choice Pain Relief Patch

Pain Relief Patch NOT FOR USE FOR 8 HOURS

LIDOCAINE 4%

Pain Relieving Ointment on a Breathable Adhesive Pad

UNSCENTED MEDICATED

MAXIMUM STRENGTH

30 Patches
3.94" x 5.5" (10cm x 14cm) each

Drug Facts

Active ingredient: Lidocaine HCl. **Proprietary Excipients:**

Use: For the temporary relief of pain.

Warnings: For external use only. Do not use if you have had a patch on your body at a time when you had a heart attack or unstable angina. Do not use on wounds. Do not use on children under 12 years of age. Do not use on children under 12 years of age. Do not use on children under 12 years of age. Do not use on children under 12 years of age.

Directions: Adults and children 12 years of age and older: Apply the patch to the affected area. Apply the patch to the affected area. Apply the patch to the affected area. Apply the patch to the affected area.

Other information: Contains 30 patches. Each patch contains 10 mg of lidocaine. Each patch contains 10 mg of lidocaine.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. For more information, call 1-800-368-9773.

Contains 30 Patches

Simple Diagnostics
100 Willow Park, NY 11994, USA
Tel: 877-643-5253
www.simplediagnostics.com
Patch in blister

NDC 62379-0201-03

Pharmacist Choice

Pain Relief Patch NOT FOR USE FOR 8 HOURS

LIDOCAINE 4%

Pain Relieving Ointment on a Breathable Adhesive Pad

UNSCENTED MEDICATED

MAXIMUM STRENGTH

30 Patches
3.94" x 5.5" (10cm x 14cm) each

Pharmacist Choice

Pain Relief Patch NOT FOR USE FOR 8 HOURS

LIDOCAINE 4%

Pain Relieving Ointment on a Breathable Adhesive Pad

- ✓ **Fast Acting**
Temporarily relieves minor pain & desensitize aggravated nerves
- ✓ **Flexible Fabric**
Flexible fabric patch, easy to apply and remove
- ✓ **No Mess**
No-Mess, single-use application which stays put

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	06/01/2023	

Labeler - simple diagnostics (004135503)

Revised: 2/2025

simple diagnostics