

TOPCARE CLEARLAX- polyethylene glycol 3350 powder, for solution
Topco Associates LLC

Topco Associates LLC. ClearLax® Drug Facts

Active ingredient (in each dose)

Polyethylene Glycol 3350, 17 g (cap filled to line)

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use

if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

taking a prescription drug

When using this product

you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - use once a day
 - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficulty swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- tamper -evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients

natural and artificial orange flavor, sucralose

Questions or comments?

1-888-423-0139

Package/Label Principal Display Panel

TopCare® health

COMPARE TO MIRALAX® ACTIVE INGREDIENT

POWDER

GRIT FREE

ClearLax®

POLYETHYLENE GLYCOL 3350 POWDER FOR SOLUTION

OSMOTIC LAXATIVE

- Relieves Occasional Constipation (Irregularity)
- Softens Stool
- Sugar-free
- Dye-free

ORANGE FLAVORED POWDER

14 ONCE-DAILY DOSES

NET WT 8.3 OZ (238 g)

NDC 76162-119-02

+TopCare[®]
health

COMPARE TO MIRALAX[®]
ACTIVE INGREDIENT*

**POWDER
GRIT FREE**

ClearLax[®]

POLYETHYLENE GLYCOL 3350 POWDER FOR SOLUTION
OSMOTIC LAXATIVE

- **Relieves Occasional Constipation (Irregularity)**
- **Softens Stool**
- **Sugar-free**
- **Dye-free**



ORANGE
FLAVORED
POWDER

14 ONCE-DAILY DOSES

NET WT 8.3 OZ (238 g)

: 7U9E1 88 F1

Peel Here

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.
Dissolves in Any Beverage

Drug Facts

Active ingredient (in each dose) **Purpose**
Polyethylene Glycol 3350,
17 g (cap filled to line).....Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days


Warnings
Allergy alert: Do not use if you are allergic to polyethylene glycol
Do not use if you have kidney disease, except under the advice and supervision of a doctor
Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug
When using this product you may have loose, watery, more frequent stools
Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of MiraLAX®.



ADHESIVE AREA • NO COPY

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - use once a day
 - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficulty swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

Other information ■ store at 20°-25°C (68°-77°F)
■ tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients natural and artificial orange flavor, sucralose

Questions or comments? 1-888-423-0139

DISTRIBUTED BY TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007
©TOPCO PERA1023 QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcarebrand.com
Visit here for more information:
http://topbrnds.com/4903bc

GLUTEN FREE
QUALITY GUARANTEED
Patent Pending

| TOPCARE CLEARLAX | | | |
|--|----------------|---------------------------|---------------|
| polyethylene glycol 3350 powder, for solution | | | |
| Product Information | | | |
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:76162-119 |
| Route of Administration | ORAL | | |
| Active Ingredient/Active Moiety | | | |
| Ingredient Name | | Basis of Strength | Strength |
| POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P) | | POLYETHYLENE GLYCOL 3350 | 17 g in 17 g |
| Inactive Ingredients | | | |
| Ingredient Name | | Strength | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | | | |
| Product Characteristics | | | |
| Color | WHITE | Score | |
| Shape | | Size | |
| Flavor | ORANGE | Imprint Code | |

Contains**Packaging**

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:76162-119-02 | 238 g in 1 BOTTLE; Type 0: Not a Combination Product | 01/30/2024 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA090685 | 01/30/2024 | |

Labeler - Topco Associates LLC (006935977)

Revised: 2/2024

Topco Associates LLC