

**KAOPECTATE PEPPERMINT FLAVOR ANTI DIARRHEAL- bismuth  
subsalylate liquid  
Kramer Laboratories**

-----  
**Drug Facts**

<b>Active ingredient (per 15 mL)</b>	<b>Purposes</b>
Bismuth subsalylate 262 mg	Anti-diarrheal Upset stomach reliever

**Uses** relieves:

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - belching
  - fullness

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy Alert: Contains salicylate.**

**Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- a sodium-restricted diet
- fever

- mucus in the stool

**Ask a doctor or pharmacist before use if you are** taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### ***Directions***

- **shake well immediately before each use**
- only use pre-measured dose cup
- adults and children 12 years of age and older:
  - 1 dose (30 mL) every 1/2 hour to 1 hour as needed.
  - do not exceed 8 doses (240 mL) in 24 hours
  - use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### ***Other Information***

- **each 15 mL dose cup contains:** sodium 6 mg and **salicylate 135mg**
- store at room temperature 20°-25° C (68°-77° F)

### ***Inactive Ingredients***

carboxymethylcellulose sodium, FD&C Red No. 40, flavor microcrystalline cellulose, purified water, sodium salicylate, sorbic acid, sucrose, xanthan gum

### **Principal Display Panel**

***NEW IMPROVED TASTE!***

**Kaopectate®**

*Bismuth Subsalicylate 262 MG*

• *Anti-Diarrheal* • *Upset Stomach Reliever*

Diarrhea &  
Upset Stomach

- ✓ Begins controlling symptoms from the first dose
- ✓ Quickly relieves urgency, gas, and cramping
- ✓ Effective on diarrhea from bacteria, viruses, and other causes

11 fl oz (325mL)

**Peppermint Flavor**

681PPT11KA0 F



# Kaopectate Peppermint 11 oz

**Do not use if inner seal is broken or missing**

Kramer Laboratories  
Bridgewater, NJ 08807  
1-800-824-4894

Lot:

Exp:

**PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION**

681PPT11KA0LB

**Kaopectate Peppermint 11oz**

**Drug Facts**

Active ingredient (per 15 mL)	Purposes
Bismuth subsalicylate 262 mg	Anti-diarrheal Upset stomach reliever

**Uses** relieves: ■ traveler's diarrhea ■ diarrhea  
■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy Alert: Contains salicylate.**

**Do not take if you are**  
■ allergic to salicylates (including aspirin)  
■ taking other salicylate products

Do not use if inner seal is broken or missing

Kramer Laboratories  
Bridgewater, NJ 08807  
1-800-824-4894

Lot: \_\_\_\_\_  
Exp: \_\_\_\_\_

**PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION**



53076 19964 2



681PPT11KA0LB

**Drug Facts (continued)**

**Do not use if you have**  
■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have**  
■ a sodium-restricted diet ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for** ■ diabetes ■ gout ■ arthritis ■ anticoagulation (thinning the blood)

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**  
■ symptoms get worse  
■ ringing in the ears or loss of hearing occurs  
■ diarrhea lasts more than 2 days

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ shake well immediately before each use  
■ only use pre-measured dose cup  
■ adults and children 12 years of age and older:  
■ 1 dose (30 mL) every 1/2 hour to 1 hour as needed.  
■ do not exceed 8 doses (240 mL) in 24 hours  
■ use until diarrhea stops but not more than 2 days

**Drug Facts (continued)**

■ children under 12 years: ask a doctor  
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**  
■ each 15 mL dose cup contains: sodium 6 mg and salicylate 135mg  
■ store at room temperature 20°-25° C (68°-77° F)

**Inactive ingredients**  
carboxymethylcellulose sodium, FD&C Red #40, flavor, microcrystalline cellulose, purified water, sodium salicylate, sorbic acid, sucrose, xanthan gum

# KAOPECTATE PEPPERMINT FLAVOR ANTI DIARRHEAL

bismuth subsalicylate liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55505-199
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD18YE)	BISMUTH SUBSALICYLATE	262 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>Carboxymethylcellulose Sodium, Unspecified</b> (UNII: K679OBS311)	
<b>Fd&amp;C Red No. 40</b> (UNII: WZB9127XOA)	
<b>Microcrystalline Cellulose</b> (UNII: OP1R32D61U)	
<b>SODIUM SALICYLATE</b> (UNII: WQ1H85SYP)	
<b>SORBIC ACID</b> (UNII: X045WJ989B)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	PEPPERMINT	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55505-199-36	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2020	
2	NDC:55505-199-64	325 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2022	08/01/2025

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	10/01/2020	

