

## **EVAXO 3% HYDROGEN PEROXIDE- hydrogen peroxide liquid**

**Evaxo International Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **EVAXO 3% Hydrogen Peroxide**

#### ***Drug Facts***

##### **Active ingredient**

Hydrogen peroxide 3%

##### **Purpose**

first aid antiseptic/ oral debriding agent

**Uses** \*First aid to help prevent the risk of infection in minor cuts, scrapes, and burns. \*Aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth.

##### **Warnings**

**For external use only.**

**Do not use** \*in the eyes or apply over large areas of the body. \*longer than one week.

**Ask a doctor before use if you have** deep or puncture wounds, animal bites, or serious burns.

**Stop use and ask a doctor if** \*swelling, rash, or fever develops. \*the condition persists or gets worse. \*sore mouth symptoms do not improve in 7 days. \*irritation, pain or redness persists or worsens.

**Keep out of reach of children.** If swallowed, get medical help or contact poison control center right away.

**Directions** For use as a first aid antiseptic: \*Clean affected area. \*if bandaged, let dry first. \*apply small amount of product on affected area 1-3 times a day. \*may be covered with a sterile bandage. **For use as an oral debriding agent (oral rinse): \*adults and children 2 years of age and over: \*mix with an equal amount of water. \*swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. \*use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor. \*children under 12 years should be supervised in the use of this product. \*children under 2 years of age: consult a dentist or doctor.**

##### **Other information**

\*Keep tightly closed and in a dark place at room temperature.

\*Do not shake bottle. Hold away from face when opening.

**Inactive ingredient** purified water

**Antiseptic**

**\* For treatment of minor cuts and abrasions.**

**\* Oral debriding agent.**

**\* Use as a gargle or rinse.**

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN  
OR MISSING**

**DISTRIBUTED BY: EVAXO INTERNATIONAL INC.**

**Tampa, FI 33619**

**All rights reserved. Made in U.S.A**

**THE GREEN SOURCES**

**Exp. 5/2023**

**Packaging**

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**DISTRIBUTED BY: EVAXO INTERNATIONAL INC.**  
Tampa, FL 33619  
All rights reserved. Made in U.S.A



Batch no 564ev  
Exp. 5/2023

# EVAXO

## 3%

### Hydrogen Peroxide

USP

### Antiseptic

- \* For treatment of minor cuts and abrasions.
- \* Oral debriding agent.
- \* Use as a gargle or rinse.

32 FL OZ (946.6ml)

Drug Facts	
Active ingredient	Purpose
Hydrogen peroxide 3%.....	first aid antiseptic/oral debriding agent
<b>Uses</b> *First aid to help prevent the risk of infection in minor cuts, scrapes, and burns. *Aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth.	
<b>Warnings</b>	
For external use only.	
Do not use *in the eyes or apply over large areas of the body. *longer than one week.	
Ask a doctor before use if you have deep or puncture wounds, animal bites, or serious burns.	
Stop use and ask a doctor if *swelling, rash, or fever develops. *the condition persists or gets worse. *sore mouth symptoms do not improve in 7 days. *irritation, pain or redness persists or worsens.	
Keep out of reach of children. If swallowed, get medical help or contact a poison control center right away.	
<b>Directions</b> For use as a first aid antiseptic: *Clean affected area. *if bandaged, let dry first. *apply small amount of product on affected area 1-3 times a day. *may be covered with a sterile bandage. For use as an oral debriding agent (oral rinse): *adults and children 2 years of age and over: *mix with an equal amount of water. *swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. *use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor. *children under 12 years should be supervised in the use of this product. *children under 2 years of age: consult a dentist or doctor.	
<b>Other information</b>	
*Keep tightly closed and in a dark place at room temperature. *Do not shake bottle. Hold away from face when opening.	
Inactive ingredient purified water	

## EVAXO 3% HYDROGEN PEROXIDE

hydrogen peroxide liquid

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:78376-234
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE -	HYDROGEN	30 mg

UNII:BBX060AN9V)	PEROXIDE	in 1 mL
------------------	----------	---------

### Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:78376-234-32	946.6 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	05/01/2020	

**Labeler** - Evaxo International Inc (003464270)

### Establishment

Name	Address	ID/FEI	Business Operations
Evaxo International Inc		003464270	manufacture(78376-234)

Revised: 6/2020

Evaxo International Inc