

**TOPCARE CLEARLAX- polyethylene glycol 3350 powder, for solution**  
**Topco Associates LLC**

-----  
**Topco Associates LLC. ClearLax® Drug Facts**

**Active ingredient (in each dose)**

Polyethylene Glycol 3350, 17 g (cap filled to line)

**Purpose**

Osmotic Laxative

**Use**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

**Warnings**

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use**

if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug

**When using this product**

you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
  - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - use once a day
  - use no more than 7 days
- children 16 years of age or under: ask a doctor

## Other information

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

## Inactive ingredients

none

## Questions or comments?

**1-888-423-0139**

## Principal Display Panel

COMPARE TO MIRALAX<sup>®</sup> ACTIVE INGREDIENT

UNFLAVORED POWDER - GRIT FREE

ClearLax<sup>®</sup>

POLYETHYLENE GLYCOL 3350 POWDER FOR SOLUTION

OSMOTIC LAXATIVE

OUR PHARMACISTS RECOMMEND

Relieves Occasional Constipation (Irregularity)

Softens Stool

2 BOTTLES – NET WT 29.9 OZ (850g) Each

TOTAL NET WT 59.8 OZ (1700g)

100 ONCE-DAILY DOSES



COMPARE TO MIRALAX® ACTIVE INGREDIENT\*

UNFLAVORED POWDER • GRIT FREE

# ClearLax®

POLYETHYLENE GLYCOL 3350 POWDER FOR SOLUTION  
OSMOTIC LAXATIVE

- Relieves Occasional Constipation (Irregularity)
- Softens Stool



2 BOTTLES - NET WT 29.9 OZ (850g) Each  
TOTAL NET WT 59.8 OZ (1700g)

100 ONCE-DAILY DOSES



**Drug Facts**

**Active ingredient (in each dose) Purpose**  
Polyethylene Glycol 3350, 17 g  
(cap filled to line).....Osmotic Laxative

**Use**

- Relieves occasional constipation (irregularity)
- Generally produces a bowel movement in 1 to 3 days

**Warnings**

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. There may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

**Drug Facts (continued)**

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

**Directions**

- do not take more than the directed dose unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older
- fill to top of white section in cap which is marked to indicate the correct dose (17 g)
- fill and dissolve in an 8 to 16 ounce of beverage (cool, hot, or room temperature) then drink
- use once a day
- use no more than 7 days
- children 10 years of age or under: ask a doctor

**Other information**

- store at 20°-25°C (68°-77°F)
- temperature-dependent: do not use if printed foil seal under cap is missing, open or broken

**Inactive ingredients** none

**Questions or comments?** 1-888-423-0139

• Sugar Free  
Dissolves in Any Beverage

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.

DISTRIBUTED BY TOPCO ASSOCIATES LLC  
1615 GARDEN LANE, EL GORDON, VA 20120  
QUESTIONS? 1-888-423-0139  
www.topcare.com  
www.pca ndr and .com



This TopCare® product is laboratory tested to guarantee the highest quality. Your total satisfaction is guaranteed.



Scan here for more information.  
Call 1-888-423-0139 for more information.

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Miralax®.

10 88 1696V



polyethylene glycol 3350 powder, for solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-306
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-306-01	119 g in 1 BOTTLE; Type 0: Not a Combination Product	10/07/2009	
2	NDC:36800-306-02	238 g in 1 BOTTLE; Type 0: Not a Combination Product	10/07/2009	
3	NDC:36800-306-03	507 g in 1 BOTTLE; Type 0: Not a Combination Product	10/07/2009	
4	NDC:36800-306-04	850 g in 1 BOTTLE; Type 0: Not a Combination Product	05/13/2011	03/05/2015
5	NDC:36800-306-19	595 g in 1 BOTTLE; Type 0: Not a Combination Product	08/21/2013	
6	NDC:36800-306-07	3 in 1 PACKAGE	10/07/2009	
6		510 g in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:36800-306-08	2 in 1 PACKAGE	10/07/2009	10/07/2009
7		238 g in 1 BOTTLE; Type 0: Not a Combination Product		
8	NDC:36800-306-20	2 in 1 PACKAGE	03/18/2020	
8		850 g in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA090685	10/07/2009	

**Labeler** - Topco Associates LLC (006935977)

Revised: 3/2020

Topco Associates LLC