VALUMEDS 24-HOUR ALLERGY RELIEF- fexofenadine hydrochloride tablet Cabinet Health P.B.C.

ValuMeds 24-Hour Allergy Relief Fexofenadine

Drug Facts

Active ingredient (in each tablet)

Fexofenadine HCI USP, 180 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings

Do not use

if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if

an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

ask a health professional before use

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years take one 180 mg tablet with water once a day; do not take

of age and over more than 1 tablet in 24 hours

children under 6 years do not use

of age adults 65 years of age

and older ask a doctor

consumers with kidney ask a doctor

disease

Other information

• safety sealed: do not use if imprinted foil under bottle cap is opened or torn

store between 20° and 25°C (68° and 77°F)

protect from excessive moisture


Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, lactose monohydrate, macrogol, magnesium stearate, microcrystalline cellulose, pregelatinized starch, titanium dioxide

Questions?

call **1-908-242-6108**Mon-Fri 8:00 AM to 5:00 PM EST

Package Labelling:



Drug Facts (continued)

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.

When using this product

■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information

- safety sealed: do not use if imprinted foil under bottle cap is opened or torn store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients colloidal silicon dioxide,

araccormollaca andium hunramallaca iran avida black iran

croscarmenose socium, hypromenose, iron oxide diack, iron oxide red, iron oxide yellow, lactose monohydrate, macrogol, magnesium stearate, microcrystalline cellulose, pregelatinized starch, titanium dioxide

Questions or comments? contact 1-908-242-6108 Mon- Fri 8:00 AM EST to 5:00 PM PST

Bottle Pack

Drug Facts	Drug Facts (continued)	
Active ingredient Purpose (in each tablet) Fexofenadine HCI USP, 180 mgAntihistamine	Directions adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ itchy, watery eyes ■ itching of the nose or throat ■ itching of the nose or throat	children under 12 years of age adults 65 years of age and older consumers with kidney	do not use ask a doctor ask a doctor
Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.	safety sealed: do not use if imprinted foil under hottle can is opened or forp.	
When using this product ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.	Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, lactose monohydrate, macrogol, magnesium stearate, microcrystalline cellulose, pregelatinized starch, titanium dioxide	
If pregnant or breast-feeding, ask a health professional before use Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	Questions or comments? contact 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST	

Blister Pack

Active ingredient (in each to Fexofenadine HCI USP, 180 mg	ablet) Purpose Antihistamin	
Uses temporarily relieves these symptoms de ■ runny nose ■ itchy, watery eyes	ue to hay fever or other upper respiratory allergies: ■ sneezing ■ itching of the nose or throat	
	orgic reaction to this product or any of its ingredients	
When using this product ■ do not take more than directed ■ do not take more than directed ■ do not take with fruit juices (see Directions) Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If prequant or breast-feeding, ask a health professional before use		
Directions	of overdose, get medical help or contact a Poison Control Center right away	
adults and children 12 years of age and over children under 12 years of age adults 65 years of age and older consumers with kidney disease	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hour do not use ask a doctor ask a doctor	
Other information ■ safety sealed: do not use if carton is 20° and 25°C (68° and 77°F) ■ pr	opened or if individual blister units are torn or opened store between otect from excessive moisture	
Inactive ingredients colloidal silicon dioxide, croscarmellose lactose monohydrate, macrogol, magne	sodium, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, sium stearate, microcrystalline cellulose, pregelatinized starch, titanium	

VALUMEDS 24-HOUR ALLERGY RELIEF

fexofenadine hydrochloride tablet

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82725-1002	
Route of Administration	ORAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
FEXOFENADINE HYDROCHLORIDE (UNII: 2S068B75ZU) (FEXOFENADINE -	FEXOFENADINE HYDROCHLORIDE	180 mg		

Inactive Ingredients		
Ingredient Name	Strength	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		

CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics				
Color	pink (Peach Color)	Score	no score	
Shape	CAPSULE	Size	17mm	
Flavor		Imprint Code	G6	
Contains				

P	Packaging				
# Item Code Package Description		Marketing Start Date	Marketing End Date		
1	NDC:82725- 1002-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/01/2024		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211075	11/01/2024	

Labeler - Cabinet Health P.B.C. (117102391)

Revised: 10/2024 Cabinet Health P.B.C.