

LIQUID BANDAGE- benzethonium chloride plus dyclonine hydrochloride liquid
Meijer Distribution Inc

Meijer Liquid Bandage

Active ingredients

Benzethonium chloride 0.2%
Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic
Topical analgesic

Uses

- first aid to prevent bacterial contamination in minor cuts, scrapes and burns
- for the temporary relief of pain in minor cuts, scrapes, and burns

Warnings

For external use only.

Flammable:

Keep away from fire or flame.

Do not use

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- conditions persists or gets worse
- infection occurs
- symptoms persist for than 7 days or clear up and occur again within a few days

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- for adults and children 2 years of age and older
- clean affected area
- apply a small amount on the area 1 to 3 times daily and let dry
- may be covered with a sterile bandage
- if bandaged, let dry first
- children under 2 years of age; do not sure, consult a doctor

Other information

- cap bottle tightly
- wipe top of bottle clean after each use
- store between 20° to 25°C (68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off
- do not allow to come into contact with fabrics, floors, counterops or other surfaces -
- will stain.

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD slohol 40

Questions

Call 1-866-964-0939

Principal Display Panel

MEIJER

For small cuts & wounds


liquid bandage

BENZETHONIUM CHLORIDE 0.2%/FIRST AID ANTISEPTIC

DYCLONINE HYDROCHLORIDE 0.75%/TOPICAL ANALGESIC

KILLS 99.9% OF GERMS TO HELP PREVENT INFECTION

NET 0.3 FL OZ (9mL)



**FIRST AID ANTISEPTIC
DYCLONINE HYDROCHLORIDE 0.75%/
TOPICAL ANALGESIC**

**KILLS 99.9% OF GERMS TO HELP
PREVENT INFECTION***


Use Liquid Bandage for:

Blisters, chapped & cracked fingertips, hangnails, paper cuts, shaving nicks & help in preventing the formation of calluses.

Great for:

Athletes, active lifestyles, outdoor sportsmen & musicians.

53-033ME-02



Liquid Bandage

FIRST AID ANTISEPTIC/
TOPICAL ANALGESIC

0.3 FL. OZ. [9 mL]

Helps prevent
infection
Seals to protect
Water-resistant

*Kills 99.9% of 3 of the top germs within 60 minutes of contact.

*This product is not manufactured or distributed by Advantice Health, LLC, owner of the registered trademark New-Skin®.



DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com
MADE IN USA WITH U.S.
AND IMPORTED PARTS

PID 369389



DO NOT USE IF NECKBAND IS BROKEN

Drug Facts

Active ingredients
Benzethonium chloride 0.2%.....First aid antiseptic
Dyclonine hydrochloride 0.75%.....Topical analgesic

Uses
■ first aid to help prevent infection in minor cuts, scrapes, and burns
■ first aid for the temporary relief of pain in minor cuts, scrapes, and burns

Warnings
For external use only.
Flammable: Keep away from fire or flame.
Do not use ■ in the eyes ■ over large areas of the body
■ longer than 1 week unless directed by a doctor
Ask a doctor before use if you have ■ deep or puncture wounds ■ animal bites
■ serious burns
Stop use and ask a doctor if ■ condition persists or gets worse ■ infection occurs
■ symptoms persist for more than 7 days or clear up and occur again within a few days
Keep out of reach of children. If swallowed, get medical help or call a Poison
Control Center right away.
Directions ■ for adults and children 2 years of age and older ■ clean the affected
area ■ apply a small amount of this product on the area 1 to 3 times daily and let dry
■ may be covered with a sterile bandage ■ if bandaged, let dry first ■ children
under 2 years of age: do not use, consult a doctor
Other information ■ cap bottle tightly ■ wipe top of bottle clean after each
use ■ store between 20° to 25°C (68° to 77°F) ■ may temporarily sting upon
application ■ to remove, apply more Liquid Bandage and quickly wipe off ■ do
not allow to come into contact with fabrics, floors, countertops or other surfaces --
will stain.
Inactive ingredients acetone, amyli acetate, castor oil, ethyl acetate,
nitrocellulose, SD alcohol 40

Questions? Call 1-866-964-0939

LIQUID BANDAGE

benzethonium chloride plus dyclonine hydrochloride liquid

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:79481-0606

Route of Administration	TOPICAL
--------------------------------	---------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	2 mg in 1 mL
DYCLONINE HYDROCHLORIDE (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q30O)	DYCLONINE HYDROCHLORIDE	7.5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ACETONE (UNII: 1364PS73AF)	
AMYL ACETATE (UNII: 92Q24NH7AS)	
CASTOR OIL (UNII: D5340Y2I9G)	
ETHYL ACETATE (UNII: 76845O8NMZ)	
PYROXYLIN (UNII: KYR8BR2X6O)	
ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0606-1	1 in 1 CARTON	08/18/2023	
1		9 mL in 1 BOTTLE, WITH APPLICATOR; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	08/18/2023	

Labeler - Meijer Distribution Inc (006959555)

Revised: 10/2025

Meijer Distribution Inc