

**LIQUID BANDAGE- benzethonium chloride plus dyclonine hydrochloride liquid**  
**Meijer Distribution Inc**

-----  
**Meijer Liquid Bandage**

***Active ingredients***

Benzethonium chloride 0.2%  
Dyclonine hydrochloride 0.75%

***Purpose***

First aid antiseptic  
Topical analgesic

***Uses***

- first aid to prevent bacterial contamination in minor cuts, scrapes and burns
- for the temporary relief of pain in minor cuts, scrapes, and burns

**Warnings**

For external use only.

**Flammable:**

Keep away from fire or flame.

**Do not use**

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

**Stop use and ask a doctor if**

- conditions persists or gets worse
- infection occurs
- symptoms persist for than 7 days or clear up and occur again within a few days

**Keep out of reach of children.**

If swallowed, get medical help or call a Poison Control Center right away.

## **Directions**

- for adults and children 2 years of age and older
- clean affected area
- apply a small amount on the area 1 to 3 times daily and let dry
- may be covered with a sterile bandage
- if bandaged, let dry first
- children under 2 years of age; do not sure, consult a doctor

## **Other information**

- cap bottle tightly
- wipe top of bottle clean after each use
- store between 20° to 25°C (68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off
- do not allow to come into contact with fabrics, floors, counterops or other surfaces -  
- will stain.

## **Inactive ingredients**

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD slohol 40

## **Questions**

Call 1-866-964-0939

## **Principal Display Panel**

MEIJER

For small cuts & wounds


liquid bandage

BENZETHONIUM CHLORIDE 0.2%/FIRST AID ANTISEPTIC

DYCLONINE HYDROCHLORIDE 0.75%/TOPICAL ANALGESIC

KILLS 99.9% OF GERMS TO HELP PREVENT INFECTION

NET 0.3 FL OZ (9mL)



FIRST AID ANTISEPTIC  
DYCLONINE HYDROCHLORIDE 0.75%/  
TOPICAL ANALGESIC

KILLS 99.9% OF GERMS TO HELP  
PREVENT INFECTION\*

**Use Liquid Bandage for:**

Blisters, chapped & cracked fingertips,  
hangnails, paper cuts, shaving nicks &  
help in preventing the formation  
of calluses.

**Great for:**

Athletes, active lifestyles, outdoor  
sportsmen & musicians.

53-033ME-02



# Liquid Bandage

FIRST AID ANTISEPTIC/  
TOPICAL ANALGESIC

0.3 FL. OZ. [9 mL]

Helps prevent  
infection  
Seals to protect  
Water-resistant

\*Kills 99.9% of 3 of the top germs within 60 minutes of contact.

\*This product is not manufactured or  
distributed by Advantice Health, LLC,  
owner of the registered trademark New-Skin®.



DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com  
MADE IN USA WITH U.S.  
AND IMPORTED PARTS

PID 369389



**DO NOT USE IF NECKBAND IS BROKEN**

**Drug Facts**

**Active ingredients**  
Benzethonium chloride 0.2%.....First aid antiseptic  
Dyclonine hydrochloride 0.75%.....Topical analgesic

**Purpose**  
First aid to help prevent infection in minor cuts, scrapes, and burns

**Uses**  
■ first aid to help prevent infection in minor cuts, scrapes, and burns  
■ first aid for the temporary relief of pain in minor cuts, scrapes, and burns

**Warnings**  
For external use only.  
Flammable: Keep away from fire or flame.  
**Do not use** ■ in the eyes ■ over large areas of the body  
■ longer than 1 week unless directed by a doctor  
**Ask a doctor before use if you have** ■ deep or puncture wounds ■ animal bites  
■ serious burns  
**Stop use and ask a doctor if** ■ condition persists or gets worse ■ infection occurs  
■ symptoms persist for more than 7 days or clear up and occur again within a few days  
**Keep out of reach of children.** If swallowed, get medical help or call a Poison  
Control Center right away.  
**Directions** ■ for adults and children 2 years of age and older ■ clean the affected  
area ■ apply a small amount of this product on the area 1 to 3 times daily and let dry  
■ may be covered with a sterile bandage ■ if bandaged, let dry first ■ children  
under 2 years of age: do not use, consult a doctor  
**Other information** ■ cap bottle tightly ■ wipe top of bottle clean after each  
use ■ store between 20° to 25°C (68° to 77°F) ■ may temporarily sting upon  
application ■ to remove, apply more Liquid Bandage and quickly wipe off ■ do  
not allow to come into contact with fabrics, floors, countertops or other surfaces --  
will stain.  
**Inactive ingredients** acetone, amyli acetate, castor oil, ethyl acetate,  
nitrocellulose, SD alcohol 40

**Questions? Call 1-866-964-0939**

**DO NOT USE IF NECKBAND IS BROKEN**

**Active ingredients:** Benzethonium Chloride 0.2% and Dyclonine Hydrochloride 0.75% **Purpose:** First aid antiseptic and topical analgesic. **Uses:** First aid to help prevent infection in minor cuts, scrapes, and burns. First aid to help prevent infection in minor cuts, scrapes, and burns.

**Warnings:** For external use only. **Flammable:** Keep away from fire or flame. Do not use in the eyes, over large areas of the body, and longer than 1 week unless directed by a doctor.

**meijer**  
FOR SMALL CUTS & WOUNDS  
**Liquid Bandage**

**BENZETHONIUM CHLORIDE 0.2%/ FIRST AID ANTISEPTIC DYCLONINE HYDROCHLORIDE 0.75%/ TOPICAL ANALGESIC**

0.3 FL. OZ. (9 mL)

Ask a doctor before use if you have deep or puncture wounds, animal bites and serious burns.

When using this product avoid contact with eyes.

Stop use and ask a doctor if condition persists or gets worse, symptoms persist for more than 7 days or clear up and occur again within a few days, and infection occurs.

Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.

**Directions:** For adults and children 2 years of age and older. Clean the affected area. Apply a small amount of this product on the area 1 to 3 times daily and let dry. May be covered with a sterile bandage. If bandaged, let dry first. Children under 2 years of age do not use, consult a doctor. **Other information:** Cap bottle tightly.

Store between 20° to 25°C (68° to 77°F). May temporarily sting upon application. To remove, apply more Liquid Bandage and quickly wipe off. Do not allow to come into contact with fabrics, floors, countertops or other surfaces -- will stain. **Inactive ingredients:** acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

**Questions?** Call 1-866-964-0939

**MADE IN USA WITH U.S. AND IMPORTED PARTS**  
DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544

50-033ME-02

## LIQUID BANDAGE

benzethonium chloride plus dyclonine hydrochloride liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79481-0606
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZETHONIUM CHLORIDE</b> (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	2 mg in 1 mL
<b>DYCLONINE HYDROCHLORIDE</b> (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q300)	DYCLONINE HYDROCHLORIDE	7.5 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ACETONE</b> (UNII: 1364PS73AF)	
<b>AMYL ACETATE</b> (UNII: 92Q24NH7AS)	
<b>CASTOR OIL</b> (UNII: D5340Y2I9G)	
<b>ETHYL ACETATE</b> (UNII: 76845O8NMZ)	
<b>PYROXYLIN</b> (UNII: KYR8BR2X6O)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-	1 in 1 CARTON	08/18/2023	

0606-1	1 III I CARTON	08/18/2023	
1	9 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product		
<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	08/18/2023	

**Labeler -** Meijer Distribution Inc (006959555)

Revised: 2/2024

Meijer Distribution Inc