HEB EFFERVESCENT PAIN RELIEF- aspirin tablet, effervescent H. E. BUTT GROCERY COMPANY

HEB Effervescent Pain Relief

Active ingredient

(in each effervescent tablet)

Buffered aspirin 325 mg (NSAID)* *Nonsteroidal anti-inflammatory drug

Purpose

(in each effervescent tablet)

Buffered aspirin 325 mg **(NSAID)***.... Pain reliever/fever reducer *Nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of minor aches and pains due to:
- headache muscle pain backache
- toothache menstrual pain colds
- minor arthritis pain
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: \blacksquare hives \blacksquare facial swelling \blacksquare asthma (wheezing) \blacksquare shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use

Do not use \blacksquare if you are allergic to aspirin or any other pain reliever/fever reducer \blacksquare if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma ■ you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are taking a prescription drug for

🔳 diabetes 🔳 gout 🔳 arthritis

When using this product

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding

- feel faint vomit blood have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose

In case of overdose, get medical help or contact Poison Control Center right away.

Directions

■ do not exceed recommended dosage

■ to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking

adults and children 12 years and over: 2 tablets every 4 hours, or as directed by a doctor - do not exceed 8 tablets in 24 hours

60 years and over: 2 tablets every 4 hours, or as directed by a doctor - do not exceed 4 tablets in 24 hours

children under 12 years: consult a doctor

Take this product any time – morning, noon or night – when you need relief from headaches, backaches, body aches or muscle pain.

Place two (2) tablets in 4 oz. of water. Dissolve tablets completely.

You do not have to drink any of the residue that may be on the bottom of the glass. The medicines are in the water.

Other information

- each tablet contains: sodium 524 mg
- store at room temperature (59º-86ºF). Avoid excessive heat.

Inactive ingredients

anhydrous citric acid, sodium bicarbonate

Questions or comments

Questions or comments? 1-888-228-6937

Principal Display Panel



HEB EFFERVESCENT aspirin tablet, effervescent	PAIN RELIEF					
Product Information						
Product Type	HUMAN OTC DRUG	Item Code (Source)		NDC:37808-861		
Route of Administration	ORAL					
Active Ingredient/Active	Moiety					
Ingredient Name			Basis of Strength		Strength	
ASPIRIN (UNII: R16C05Y76E) (ASPI		ASPIRIN		325 mg		
Inactive Ingredients						
Ingredient Name					Strength	

SODIUM BICARBON	NATE (UNI	I: 8MDF5V3900)								
ANHYDROUS CITRI										
Product Characteristics										
Color		white	Score		no score					
Shape		ROUND	Size		25mm					
Flavor			Imprint Code		L421					
Contains										
Packaging										
# Item Code		Package Description		ſ	Marketing Start Date	Marketing End Date				
1 NDC:37808-861- 36	18 in 1 C	CARTON		07/	31/2024					
1	2 in 1 PO Product	in 1 POUCH; Type 0: Not a Combination roduct								
Marketing Information										
Marketing Category	Арр	Application Number or Monograph Citation		ı	Marketing Start Date	Marketing End Date				
OTC Monograph Dru	g M013	M013			07/31/2024					

Labeler - H. E. BUTT GROCERY COMPANY (007924756)

Revised: 12/2024

H. E. BUTT GROCERY COMPANY