

IVERMECTIN- ivermectin paste
Guangzhou Aimaixun Technology Co., Ltd.

Zimorven

Ivermectin Horse Paste 1.87

For Oral Use in Horses Only

IVERMECTIN

(ivermectin paste) 1.87%

Net Wt. 0.21 oz (6.08 g)

Contents treat up to 1,250 lb body weight

Approved by FDA under ANADA

Apple Flavored

INDICATIONS: Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. IVERMECTIN (ivermectin paste) provides effective treatment and control of the following parasites in horses:

Large Strongyles (adults): *Strongylus vulgaris* (also early forms in blood vessels), *S. edentatus* (also tissue stages), *S. equinus*, *Triodontophorus* spp. including *T. brevicauda* and *T. serratus*, and *Craterostomum acuticaudatum*.

Small Strongyles (adults, including those resistant to some benzimidazole class compounds): *Coronocyclus* spp. including *C. coronatus*, *C. labiatus* and *C. labratus*; *Cyathostomum* spp. including *C. catinatum* and *C. pateratum*; *Cylicocyclus* spp. including *C. insigne*, *C. leptostomum*, *C. nassatus* and *C. brevicapsulatus*; *Cylicodontophorus* spp.; *Cylicostephanus* spp. including *C. calicatus*, *C. goldi*, *C. longibursatus* and *C. minutus*; and *Petrovinema poculatum*.

Small Strongyles: Fourth-stage larvae.

Pinworms (adults and fourth-stage larvae): *Oxyuris equi*.

Ascarids (adults and third- and fourth-stage larvae): *Parascaris equorum*.

Hairworms (adults): *Trichostrongylus axei*.

Large-mouth Stomach Worms (adults): *Habronema muscae*.

Bots (oral and gastric stages): *Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*.

Lungworms (adults and fourth-stage larvae): *Dictyocaulus arnfieldi*.

Intestinal Threadworms (adults): *Strongyloides westeri*.

Summer Sores: Caused by *Habronema* and *Draschia* spp. cutaneous third-stage larvae.

Dermatitis: Caused by neck threadworm microfilariae, *Onchocerca* sp.

DOSAGE: This syringe contains sufficient paste to treat one 1250 lb horse at the recommended dose rate of 91 mcg ivermectin per lb (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 lb body weight. Do not underdose. Ensure each animal receives a complete dose based on a

current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

PARASITE CONTROL PROGRAM: All horses should be included in a regular parasite control program with particular attention being paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. IVERMECTIN (ivermectin paste) effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the chances of verminous arteritis caused by *Strongylus vulgaris*.

PRODUCT ADVANTAGES: Broad-spectrum Control - IVERMECTIN controls important internal parasites, including bots and the arterial stages of *S. vulgaris*, with a single dose. IVERMECTIN is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

ANIMAL SAFETY: IVERMECTIN (ivermectin paste) may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

STORAGE INFORMATION: Store at 68°F-77°F (20°C-25°C). Excursions between 59°F-86°F (15°C-30°C) are permitted.

ADMINISTRATION:

- (1) While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking, aligning the arrow on the ring to the line between the weight and lbs, as shown in the pictogram.
- (2) Lock the ring in place by making a 1/4 turn to the right. Ensure it is locked (it should no longer slide).
- (3) Make sure that the horse's mouth contains no feed.
- (4) Remove the cover from the tip of the syringe.
- (5) Insert the syringe tip into the horse's mouth at the space between the teeth.
- (6) Depress the plunger as far as it will go, depositing paste on the back of the tongue.
- (7) Immediately raise the horse's head for a few seconds after dosing.





WARNING: Do not use in horses intended for human consumption. **Not for use in humans. Keep this and all drugs out of the reach of children.** Refrain from smoking or eating when handling this product. Wash hands after use. Avoid contact with eyes.

PRECAUTIONS: Ivermectin paste has been formulated specifically for use in horses only. This product should not be used in other animal species, as severe adverse reactions, including fatalities in dogs, may result.

OTHER WARNINGS: Parasite resistance may develop to any dewormer. Resistance has

been reported for most classes of dewormers. Frequent treatment with a dewormer, used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated, may slow the development of parasite resistance. Clinical examinations, diagnostic tests, and parasite management history should be used to determine whether this product is appropriate for the herd prior to the use of any dewormer. Following the use of any dewormer, the effectiveness of treatment should be monitored (for example, through fecal egg count reduction tests or other appropriate methods). A decrease in a drug's effectiveness over time, as determined by fecal egg count reduction tests, may indicate the development of resistance to the dewormer administered. Parasite management programs should be adjusted accordingly based on regular monitoring.

ENVIRONMENTAL SAFETY: Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

INFORMATION FOR HORSE OWNERS: Swelling and itching reactions after treatment with ivermectin paste have occurred in horses carrying heavy infections of neck threadworm (*Onchocerca* spp.) microfilariae. These reactions are most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable. Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require additional appropriate therapy in conjunction with treatment with ivermectin paste. Reinfection and measures for its prevention should also be considered. Consult your veterinarian if the condition does not improve.

Distributed by:

GuangZhouShi AiMaiXun KeJi YouXian GongSi

Rm G381, Units 607-613, No. 36 Gaoke Road

Tianhe District, Guangzhou,

Guangdong, China

PRINCIPAL DISPLAY PANEL - 0.21 oz (6.08 g)

Keep out of the reach of children.

For Oral Use in Horses Only

NDC: 86249-001-01

IVERMECTIN


(ivermectin paste) 1.87%

Net Wt. 0.21 oz (6.08 g)

Contents treat up to 1,250 lb body weight

Approved by FDA under ANADA

Apple-Flavored



IVERMECTIN

ivermectin paste

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:86260-001
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IVERMECTIN (UNII: 8883YP2R6D) (IVERMECTIN - UNII:8883YP2R6D)	IVERMECTIN	18.7 mg in 1 g

Product Characteristics

Color		Score	
Shape		Size	
Flavor	APPLE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86260-001-01	6.08 g in 1 SYRINGE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200326	05/11/2026	

Labeler - Guangzhou Aimaixun Technology Co., Ltd. (516797312)

Registrant - Guangzhou Aimaixun Technology Co., Ltd. (516797312)

Establishment

Name	Address	ID/FEI	Business Operations
Oscar (Guangzhou) Pet Co., Ltd.		569802289	api manufacture

Revised: 5/2026

Guangzhou Aimaixun Technology Co., Ltd.