

**VICKS DAYQUIL NYQUIL HIGH BLOOD PRESSURE COLD AND FLU-  
acetaminophen, dextromethorphan hbr, doxylamine succinate  
The Procter & Gamble Manufacturing Company**

-----  
**VICKS® DayQuil™ NyQuil™ HIGH BLOOD PRESSURE COLD & FLU, LiquiCaps™  
Convenience Pack**

**VICKS® DayQuil™ HIGH BLOOD PRESSURE COLD & FLU, LiquiCaps™**

**Drug Facts**

**DayQuil™ Cold & Flu**

**Active Ingredient (in each LiquiCap)**

**Acetaminophen** 325 mg

Dextromethorphan HBr 10 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat • headache • minor aches & pains
- fever

**Warnings**

**Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 Liquicaps in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain or cough get worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed

- do not exceed 8 LiquiCaps per 24 hrs

---

adults & children 12 yrs & over	2 LiquiCaps with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

---

### **Other information**

store at no greater than 25°C (77°F)

### **Inactive ingredients**

FD&C Yellow No. 5, FD&C Yellow No. 6, gelatin, glycerin, lecithin, mica, polyethylene glycol, polyvinyl acetate phthalate, povidone, sorbitol sorbitan solution, titanium dioxide, water

### **Questions?**

**1-800-362-1683**

**TAMPER EVIDENT: DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED.**

**DIST. BY PROCTER & GAMBLE,**

**CINCINNATI OH 45202**

**VICKS® NyQuil™ HIGH BLOOD PRESSURE COLD & FLU, LiquiCaps™**

### **Drug Facts**

**NyQuil™ Cold & Flu**

### **Active Ingredient (in each LiquiCap)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

### **Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

### **Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing

## **Warnings**

### **Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 Liquicaps in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Allergy Alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

### **Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma,

chronic bronchitis, or emphysema

- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- do not exceed 8 LiquiCaps per 24 hrs

---

adults & children 12 yrs & over  
children 4 to under 12 yrs

2 LiquiCaps with water every 4 hrs  
ask a doctor

children under 4 yrs

do not use

---

## **Other information**

store at no greater than 25°C (77°F)

## **Inactive ingredients**

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, lecithin, mica, polyethylene glycol, polyvinyl acetate phthalate, povidone, sorbitol sorbitan solution, titanium dioxide, water

## **Questions?**

**1-800-362-1683**

**TAMPER EVIDENT: DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED.**

**DIST. BY PROCTER & GAMBLE,  
CINCINNATI OH 45202**

## **PRINCIPAL DISPLAY PANEL**

MAX STRENGTH

EASY TO OPEN BOTTLE†

VICKS®

DayQuil™

**Acetaminophen**, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Cough

DECONGESTANT FREE

NyQuil™ HIGH BLOOD PRESSURE COLD & FLU

**Acetaminophen**, Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Sneezing, Runny Nose

Cough

HBP

25% SMALLER\*

24 DAYQUIL LIQUICAPS™ 24 NYQUIL LIQUICAPS™ 48 TOTAL LIQUICAPS™



# VICKS DAYQUIL NYQUIL HIGH BLOOD PRESSURE COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate kit

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-805

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-805-48	1 in 1 CARTON; Type 0: Not a Combination Product	07/11/2023	

Quantity of Parts		
Part #	Package Quantity	Total Product Quantity
Part 1	48	
Part 2	48	

## Part 1 of 2

# VICKS DAYQUIL HIGH BLOOD PRESSURE COLD AND FLU

acetaminophen, dextromethorphan hbr capsule, liquid filled

## Product Information

**Item Code (Source)** NDC:69423-803

**Route of Administration** ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>POLYVINYL ACETATE PHTHALATE</b> (UNII: 58QVG85GW3)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MICA</b> (UNII: V8A1AW0880)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FD&amp;C YELLOW NO. 5</b> (UNII: I753WB2F1M)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	

## Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	15mm
<b>Flavor</b>		<b>Imprint Code</b>	HBP
<b>Contains</b>			

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/11/2023	

## Part 2 of 2

**VICKS NYQUIL HIGH BLOOD PRESSURE COLD AND FLU**



acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

### Product Information

**Item Code (Source)** NDC:69423-804

**Route of Administration** ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MICA</b> (UNII: V8A1AW0880)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>POLYVINYL ACETATE PHTHALATE</b> (UNII: 58QVG85GW3)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	15mm
<b>Flavor</b>		<b>Imprint Code</b>	NQ
<b>Contains</b>			

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/11/2023	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/11/2023	

**Labeler** - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2024

The Procter & Gamble Manufacturing Company