

**PAIN RELIEVER PM EXTRA STRENGTH- acetaminophen, diphenhydramine  
hcl tablet, film coated  
L.N.K. International, Inc.**

-----  
**Quality Plus 44-235**

***Active ingredients (in each caplet)***

Acetaminophen 500 mg  
Diphenhydramine HCl 25 mg

***Purpose***

Pain reliever  
Nighttime sleep-aid

***Uses***

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis

- liver disease
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- avoid alcoholic beverages
- drowsiness will occur
- do not drive a motor vehicle or operate machinery

**Stop use and ask a doctor if**

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- **do not take more than directed**
- adults and children 12 years and over: take 2 caplets at bedtime. Do not take more than 2 caplets of this product in 24 hours.
- children under 12 years: do not use

***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

***Inactive ingredients***

colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C blue #1 aluminum lake, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, stearic acid, talc, titanium dioxide

**Questions or comments?**

**1-800-426-9391**

***Principal Display Panel***

**QUALITY**

**+PLUS**

NDC 50844-235-15

\*Compare to active ingredients in  
Extra Strength Tylenol® PM

**EXTRA STRENGTH**

**Pain Reliever PM**

**Acetaminophen**, Diphenhydramine HCl

PAIN RELIEVER/NIGHTTIME SLEEP-AID

**50 Caplets**

Non-habit Forming

ACTUAL

SIZE

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*This product is not manufactured or distributed  
by Johnson & Johnson Corporation, owner of the  
registered trademark Extra Strength Tylenol®  
PM.

50844 REV0521K23515

Distributed by

**LNK International, Inc.**

60 Arkay Drive

Hauppauge, NY 11788

USA

No Print  
Glue Area

**Drug Facts (continued)**

Ask a doctor before use if you have  
 ■ breathing problems such as emphysema or chronic bronchitis ■ liver disease ■ glaucoma  
 ■ difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are  
 ■ taking the blood thinning drug warfarin  
 ■ taking sedatives or tranquilizers

**When using this product**

■ avoid alcoholic beverages ■ drowsiness will occur  
 ■ do not drive a motor vehicle or operate machinery

**Stop use and ask a doctor if**

■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. ■ new symptoms occur  
 ■ redness or swelling is present  
 ■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days  
 These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

■ do not take more than directed  
 ■ adults and children 12 years and over: take 2 caplets at bedtime. Do not take more than 2 caplets of this product in 24 hours.  
 ■ children under 12 years: do not use

No Print  
Glue Area

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

**Drug Facts**

**Active ingredients (in each caplet)**  
 Acetaminophen 500 mg ..... Pain reliever  
 Diphenhydramine HCl 25 mg ..... Nighttime sleep-aid

**Uses** temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
 ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin redness ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

■ with any other product containing diphenhydramine, even one used on skin  
 ■ in children under 12 years of age  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

B-1603-235-15-R  
 REV0521K23515

No print/No varnish  
 Lot & Exp date



TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

QUALITY PLUS

NDC 50844-235-15

\*Compare to active ingredients in Extra Strength Tylenol® PM

EXTRA STRENGTH

PAIN RELIEVER PM

Acetaminophen, Diphenhydramine HCl

PAIN RELIEVER/NIGHTTIME SLEEP-AID

50 Caplets

Non-habit Forming



ACTUAL SIZE

QUALITY PLUS

EXTRA STRENGTH

PAIN RELIEVER PM

Acetaminophen, Diphenhydramine HCl

50 Caplets

**Drug Facts (continued)**

**Other information**

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
 ■ see end flap for expiration date and lot number

**Inactive ingredients** colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C blue #1, aluminum lake, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, stearic acid, talc, titanium dioxide

**Questions or comments? 1-800-426-9391**

\*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Extra Strength Tylenol® PM.

50844 REV0521K23515  
 Distributed by  
 LNK International, Inc.  
 60 Arkey Drive  
 Hauppauge, NY 11788  
 USA

Quality Plus 44-235

**PAIN RELIEVER PM EXTRA STRENGTH**

acetaminophen, diphenhydramine hcl tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50844-235
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	44;235
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-235-15	1 in 1 CARTON	05/15/1994	
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/15/1994	

**Labeler** - L.N.K. International, Inc. (038154464)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-235)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-235) , pack(50844-235)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-235)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(50844-235)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-235)

Revised: 7/2025

L.N.K. International, Inc.