

CAREONE IBUPROFEN- ibuprofen tablet, film coated
American Sales Company

American Sales Company Ibuprofen Drug Facts

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

Adults and children 12 years and older:

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

Children under 12 years: ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- see end panel for lot number and expiration date

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, iron oxides, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

CAREone®

Compare to the active ingredient in Motrin[®] IB

IBUPROFEN

Tablets, 200mg

Actual Size

Pain Reliever/Fever Reducer (NSAID)

OUR PHARMACISTS RECOMMEND

Gluten Free

See Revised Warning

50 CAPLETS**

**Capsule-Shaped Tablets

51771 OF C4



Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active Ingredient (in each caplet)
Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer
Nonsteroidal anti-inflammatory drug

Purposes
Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer

Warnings

Always alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or a longer time than directed. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer; if you have or after heart surgery.
Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers; the stomach bleeding warning applies to you; you have a history of stomach problems, such as:

Drug Facts (continued)

asthma; heartburn; you have high blood pressure, heart disease, liver, stomach, kidney disease, asthma, or had a stroke; you are taking a diuretic. Ask a doctor or pharmacist before use if you are: taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin; under a doctor's care for any serious condition; taking any other drug.
When using this product: take with food or milk if stomach upset occurs.
Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint; vomit blood; have bloody or black stools; have stomach pain that does not get better; you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, pain, gets worse or lasts more than 10 days, faint or gets worse or lasts more than 3 days, redness or swelling, or present in the pain/relief area; any new symptoms appear.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by your doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions Do not take more than directed. The smallest effective dose should be used. Adults and children 12 years and older: take 1 caplet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 caplet, 2 caplets may be used; do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ask a doctor.

Drug Facts (continued)

Other information: read all warnings and directions before use; store below 25°C (77°F); avoid high humidity; and excessive heat above 40°C (104°F); see expiration date and expiration date.
Keep five ingredients outside silicon dioxide, croscarmellose sodium, FD &C red #40, aluminum lake, FD &C yellow #5, aluminum lake, iron oxides, m-cresol, titanium dioxide, polyethylene glycol, polyvinyl alcohol, stearic acid, talc, titanium dioxide.
Questions or comments? 1-800-719-9260
DO NOT USE IF PRINTED SERIAL NUMBER IS BROKEN OR MISSING
Johnson Consumer Inc., distributor of Motrin® IB, Pain Reliever/NSAID, is not manufactured or distributed by Johnson & Johnson Consumer Inc. ADUSADISTRIBUTORS LLC
SALES BUREAU, INC. 28147
1-833-992-3872
Quality guaranteed or your money back.

Actual Size



CAREone® NDC 41520-495-71
IBUPROFEN
Tablets, 200mg
Pain Reliever/Fever Reducer
(NSAID)
Gluten Free
"Capsule-Shaped" Tablets

CAREone® NDC 41520-495-71

Compare to the active ingredient in Motrin® IB***

IBUPROFEN

Tablets, 200mg
Pain Reliever/
Fever Reducer (NSAID)



Gluten Free

See Revised Warning

50 CAPLETS**
**Capsule-Shaped Tablets

CAREONE IBUPROFEN
ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-495
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	I2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-495-71	1 in 1 CARTON	07/20/2016	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41520-495-78	1 in 1 CARTON	07/20/2016	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077349	07/20/2016	

Labeler - American Sales Company (809183973)

Revised: 11/2023

American Sales Company