

**FLUZONE HIGH-DOSE QUADRIVALENT NORTHERN HEMISPHERE- influenza a virus a/victoria/4897/2022 ivr-238 (h1n1) antigen (formaldehyde inactivated), influenza a virus a/croatia/10136rv/2023 x-425a (h3n2) antigen (formaldehyde inactivated), influenza b virus b/phuket/3073/2013 antigen (formaldehyde inactivated), and influenza b virus b/michigan/01/2021 antigen (formaldehyde inactivated) injection, suspension
Sanofi Pasteur Inc.**

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Fluzone[®] High-Dose Quadrivalent safely and effectively. See full prescribing information for Fluzone High-Dose Quadrivalent.

**Fluzone High-Dose Quadrivalent (Influenza Vaccine)
injectable suspension, for intramuscular use
2025-2026 Formula
Initial U.S. Approval: 2019**

----- **INDICATIONS AND USAGE** -----

Fluzone High-Dose Quadrivalent is a vaccine indicated for active immunization for the prevention of disease caused by influenza A subtype viruses and type B viruses contained in the vaccine. (1)

Fluzone High-Dose Quadrivalent is approved for use in persons 65 years of age and older. (1)

----- **DOSAGE AND ADMINISTRATION** -----

For intramuscular use

Administer Fluzone High-Dose Quadrivalent as a single 0.7 mL dose. (2)

----- **DOSAGE FORMS AND STRENGTHS** -----

Fluzone High-Dose Quadrivalent is an injectable suspension. A single dose is 0.7 mL (3)

----- **CONTRAINDICATIONS** -----

Do not administer Fluzone High-Dose Quadrivalent to anyone with a history of severe allergic reaction (e.g., anaphylaxis) to any component of the vaccine, including egg protein, or to a previous dose of any influenza vaccine (4, 11)

----- **WARNINGS AND PRECAUTIONS** -----

If Guillain-Barré Syndrome (GBS) has occurred within 6 weeks following previous influenza vaccination, the decision to give Fluzone High-Dose Quadrivalent should be based on careful consideration of the potential benefits and risks. (5.1)

----- **ADVERSE REACTIONS** -----

In adults \geq 65 years of age, the most common ($>10\%$) injection-site adverse reaction was pain (41.3%); the most common solicited systemic adverse reactions were myalgia (22.7%), headache (14.4%) and malaise (13.2%). (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Sanofi Pasteur Inc., Discovery Drive, Swiftwater, PA 18370 at 1-800-822-2463 (1-800-VACCINE) or VAERS at 1-800-822-7967 or <https://vaers.hhs.gov>.

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 7/2025

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Dose and Schedule

2.2 Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Guillain-Barré Syndrome
- 5.2 Preventing and Managing Allergic Reactions
- 5.3 Altered Immunocompetence
- 5.4 Limitations of Vaccine Effectiveness
- 5.5 Syncope

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action

13 NONCLINICAL TOXICOLOGY

14 CLINICAL STUDIES

- 14.1 Immunogenicity of Fluzone High-Dose Quadrivalent in Adults 65 Years of Age and Older
- 14.2 Efficacy of Fluzone High-Dose in Adults 65 Years of Age and Older

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Fluzone[®] High-Dose Quadrivalent is a vaccine indicated for active immunization for the prevention of disease caused by influenza A subtype viruses and type B viruses contained in the vaccine.

Fluzone High-Dose Quadrivalent is indicated for use in persons 65 years of age and older.

2 DOSAGE AND ADMINISTRATION

For intramuscular use

2.1 Dose and Schedule

Administer Fluzone High-Dose Quadrivalent as a single 0.7 mL dose.

2.2 Administration

Fluzone High-Dose Quadrivalent is a colorless opalescent liquid. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. If either of these conditions exist, the vaccine should not be administered.

Before administering a dose of vaccine, shake the prefilled syringe.

Administer Fluzone High-Dose Quadrivalent intramuscularly.

3 DOSAGE FORMS AND STRENGTHS

Fluzone High-Dose Quadrivalent is an injectable suspension.

A single dose is 0.7 mL.

4 CONTRAINDICATIONS

Do not administer Fluzone High-Dose Quadrivalent to anyone with a history of severe allergic reaction (e.g., anaphylaxis) to any component of the vaccine [see *Description (11)*], including egg protein, or to a previous dose of any influenza vaccine.

5 WARNINGS AND PRECAUTIONS

5.1 Guillain-Barré Syndrome

If Guillain-Barré Syndrome (GBS) has occurred within 6 weeks following any previous influenza vaccination, the decision to give Fluzone High-Dose Quadrivalent should be based on careful consideration of the potential benefits and risks.

The 1976 swine influenza vaccine was associated with an elevated risk of GBS. Evidence for a causal relation of GBS with other influenza vaccines is inconclusive; if an excess risk exists, it is probably slightly more than 1 additional case per 1 million persons vaccinated. (See reference 1.)

5.2 Preventing and Managing Allergic Reactions

Appropriate medical treatment must be immediately available to manage potential anaphylactic reactions following administration of the vaccine.

5.3 Altered Immunocompetence

If Fluzone High-Dose Quadrivalent is administered to immunocompromised persons, including those receiving immunosuppressive therapy, the immune response may be lower than expected.

5.4 Limitations of Vaccine Effectiveness

Vaccination with Fluzone High-Dose Quadrivalent may not protect all recipients.

5.5 Syncope

Syncope (fainting) has been reported following vaccination with Fluzone High-Dose Quadrivalent. Procedures should be in place to avoid injury from fainting.

6 ADVERSE REACTIONS

In adults ≥ 65 years of age, the most common ($>10\%$) injection-site adverse reaction was pain (41.3%); the most common solicited systemic adverse reactions were myalgia (22.7%), headache (14.4%) and malaise (13.2%).

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse event rates observed in the clinical trial(s) of a vaccine cannot be directly compared to rates in the clinical trial(s) of another vaccine and may not reflect the rates observed in practice. One clinical study has evaluated the safety of Fluzone High-Dose Quadrivalent.

Study 1 (NCT03282240) was a randomized, active-controlled, modified double-blind pre-licensure trial conducted in the U.S. The study compared the safety and immunogenicity of Fluzone High-Dose Quadrivalent to those of Fluzone High-Dose (trivalent formulation). The safety analysis set included 1777 Fluzone High-Dose Quadrivalent recipients, 443 Fluzone High-Dose recipients, and 450 investigational Fluzone High-Dose containing the alternate B influenza strain recipients.

The most common reactions occurring after Fluzone High-Dose Quadrivalent administration were injection-site pain (41.3%), myalgia (22.7%), headache (14.4%), and malaise (13.2%). Onset usually occurred within the first 3 days after vaccination. The majority of solicited reactions resolved within three days of vaccination.

Table 1 displays solicited adverse reactions for Fluzone High-Dose Quadrivalent compared to Fluzone High-Dose reported within 7 days after vaccination and collected using standardized diary cards.

Table 1: Frequency of Solicited Injection-Site Systemic Adverse Reactions within 7 Days after Vaccination with Fluzone High-Dose Quadrivalent or Fluzone High-Dose, Adults 65 Years of Age and Older (Study 1*)

| | Fluzone High-Dose Quadrivalent (N [†] =1761-1768) Percentage | | Fluzone High-Dose [‡] (N [†] =885-889) Percentage | |
|--------------------------------------|---|---------|---|---------|
| | Any | Grade 3 | Any | Grade 3 |
| Local Reactions | | | | |
| Injection Site Pain [§] | 41.3 | 0.7 | 36.4 | 0.2 |
| Injection Site Erythema [¶] | 6.2 | 0.6 | 5.7 | 0.2 |
| Injection Site Swelling [¶] | 4.9 | 0.3 | 4.7 | 0.1 |

| | | | | |
|--|------|-----|------|-----|
| Injection Site Induration [¶] | 3.7 | 0.2 | 3.5 | 0.1 |
| Injection Site Bruising [¶] | 1.3 | 0.0 | 1.1 | 0.0 |
| Systemic Reactions | | | | |
| Myalgia [§] | 22.7 | 0.9 | 18.9 | 0.7 |
| Headache [§] | 14.4 | 0.6 | 13.6 | 0.4 |
| Malaise [§] | 13.2 | 0.7 | 13.4 | 0.4 |
| Shivering [§] | 5.4 | 0.3 | 4.7 | 0.3 |
| Fever [#] | 0.4 | 0.2 | 0.9 | 0.2 |

* NCT03282240

† N is the number of vaccinated participants with available data for the events listed

‡ Safety results for the Fluzone High-Dose and investigational Fluzone High-Dose containing the alternate B influenza strain recipients were pooled for the analysis.

§ Grade 3: A type of AE that interrupts usual activities of daily living, or significantly affects clinical status, or may require intensive therapeutic intervention.

¶ Grade 3: > 100 mm

Grade 3: ≥ 102.1°F (39.0°C)

Based on data from Fluzone High-Dose, solicited injection site reactions and systemic adverse reactions were slightly more frequent after vaccination with Fluzone High-Dose compared to a standard-dose vaccine.

Unsolicited non-serious adverse events were reported in 279 (15.7%) recipients in the Fluzone High-Dose Quadrivalent group and 140 (15.7%) recipients in the Fluzone High-Dose group. The most commonly reported unsolicited adverse event was cough.

Within 180 days post-vaccination, 80 (4.5%) Fluzone High-Dose Quadrivalent recipients and 48 (5.4%) Fluzone High-Dose recipients experienced a serious adverse event (SAE). None of the SAEs were assessed as related to the study vaccines.

6.2 Postmarketing Experience

The following additional adverse events have been spontaneously reported during the postmarketing use of Fluzone High-Dose, Fluzone, Fluzone Quadrivalent, or Fluzone High-Dose Quadrivalent. Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to vaccine exposure. Adverse events were included based on one or more of the following factors: severity, frequency of reporting, or strength of evidence for a causal relationship to Fluzone High-Dose, Fluzone Fluzone Quadrivalent or Fluzone High-Dose Quadrivalent.

- *Blood and Lymphatic System Disorders:* Thrombocytopenia, lymphadenopathy
- *Immune System Disorders:* Anaphylaxis, other allergic/hypersensitivity reactions (including urticaria, angioedema)
- *Eye Disorders:* Ocular hyperemia
- *Nervous System Disorders:* Guillain-Barré syndrome (GBS), convulsions, febrile convulsions, myelitis (including encephalomyelitis and transverse myelitis), facial palsy (Bell's palsy), optic neuritis/neuropathy, brachial neuritis, syncope (shortly after vaccination), dizziness, paresthesia
- *Vascular Disorders:* Vasculitis, vasodilatation
- *Respiratory, Thoracic and Mediastinal Disorders:* Dyspnea, cough, wheezing, throat

tightness, oropharyngeal pain, and rhinorrhea

- *Gastrointestinal Disorders*: Vomiting
- *Skin and Subcutaneous Tissue Disorders*: Stevens-Johnson syndrome
- *General Disorders and Administration Site Conditions*: pruritus, asthenia/fatigue, chest pain, chills

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Fluzone High-Dose Quadrivalent is not approved for use in persons <65 years of age. There are no human or animal data available on Fluzone High-Dose Quadrivalent to establish whether there is a vaccine-associated risk with use of Fluzone High-Dose Quadrivalent in pregnancy.

8.2 Lactation

Fluzone High-Dose Quadrivalent is not approved for use in persons <65 years of age. No human or animal data are available to assess the effects of Fluzone High-Dose Quadrivalent on the breastfed infant or on milk production/excretion.

8.4 Pediatric Use

Safety and effectiveness of Fluzone High-Dose Quadrivalent in children younger than 18 years of age have not been established.

8.5 Geriatric Use

Safety, immunogenicity, and efficacy of Fluzone High-Dose Quadrivalent have been evaluated in adults 65 years of age and older [see *Adverse Reactions (6.1) and Clinical Studies (14)*].

11 DESCRIPTION

Fluzone High-Dose Quadrivalent for intramuscular use is an inactivated influenza vaccine, prepared from influenza viruses propagated in embryonated chicken eggs. The virus-containing allantoic fluid is harvested and inactivated with formaldehyde. Influenza virus is concentrated and purified in a linear sucrose density gradient solution using a continuous flow centrifuge. The virus is then chemically disrupted using a non-ionic surfactant, octylphenol ethoxylate (Triton[®] X-100), producing a "split virus." The split virus containing hemagglutinin (HA) antigen is further purified and then suspended in sodium phosphate-buffered isotonic sodium chloride solution. The Fluzone High-Dose Quadrivalent process uses an additional concentration factor after the ultrafiltration step in order to obtain a higher HA antigen concentration. The purified split virus from the four strains included in the vaccine are produced separately and then combined to make the quadrivalent formulation.

Fluzone High-Dose Quadrivalent is an injectable suspension and is a colorless opalescent liquid.

Neither antibiotics nor preservative are used in the manufacture of Fluzone High-Dose

Quadrivalent.

The Fluzone High-Dose Quadrivalent prefilled syringe presentation is not made with natural rubber latex.

Fluzone High-Dose Quadrivalent is standardized according to United States Public Health Service requirements and is formulated to contain HA of each of the following four influenza strains recommended for the 2025-2026 influenza season:

A/Victoria/4897/2022 IVR-238 (H1N1), A/Croatia/10136RV/2023 X-425A (H3N2), B/Phuket/3073/2013 (B Yamagata lineage), and B/Michigan/01/2021 (a B/Austria/1359417/2021-like virus, B Victoria lineage).

The amounts of HA and other ingredients per dose of vaccine are listed in Table 2.

Table 2: Fluzone High-Dose Quadrivalent Ingredients

| Ingredient | Quantity (per dose) |
|--|--|
| | Fluzone High-Dose Quadrivalent 0.7 mL Dose |
| Active Substance: Split influenza virus, inactivated strains* | 240 mcg HA total |
| A (H1N1) | 60 mcg HA |
| A (H3N2) | 60 mcg HA |
| B (Victoria Lineage) | 60 mcg HA |
| B (Yamagata Lineage) | 60 mcg HA |
| Other: | |
| Sodium phosphate-buffered isotonic sodium chloride solution | QS [†] to appropriate volume |
| Formaldehyde | ≤140 mcg |
| Octylphenol ethoxylate | ≤350 mcg |
| Gelatin | None |
| Preservative | None |

* per United States Public Health Service (USPHS) requirement

† Quantity sufficient

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Specific levels of hemagglutination inhibition (HI) antibody titer post-vaccination with inactivated influenza virus vaccines have not been correlated with protection from influenza virus infection. In some human studies, antibody titers $\geq 1:40$ have been associated with protection from influenza illness in up to 50% of participants. (See references 2 and 3.)

Antibodies against one influenza virus type or subtype confer limited or no protection against another. Furthermore, antibodies to one antigenic variant of influenza virus might not protect against a new antigenic variant of the same type or subtype. Frequent development of antigenic variants through antigenic drift is the virologic basis for

seasonal epidemics and the reason for the usual change of one or more new strains in each year's influenza vaccine.

13 NONCLINICAL TOXICOLOGY

Carcinogenesis, Mutagenesis, Impairment of Fertility

Fluzone High-Dose Quadrivalent has not been evaluated for carcinogenic or mutagenic potential or for impairment of fertility.

14 CLINICAL STUDIES

14.1 Immunogenicity of Fluzone High-Dose Quadrivalent in Adults 65 Years of Age and Older

Study 1 (NCT03282240, see <http://clinicaltrials.gov>) was a randomized, active-controlled, modified double-blind trial in adults 65 years of age and older conducted in the US. The study compared the safety and immunogenicity of Fluzone High-Dose Quadrivalent to those of Fluzone High-Dose. The objective was to demonstrate immunologic non-inferiority of Fluzone High-Dose Quadrivalent to Fluzone High-Dose, as assessed by HAI geometric mean antibody titers (GMTs) at Day 28 and seroconversion rates, to strains common to formulations of both vaccines, based on pre-specified criteria.

A total of 2670 adults from 65 years of age were randomized (4:1:1) to receive one dose of either Fluzone High-Dose Quadrivalent or one of two formulations of Fluzone High-Dose (one formulation contained a B strain of the Victoria lineage [TIV-HD1] while the other contained a B strain of the Yamagata lineage [TIV-HD2]).

Females accounted for 58.2% of participants in the Fluzone High-Dose Quadrivalent group and 57.4% of participants in the Fluzone High-Dose group (TIV-HD1 and TIV-HD2, pooled). The mean age was 72.9 years (range: 65 through 100 years) in the Fluzone High-Dose Quadrivalent group and the mean age was 73.0 (range: 65 through 95 years) in the Fluzone High-Dose group. The percentage of subjects 75 years of age or older was 35.4% in the Fluzone High-Dose Quadrivalent group and 35.8% in the Fluzone High-Dose group. Most participants were White (91.2% and 89.7%), followed by Black (6.8% and 8.0%), and Hispanic (2.8% and 2.6%) in the Fluzone High-Dose Quadrivalent and Fluzone High-Dose groups, respectively.

The immunogenicity results of Study 1 are summarized in Table 3 and Table 4 below.

Table 3: Post-vaccination HAI Antibody GMTs and Analyses of Non-inferiority of Fluzone High-Dose Quadrivalent Relative to Fluzone High-Dose, Adults 65 Years of Age and Older, Per-Protocol Analysis Set (Study 1*)

| Influenza Strain | GMT | | | GMT Ratio | Met Predefined Non-inferiority Criteria [†] |
|------------------|---------------------------------|--|--|--------------------------------|--|
| | QIV-HD N [¶] =1679- | TIV-HD1 [‡] (B1 Victoria) N [¶] =423 | TIV-HD2 [§] (B2 Yamagata) N [¶] =430 | QIV-HD over TIV-HD (95% CI) | |
| | | | | | |

| | | | | | |
|-----------------------------|-------------|-----|-----|---------------------------|-----|
| | 1680 | | | | |
| A (H1N1)[#] | 312 | 374 | | 0.83 (0.744; 0.932) | Yes |
| A (H3N2)[#] | 563 | 594 | | 0.95 (0.842; 1.066) | Yes |
| B1 (Victoria) | 516 | 476 | -- | 1.08 (0.958; 1.224) | Yes |
| B2 (Yamagata) | 578 | -- | 580 | 1.00 (0.881; 1.129) | Yes |

* NCT03282240

† Predefined noninferiority criterion for the GMT ratio: the lower limit of the 95% CI of the GMT ratio (QIV-HD divided by TIV-HD) is >0.667

‡ TIV-HD1 contained A/Michigan/45/2015 (H1N1), A/Hong Kong/4801/2014 (H3N2), and B/Brisbane/60/2008 (B1, Victoria lineage)

§ TIV-HD2 contained A/Michigan/45/2015 (H1N1), A/Hong Kong/4801/2014 (H3N2), and B/Phuket/3073/2013 (B2, Yamagata lineage)

¶ N is the number of vaccinated participants with available data for the immunologic endpoint listed

Pooled TIV-HD group includes subjects vaccinated with either TIV-HD1 or TIV-HD2 for the A strain comparison

Table 4: Seroconversion Rates and Analyses of Non-inferiority of Fluzone High-Dose Quadrivalent Relative to Fluzone High-Dose, Adults 65 Years of Age and Older, Per-Protocol Analysis Set (Study 1*)

| Influenza Strain | Seroconversion Rates (Percentage) [†] | | | Difference of Seroconversion Rates | Met Predefined Non-inferiority Criteria [‡] |
|-----------------------------|--|---|---|------------------------------------|--|
| | QIV-HD N [#] =1668- 1669 | TIV-HD1 [§] (B1 Victoria) N [#] =420- 421 | TIV-HD2 [¶] (B2 Yamagata) N [#] =428 | QIV-HD minus TIV-HD (95% CI) | |
| A (H1N1)[¶] | 50.4 | 53.7 | | -3.27 (-7.37; 0.86) | Yes |
| A (H3N2)[¶] | 49.8 | 50.5 | | -0.71 (-4.83; 3.42) | Yes |
| B1 (Victoria) | 36.5 | 39.0 | -- | -2.41 (-7.66; 2.70) | Yes |
| B2 (Yamagata) | 46.6 | -- | 48.4 | -1.75 (-7.04; 3.53) | Yes |

* NCT03282240

† Seroconversion Rates: For subjects with a pre-vaccination titer <10 (1/dil), proportion of subjects with a post-vaccination titer ≥40 (1/dil) and for subjects with a pre-vaccination titer ≥10 (1/dil), proportion of subjects with a ≥four-fold increase from pre-vaccination to post-vaccination titer

‡ Predefined noninferiority criterion for seroconversion: the lower limit of the two-sided 95% CI of the difference of the seroconversion rates (QIV-HD minus TIV-HD) is >-10%

- § TIV-HD1 contained A/Michigan/45/2015 (H1N1), A/Hong Kong/4801/2014 (H3N2), and B/Brisbane/60/2008 (B1, Victoria lineage)
- ¶ TIV-HD2 contained A/Michigan/45/2015 (H1N1), A/Hong Kong/4801/2014 (H3N2), and B/Phuket/3073/2013 (B2, Yamagata lineage)
- # N is the number of vaccinated participants with available data for the immunologic endpoint listed
- p Pooled TIV-HD group includes subjects vaccinated with either TIV-HD1 or TIV-HD2 for the A strain comparison

Fluzone High-Dose Quadrivalent was as immunogenic as Fluzone High-Dose for GMTs and seroconversion rates for the common influenza strains. Fluzone High-Dose Quadrivalent induced a superior immune response, based on a pre-specified superiority criterion, with respect to the additional B strain than the immune response induced by Fluzone High-Dose formulation that did not contain the additional B strain.

14.2 Efficacy of Fluzone High-Dose in Adults 65 Years of Age and Older

The efficacy of Fluzone High-Dose (trivalent formulation) is relevant to Fluzone High-Dose Quadrivalent since both vaccines are manufactured according to the same process and have overlapping compositions.

Study 2 (NCT01427309) was a multi-center, double-blind, post-licensure efficacy trial conducted in the U.S. and Canada in which adults 65 years of age and older were randomized (1:1) to receive either Fluzone High-Dose or Fluzone. The study was conducted over two influenza seasons (2011-2012 and 2012-2013); 53% of participants enrolled in the first year of the study were re-enrolled and re-randomized in the second year. The per-protocol analysis set for efficacy assessments included 15,892 Fluzone High-Dose recipients and 15,911 Fluzone recipients. The majority (67%) of participants in the per-protocol analysis set for efficacy had one or more high-risk chronic comorbid conditions.

In the per-protocol analysis set, females accounted for 57.2% of participants in the Fluzone High-Dose group and 56.1% of participants in the Fluzone group. In both groups, the median age was 72.2 years (range 65 through 100 years). Overall, most participants in the study were White (95%); approximately 4% of study participants were Black, and approximately 6% reported Hispanic ethnicity.

The primary endpoint of the study was the occurrence of laboratory-confirmed influenza (as determined by culture or polymerase chain reaction) caused by any influenza viral type/subtype in association with influenza-like illness (ILI), defined as the occurrence of at least one of the following respiratory symptoms: sore throat, cough, sputum production, wheezing, or difficulty breathing; concurrent with at least one of the following systemic signs or symptoms: temperature >99.0°F, chills, tiredness, headaches or myalgia. Participants were monitored for the occurrence of a respiratory illness by both active and passive surveillance, starting 2 weeks post-vaccination for approximately 7 months. After an episode of respiratory illness, nasopharyngeal swab samples were collected for analysis; attack rates and vaccine efficacy were calculated (see Table 5).

Table 5: Relative Efficacy Against Laboratory-Confirmed Influenza* Regardless of Similarity to the Vaccine Components, Associated with Influenza-Like Illness[†], Adults 65 Years of Age and Older (Study 2[‡])

| | Fluzone High-Dose N[§]=15,892 n[¶] (%) | Fluzone N[§]=15,911 n[¶] (%) | Relative Efficacy % (95% CI) |
|-------------------------------------|---|---|---|
| Any type/subtype[#] | 227 (1.43) | 300 (1.89) | 24.2 (9.7; 36.5) ^p |
| Influenza A | 190 (1.20) | 249 (1.56) | 23.6 (7.4; 37.1) |
| A (H1N1) | 8 (0.05) | 9 (0.06) | 11.0 (-159.9; 70.1) |
| A (H3N2) | 171 (1.08) | 222 (1.40) | 22.9 (5.4; 37.2) |
| Influenza B^β | 37 (0.23) | 51 (0.32) | 27.4 (-13.1; 53.8) |

* Laboratory-confirmed: culture or polymerase-chain-reaction-confirmed

† Occurrence of at least one of the following respiratory symptoms: sore throat, cough, sputum production, wheezing, or difficulty breathing; concurrent with at least one of the following systemic signs or symptoms: temperature >99.0°F, chills, tiredness, headaches or myalgia

‡ NCT01427309

§ N is the number of vaccinated participants in the per-protocol analysis set for efficacy assessments

¶ n is the number of participants with protocol-defined influenza-like illness with laboratory confirmation

Primary endpoint

p The prespecified statistical superiority criterion for the primary endpoint (lower limit of the 2-sided 95% CI of the vaccine efficacy of Fluzone High-Dose relative to Fluzone >9.1%) was met.

β In the first year of the study the influenza B component of the vaccine and the majority of influenza B cases were of the Victoria lineage; in the second year the influenza B component of the vaccine and the majority of influenza B cases were of the Yamagata lineage

A secondary endpoint of the study was the occurrence of culture-confirmed influenza caused by viral types/subtypes antigenically similar to those contained in the respective annual vaccine formulations in association with a modified CDC-defined ILI, defined as the occurrence of a temperature >99.0°F (>37.2°C) with cough or sore throat. The efficacy of Fluzone High-Dose relative to Fluzone for this endpoint was 51.1% (95% CI: 16.8; 72.0).

15 REFERENCES

- 1 Lasky T, Terracciano GJ, Magder L, et al. The Guillain-Barré syndrome and the 1992-1993 and 1993-1994 influenza vaccines. *N Engl J Med* 1998;339:1797-802.
- 2 Hannoun C, Megas F, Piercy J. Immunogenicity and protective efficacy of influenza vaccination. *Virus Res* 2004;103:133-138.
- 3 Hobson D, Curry RL, Beare AS, Ward-Gardner A. The role of serum haemagglutination-inhibiting antibody in protection against challenge infection with influenza A2 and B viruses. *J Hyg Camb* 1972;70:767-777.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Single-dose, prefilled syringe, without needle, 0.7 mL (NDC 49281-226-88) (not made

with natural rubber latex). Supplied as package of 10 (NDC 49281-226-65).

16.2 Storage and Handling

Store Fluzone High-Dose Quadrivalent refrigerated at 2° to 8°C (35° to 46°F). DO NOT FREEZE. Discard if vaccine has been frozen.

Do not use after the expiration date shown on the label.

17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Patient Information).

- Inform the patient or caregiver that Fluzone High-Dose Quadrivalent contains killed viruses and cannot cause influenza.
- Fluzone High-Dose Quadrivalent stimulates the immune system to produce antibodies that help protect against influenza.
- Instruct that annual influenza vaccination is recommended.
- Instruct vaccine recipients and caregivers to report adverse reactions to their healthcare provider and/or to Vaccine Adverse Event Reporting System (VAERS).
- Give the Vaccine Information Statements to recipients or caregivers, which are required by the National Childhood Vaccine Injury Act of 1986 prior to each immunization. These materials are available free of charge at the Centers for Disease Control and Prevention (CDC) website (www.cdc.gov/vaccines).

Fluzone is a registered trademark of Sanofi Pasteur Inc.

Manufactured by:

Sanofi Pasteur Inc.

Swiftwater, PA 18370 USA

Patient Information Sheet Fluzone® High-Dose Quadrivalent Influenza Vaccine

Please read this information sheet before getting Fluzone High-Dose Quadrivalent vaccine. This summary is not intended to take the place of talking with your healthcare provider. If you have questions or would like more information, please talk with your healthcare provider.

What is Fluzone High-Dose Quadrivalent vaccine?

Fluzone High-Dose Quadrivalent is a vaccine that helps protect against influenza illness (flu).

Fluzone High-Dose Quadrivalent vaccine is for people 65 years of age and older.

Vaccination with Fluzone High-Dose Quadrivalent vaccine may not protect all people who receive the vaccine.

Who should not get Fluzone High-Dose Quadrivalent vaccine?

You should not get Fluzone High-Dose Quadrivalent vaccine if you:

- ever had a severe allergic reaction to eggs or egg products.

- ever had a severe allergic reaction after getting any influenza vaccine.

Tell your healthcare provider if you have or have had:

- Guillain-Barré Syndrome (severe muscle weakness) after getting an influenza vaccine.
- problems with your immune system as the immune response may be diminished.

How is Fluzone High-Dose Quadrivalent vaccine given?

Fluzone High-Dose Quadrivalent vaccine is given as an injection into the muscle.

What are the possible side effects of Fluzone High-Dose Quadrivalent vaccine?

The most common side effects of Fluzone High-Dose Quadrivalent vaccine are:

- pain where you got the injection
- muscle ache
- tiredness
- headache

These are not all of the possible side effects of Fluzone High-Dose Quadrivalent vaccine. Ask your healthcare provider about other side effects.

Call your healthcare provider for advice about any side effects that concern you. You may report side effects to the Vaccine Adverse Event Reporting System (VAERS) at 1-800-822-7967 or <https://vaers.hhs.gov>.

Why should I get Fluzone High-Dose Quadrivalent vaccine instead of a standard-dose quadrivalent influenza vaccine?

Among persons 65 years of age and older, Fluzone High-Dose Quadrivalent generated a similar immune response to Fluzone High-Dose and is expected to provide better protection against influenza compared to standard-dose quadrivalent influenza vaccines.

What are the ingredients in Fluzone High-Dose Quadrivalent vaccine?

Fluzone High-Dose Quadrivalent vaccine contains 4 killed influenza virus strains. There is no live influenza virus in Fluzone High-Dose Quadrivalent. Fluzone High-Dose Quadrivalent cannot cause influenza.

Inactive ingredients include formaldehyde and octylphenol ethoxylate.

Manufactured by: Sanofi Pasteur Inc.
Swiftwater, PA 18370 USA

PRINCIPAL DISPLAY PANEL - 0.7 mL Syringe Label

NDC 49281-226-88

Influenza Vaccine
Fluzone[®] High-Dose
Quadrivalent

2025-2026 Formula
No Preservative
0.7 mL Single-Dose

For 65 yrs of age and older
Mfd by: Sanofi Pasteur Inc.

Rx only

For IM Use

000000

Lot
EXP

XXXXXXX
XXXXXXX



NDC 49281-226-88

Influenza Vaccine

Fluzone® High-Dose

Quadrivalent

2025-2026 Formula

R_x only

No Preservative

0.7 mL Single-Dose

For IM Use

For 65 yrs of age and older

Mfd by: **Sanofi Pasteur Inc.**

Fluzone® HD QIV/NDC 49281-226-88

Lot
EXP

XXXXXXX
XXXXXXX



PRINCIPAL DISPLAY PANEL - 0.7 mL Syringe Package

NDC 49281-226-65

FOR ADULTS

65+

Influenza Vaccine
Fluzone® High-Dose
Quadrivalent

2025-2026 Formula

For Intramuscular Use

10 single-dose prefilled syringes 0.7 mL each

Rx only

NDC 49281-226-65

Influenza Vaccine
Fluzone® High-Dose
Quadrivalent
2025-2026 Formula
For Intramuscular Use
10 single-dose prefilled syringes 0.7 mL each

FOR ADULTS
65+

Rx only

DO NOT FREEZE. Store at 2° to 8°C (35° to 46°F).

SHAKE WELL.

FOR INTRAMUSCULAR USE.

See full prescribing information for additional details.

Prepared from influenza viruses propagated in embryonated chicken eggs and inactivated with formaldehyde. A nonionic surfactant (Triton® X-100*) is added during manufacture. This vaccine has been standardized according to USPHS requirements for the 2025-2026 influenza season and is formulated to contain 240 micrograms (mcg) hemagglutinin (HA) per 0.7 mL dose, in the recommended ratio of 60 mcg HA each, representative of the following prototype strains: A/Victoria/4897/2022 IVR-238 (H1N1), A/Croatia/10136RV/2023 X-425A (H3N2), B/Phuket/3073/2013 (B Yamagata lineage), and B/Michigan/01/2021 (a B/Austria/1359417/2021-like virus, B Victoria lineage).

Contains no preservative.

Not made with natural rubber latex.

*Triton® X-100 – Registered trademark of Union Carbide, Co., USA.



522

US Govt License #1725
Manufactured by:
Sanofi Pasteur Inc.
Swiftwater, PA 18370 USA



000000

GTIN: 00000000000000



XXXXXXXXXXXX

XXXXXX
XXXXXX

N/S
EXP
Lot

NDC 49281-226-65

2025-2026
Formula

Influenza Vaccine
Fluzone® High-Dose
Quadrivalent

Rx only

FOR ADULTS
65+

For Intramuscular Use
10 single-dose prefilled syringes **0.7 mL** each



sanofi

FLUZONE HIGH-DOSE QUADRIVALENT NORTHERN HEMISPHERE

influenza a virus a/victoria/4897/2022 ivr-238 (h1n1) antigen (formaldehyde inactivated), influenza a virus a/croatia/10136rv/2023 x-425a (h3n2) antigen (formaldehyde inactivated), influenza b virus b/phuket/3073/2013 antigen (formaldehyde inactivated), and influenza b virus b/michigan/01/2021 antigen (formaldehyde inactivated) injection, suspension

Product Information

| | | | |
|--------------------------------|---------------|---------------------------|---------------|
| Product Type | VACCINE | Item Code (Source) | NDC:49281-226 |
| Route of Administration | INTRAMUSCULAR | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|---|--------------------|
| INFLUENZA A VIRUS A/VICTORIA/4897/2022 IVR-238 (H1N1) ANTIGEN (FORMALDEHYDE INACTIVATED) (UNII: AU5C98U4BB) (INFLUENZA A VIRUS A/VICTORIA/4897/2022 IVR-238 (H1N1) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) - UNII:C46XJT9FQ9) | INFLUENZA A VIRUS A/VICTORIA/4897/2022 IVR-238 (H1N1) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) | 60 ug in 0.7 mL |
| INFLUENZA A VIRUS A/CROATIA/10136RV/2023 X-425A (H3N2) ANTIGEN (FORMALDEHYDE INACTIVATED) (UNII: HQW9FZS4YK) (INFLUENZA A VIRUS A/CROATIA/10136RV/2023 X-425A (H3N2) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) - UNII:SD82XNG5M6) | INFLUENZA A VIRUS A/CROATIA/10136RV/2023 X-425A (H3N2) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) | 60 ug in 0.7 mL |
| INFLUENZA B VIRUS B/PHUKET/3073/2013 ANTIGEN (FORMALDEHYDE INACTIVATED) (UNII: B93BQX9789) (INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) - UNII:9HB0XUS9TM) | INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) | 60 ug in 0.7 mL |
| INFLUENZA B VIRUS B/MICHIGAN/01/2021 ANTIGEN (FORMALDEHYDE INACTIVATED) (UNII: FF9YP4D23C) (INFLUENZA B VIRUS B/MICHIGAN/01/2021 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) - UNII:CQV855H5FG) | INFLUENZA B VIRUS B/MICHIGAN/01/2021 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) | 60 ug in 0.7 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| GELATIN, UNSPECIFIED (UNII: 2G86QN327L) | |
| SODIUM PHOSPHATE, DIBASIC, ANHYDROUS (UNII: 22ADO53M6F) | |
| SODIUM PHOSPHATE, MONOBASIC, ANHYDROUS (UNII: KH7I04HPUU) | |
| WATER (UNII: 059QF0KO0R) | |
| OCTOXYNOL-9 (UNII: 7JPC6Y25QS) | |
| FORMALDEHYDE (UNII: 1HG84L3525) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:49281-226-65 | 10 in 1 PACKAGE | | |
| 1 | NDC:49281-226-88 | 0.7 mL in 1 SYRINGE, GLASS; Type 3: Prefilled Biologic Delivery Device/System (syringe, patch, etc.) | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| BLA | BLA103914 | 07/01/2025 | 06/30/2026 |

Labeler - Sanofi Pasteur Inc. (086723285)

Registrant - Sanofi Pasteur Inc. (086723285)

Establishment

| Name | Address | ID/FEI | Business Operations |
|---------------------|---------|-----------|---|
| Sanofi Pasteur Inc. | | 086723285 | MANUFACTURE(49281-226) , PACK(49281-226) , LABEL(49281-226) |

Revised: 7/2025

Sanofi Pasteur Inc.