POCKET PEPTO PEPTO BISMOL- bismuth subsalicylate tablet Procter & Gamble Manufactura S de RL de CV

POCKET PEPTO Pepto Bismol [®] Bismuth Subsalicylate Upset Stomach Reliever/Antidiarrheal CHEWABLES CHERRY

Drug Facts

Active ingredient (in each tablet)

Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever and antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - belching
 - fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

• allergic to salicylates (including aspirin) • taking other salicylate products

Do not use if

you have

an ulcer

- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- chew or dissolve in mouth
- adults and children 12 years and over: 2 tablets (1 dose) every 1/2 hour or 4 tablets (2 doses) every hour as needed for diarrhea
- 2 tablets (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- each tablet contains: calcium 140 mg, magnesium 9 mg
- salicylate 99 mg
- low sodium

• avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

adipic acid, calcium carbonate, D&C Red No. 27 aluminum lake, FD&C Red No. 40 aluminum lake, flavor, magnesium stearate, mannitol, povidone, saccharin sodium, talc

Questions? 1-800-717-3786 TAMPER EVIDENT: Do not use if the printed inner wrap is broken or missing. MADE IN MEXICO DIST. BY PROCTER & GAMBLE CINCINNATI, OH 45202

PRINCIPAL DISPLAY PANEL - 12 Tablet Carton

РОСКЕТ РЕРТО

Pepto Bismo®

Bismuth Subsalicylate Upset Stomach Reliever/ Antidiarrheal

CHEWABLES

CHERRY

5 SYMPTOMRELIEF

NAUSEA HEARTBURN INDIGESTION UPSET STOMACH DIARRHEA

12 CHEWABLE TABLETS

	POCKET	рерто		TAMPER EVIDENT: Do not use if inner printed seal on bottle is broken or missing.	91633815 MADE IN MEXICO DIST. BY PRIOTER & GAMBLE, Cancimiant of 45202	
	Pep Bismol CHEWAB CHERRY	CVMDTON		Drug Facts	Drug Facts (continued) Step use and six a doctor if Step use and six a doctor if	8
Performance Wardpscom Paters: www.gepto-bismed.com		12 CHEWABLE TABLETS	CONTROL BAP DATE			
	Drug Facts (continued) flow maynesism poviden, search Outestions 1-800-117-3786	⁹				

ΡΟСΚΕΤ ΡΕΡΤΟ ΡΕΡΊ	TO BISMOL					
bismuth subsalicylate tablet						
Product Information						
Product Type	HUMAN OTC DRUG Item Code (Source)				IDC:58933-017	
Route of Administration	ORAL					
Active Ingredient/Active	Moiety					
Ingr	edient Name		Basis of S	trength	Strength	
				BISMUTH SUBSALICYLATE		
UNII.0414PZ4LPZ)			SUBSALICYLA	IE	262 mg	
UNII.0414FZ4LFZ)			SUBSALICYLA	IE		
Inactive Ingredients			SUBSALICYLA	IE		
	Ingredient Name		SUBSALICYLA		rength	
	Ingredient Name		SUBSALICYLA			
Inactive Ingredients			SUBSALICYLA			
Inactive Ingredients ADIPIC ACID (UNII: 76A0JE0FKJ)	KE (UNII: ZK64F7XSTX)		SUBSALICYLA			
Inactive Ingredients ADIPIC ACID (UNII: 76A0JE0FKJ) D&C RED NO. 27 ALUMINUM LAI	KE (UNII: ZK64F7XSTX)		SUBSALICYLA			
Inactive Ingredients ADIPIC ACID (UNII: 76A0JE0FKJ) D&C RED NO. 27 ALUMINUM LAI SACCHARIN SODIUM (UNII: SB8Z	KE (UNII: ZK64F7XSTX) UX40TY)		SUBSALICYLA			
Inactive Ingredients ADIPIC ACID (UNII: 76A0JE0FKJ) D&C RED NO. 27 ALUMINUM LAI SACCHARIN SODIUM (UNII: SB8Z TALC (UNII: 7SEV7J4R1U)	KE (UNII: ZK64F7XSTX) UX40TY) 7XOA)		SUBSALICYLA			
Inactive Ingredients ADIPIC ACID (UNII: 76A0JE0FKJ) D&C RED NO. 27 ALUMINUM LAU SACCHARIN SODIUM (UNII: SB8Z TALC (UNII: 7SEV7J4R1U) FD&C RED NO. 40 (UNII: WZB912	KE (UNII: ZK64F7XSTX) UX40TY) 7XOA) 933)		SUBSALICYLA			
Inactive Ingredients ADIPIC ACID (UNII: 76A0JE0FKJ) D&C RED NO. 27 ALUMINUM LAU SACCHARIN SODIUM (UNII: SB8Z TALC (UNII: 7SEV7J4R1U) FD&C RED NO. 40 (UNII: WZB912 ALUMINUM OXIDE (UNII: LMI2606	KE (UNII: ZK64F7XSTX) UX40TY) 7XOA) 933) 9339FGK)		SUBSALICYLA			
Inactive Ingredients ADIPIC ACID (UNII: 76A0JE0FKJ) D&C RED NO. 27 ALUMINUM LAU SACCHARIN SODIUM (UNII: SB8Z TALC (UNII: 7SEV7J4R1U) FD&C RED NO. 40 (UNII: WZB912 ALUMINUM OXIDE (UNII: LMI2606 CALCIUM CARBONATE (UNII: H0G	KE (UNII: ZK64F7XSTX) UX40TY) 7XOA) 933) 9339FGK)		SUBSALICYLA			

P	roduct Chara	acteristics						
Color		pink	Score	Score		no score		
Shape		ROUND	Size	Size		16mm		
Flavor		CHERRY	Imprint Code	Imprint Code		Pepto;Bismol		
Co	ontains							
P	ackaging							
#	ltem Code	Packag	Package Description		Start	Marketing End Date		
1	NDC:58933- 017-12	1 in 1 CARTON	in 1 CARTON					
1		12 in 1 VIAL, PLASTIC Product	n					
2 NDC:58933- 017-24 2 in 1 C		2 in 1 CARTON	CARTON					
2 12 in 1 VIAL, PLASTIC; Type 0: Not a Combination Product			n					
M	larketing	Information						
	Marketing Category	Application I	Number or Monograph Citation	Marketing Date		Marketing End Date		
		Ig M008						

Labeler - Procter & Gamble Manufactura S de RL de CV (812807550)

Revised: 10/2023

Procter & Gamble Manufactura S de RL de CV