HURRICAINE TOPICAL ANESTHETIC- benzocaine gel Beutlich Pharmaceuticals, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

HurriCaine Gel Pina Colada 1.5oz tube

Active ingredient

Benzocaine 20%

Purpose

Oral Anesthetic

Uses

for the temporary relief of occasional minor irritation and pain, associated with:

- canker sores
- sore mouth and throat
- minor injury of the mouth and gums
- minor dental procedures
- minor irritation of the mouth and gums caused by dentures or orthodontic appliances

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert:Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Sore throat warning:if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or irritation, pain, or redness persists or worsens, see your dentist or doctor promptly.

Do not use

Do not use:

- if the tube has been punctured or previously opened
- for teething
- in children under 2 years of age

When using this product

avoid contact with eyes

Keep out of reach of children.

If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

Directions

Directions:

Do not exceed recommended dosage.

- adults and children 2 years of age and older: apply to the affected area. Allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor.
- children under 12 years of age: should be supervised in the use of the product
- children under 2 years of age: do not use

Other information

• store at 20°-25° C (68°-77° F)

Inactive ingredients

• flavor, polyethylene glycol, sodium saccharin

Questions or comments?

1-800-238-8542

M-F: 8:00 a.m. - 4:30 p.m. ET

Principal Display Panel



Drug Facts

Active ingredient

Purpose

Uses for the temporary relief of odcasional minor irritation and pain, associated with - canker sores - spre mouth and throat - minor injury of the mouth and gums = minor dental procedures = minor irritation of the mouth and gums caused by dentures or orthodontic appliances

Warnings Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: ■ pale, gray, or blue colored skin (cyanosis) = headache = rapid heart rate = shortness of breath dizziness or lightheadedness
 fatigue or lack of energy

Allergy alert: Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.

Do not use ■ for teething ■ in children under 2 years of age

When using this product avoid contact with eyes

Directions do not exceed recommended dosage

adults and children 2 years of age and older

apply to the affected area. Allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor.

children under 12 years of age should be supervised in the use of the product children under 2 years of age do not use

Other information store at 20-25 °C (68-77 °F)

Inactive ingredients flavor, polyethylene glycol, sodium saccharin

Questions or comments? 1-800-238-8542 M-F: 8:00 a.m. - 4:30 p.m. ET

HURRICAINE®

Topical Anesthetic GEL

20% Benzocaine Oral Anesthetic

PINA COLADA

Manufactured for:

Beutlich® Pharmaceuticals, LLC

Bunnell, FL

Made in the USA

1 Tube - NET WT. 1.5 oz. (42.52 g)

HURRICAINE TOPICAL ANESTHETIC

benzocaine gel

Product Information

HUMAN OTC DRUG Item Code (Source) NDC:0283-1820 **Product Type**

Active Ingredient/Active Moiety

Ш	, icare 1119, carein, icare 1 10101,		
	Ingredient Name	Basis of Strength	Strength
	BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

Inactive Ingredients			
Ingredient Name	Strength		
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)			
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			

Product Characteristics		
Color		Score
Shape		Size
Flavor	COCONUT (Pina Colada)	Imprint Code
Contains		

l	P	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
	1	NDC:0283-1820- 27	42.52 g in 1 TUBE; Type 0: Not a Combination Product	08/31/2023		

			Marketing Information			
Marketing Applicat Category	tion Number or Monograph Citation	Marketing Start Date	Marketing End Date			
OTC monograph not part356		08/31/2023				

Labeler - Beutlich Pharmaceuticals, LLC (005209325)

Registrant - Beutlich Pharmaceuticals, LLC (005209325)

Establishment					
Name	Address	ID/FEI	Business Operations		
Dental Technologies, Inc.		148312838	manufacture(0283-1820)		

Revised: 8/2023 Beutlich Pharmaceuticals, LLC