

**COLD AND FLU SEVERE DAYTIME, NIGHTTIME- acetaminophen,
chlorpheniramine maleate, dextromethorphan hbr, guaifenesin,
phenylephrine hcl
Meijer Distribution, Inc.**

Meijer 44-503C473A-08

Active ingredients (in each caplet) (Daytime Cold & Flu Severe)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Active ingredients (in each caplet) (Nighttime Cold & Flu Severe)

Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Antihistamine
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - sore throat
 - headache
 - nasal congestion
 - minor aches and pains
 - sinus congestion and pressure
 - sneezing and runny nose (***Nighttime only***)
- helps clear nasal passages

- relieves cough to help you sleep
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive (**Daytime only**)
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- thyroid disease
- diabetes
- high blood pressure
- heart disease
- glaucoma (**Nighttime only**)
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a breathing problem such as emphysema or chronic bronchitis (**Nighttime only**)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**Nighttime only**)

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children (**Nighttime only**)
- marked drowsiness may occur (**Nighttime only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**Nighttime only**)
- avoid alcoholic beverages (**Nighttime only**)
- use caution when driving a motor vehicle or operating machinery (**Nighttime only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Do not take DAYTIME and NIGHTTIME products at the same time.

Directions

- **do not take more than directed**
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole - do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- **each caplet contains:** sodium 3 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients (Daytime only)

corn starch, crospovidone, D&C yellow #10 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Inactive ingredients (Nighttime only)

corn starch, crospovidone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal display panel

meijer®

NDC 79481-5034-8

**COMBO PACK
24 CAPLETS TOTAL**

<p>COMPARE TO TYLENOL® COLD + FLU SEVERE DAY ACTIVE INGREDIENTS*</p> <p>Daytime COLD & FLU SEVERE ACETAMINOPHEN DEXTROMETHORPHAN HBr GUAIFENESIN PHENYLEPHRINE HCl Pain Reliever/Fever Reducer Cough Suppressant Expectorant Nasal Decongestant Fever, Headache, Sore Throat, Nasal Congestion, Cough, Mucus, Chest Congestion Actual Size 16 DAYTIME CAPLETS</p>	<p>COMPARE TO TYLENOL® COLD + FLU SEVERE NIGHT ACTIVE INGREDIENTS*</p> <p>Nighttime COLD & FLU SEVERE ACETAMINOPHEN CHLORPHENIRAMINE MALEATE DEXTROMETHORPHAN HBr PHENYLEPHRINE HCl Pain Reliever/Fever Reducer Antihistamine Cough Suppressant Nasal Decongestant Fever, Headache, Sore Throat, Runny Nose, Cough, Nasal Congestion Actual Size 8 NIGHTTIME CAPLETS</p>
--	---

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER

UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Do Not Take Daytime and Nighttime Products at the Same Time.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

DIST. BY MEIJER

DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com

*This product is not manufactured or distributed by
Kenvue Inc., owner of the registered trademark
Tylenol® COLD + FLU SEVERE Day & Night.

50844

ORG092250347308



1 3733 97191 7

PLD 4109448

50844 0R092250347308

Tylenol® COLD + FLU SEVERE Day & Night.
Kenner Inc., owner of the registered trademark.
*This product is not manufactured or distributed by

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

No Print/No Varnish
Lot & Exp

Drug Facts **KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

Active ingredients (in each caplet) (Daytime Cold & Flu Severe)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Active ingredients (in each caplet) (Nighttime Cold & Flu Severe)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses ■ temporarily relieves these common cold and flu symptoms:
 ■ cough ■ sore throat ■ headache ■ nasal congestion
 ■ minor aches and pains ■ sinus congestion and pressure
 ■ sneezing and runny nose (Nighttime only)
 ■ helps clear nasal passages
 ■ relieves cough to help you sleep
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive (Daytime only)

Drug Facts (continued)

■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug

TAMPER EVIDENT: DO NOT USE IF PACK AGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

OUR QUALITY GUARANTEE

 DIST. BY MEIJER DISTRIBUTION, INC.
 GRAND RAPIDS, MI 49544
www.meijer.com

meijer

Daytime
COLD & FLU SEVERE

COMBO PACK
24 CAPLETS TOTAL

Nighttime
COLD & FLU SEVERE

meijer

Daytime
COLD & FLU SEVERE

**ACETAMINOPHEN
DEXTROMETHORPHAN HBR
GUAIFENESIN
PHENYLEPHRINE HCl**

Pain Reliever/Fever Reducer
Cough Suppressant
Expectorant
Nasal Decongestant

Fever, Headache,
Sore Throat,
Nasal Congestion,
Cough, Mucus,
Chest Congestion



16
DAYTIME
CAPLETS

Actual Size

**COMPARE TO TYLENOL®
COLD + FLU SEVERE DAY
ACTIVE INGREDIENTS***

**COMPARE TO TYLENOL®
COLD + FLU SEVERE NIGHT
ACTIVE INGREDIENTS***

Nighttime
COLD & FLU SEVERE

**ACETAMINOPHEN
CHLORPHENIRAMINE MALEATE
DEXTROMETHORPHAN HBR
PHENYLEPHRINE HCl**

Pain Reliever/Fever Reducer
Antihistamine
Cough Suppressant
Nasal Decongestant

Fever, Headache,
Sore Throat,
Runny Nose, Cough,
Nasal Congestion



8
NIGHTTIME
CAPLETS

Actual Size

Drug Facts (continued)

contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ liver disease
 ■ thyroid disease ■ diabetes ■ high blood pressure
 ■ heart disease ■ glaucoma (Nighttime only)
 ■ cough that occurs with too much phlegm (mucus)
 ■ difficulty in urination due to enlargement of the prostate gland
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
 ■ a breathing problem such as emphysema or chronic bronchitis (Nighttime only)

Ask a doctor or pharmacist before use if you are
 ■ taking the blood thinning drug warfarin
 ■ taking sedatives or tranquilizers (Nighttime only)

When using this product
 ■ do not exceed recommended dosage
 ■ excitability may occur, especially in children (Nighttime only)
 ■ marked drowsiness may occur (Nighttime only)
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness (Nighttime only)

Drug Facts (continued)

■ avoid alcoholic beverages (Nighttime only)
 ■ use caution when driving a motor vehicle or operating machinery (Nighttime only)

Stop use and ask a doctor if
 ■ nervousness, dizziness, or sleeplessness occur
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ new symptoms occur
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Do not take DAYTIME and NIGHTTIME products at the same time.

Drug Facts (continued)

Directions
 ■ do not take more than directed
 ■ adults and children 12 years and over
 ■ take 2 caplets every 4 hours
 ■ swallow whole – do not crush, chew, or dissolve
 ■ do not take more than 10 caplets in 24 hours
 ■ children under 12 years: ask a doctor

Other information
 ■ each caplet contains: sodium 3 mg
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
 ■ see end flap for expiration date and lot number

Inactive ingredients (Daytime only)
 corn starch, croscopolone, D&C yellow #10 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Drug Facts (continued)

Inactive ingredients (Nighttime only)
 corn starch, croscopolone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments? 1-800-426-9391

*This product is not manufactured or distributed by Kenner Inc., owner of the registered trademark Tylenol® COLD + FLU SEVERE Day & Night.

DIST. BY MEIJER DISTRIBUTION, INC.
 GRAND RAPIDS, MI 49544
www.meijer.com

OUR QUALITY GUARANTEE

 WWW.MEIJER.COM/SUBSCRIPTION

Do Not Take Daytime and Nighttime Products at the Same Time.

50844 ORG092250347308
 R-1214-503A473C-08DF

Meijer 44-503A473C-08

COLD AND FLU SEVERE DAYTIME, NIGHTTIME
 acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, guaifenesin, phenylephrine hcl

kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-5034
---------------------	----------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-5034-8	2 in 1 CARTON	11/08/2024	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1		8
Part 2		4

Part 1 of 2

COLD AND FLU SEVERE DAYTIME

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

Product Information

Item Code (Source)	NDC:79481-9503
---------------------------	----------------

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

MALTODEXTRIN (UNII: 7CVR7L4A2D)
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)
STEARIC ACID (UNII: 4ELV7Z65AP)
SUCRALOSE (UNII: 96K6UQ3ZD4)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	19mm
Flavor	MENTHOL	Imprint Code	44;503
Contains			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/08/2024	

Part 2 of 2

COLD AND FLU SEVERE NIGHTTIME

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl tablet, film coated

Product Information

Item Code (Source)	NDC:79481-9473
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	17mm
Flavor	MENTHOL	Imprint Code	44;473
Contains			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/08/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/08/2024	

Labeler - Meijer Distribution, Inc. (006959555)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(79481-5034) , pack(79481-5034)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		832867894	manufacture(79481-5034)
-------------------------	--	-----------	-------------------------

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(79481-5034)

Revised: 11/2025

Meijer Distribution, Inc.