

**FIRST AID ANTIBIOTIC PLUS BURN RELIEF-
bacitracinzinc,neomycinsulfate,polymyxinbsulfate,pramoxinehci ointment
Trifecta Pharmaceuticals USA LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Globe First Aid Antibiotic Plus Burn Relief

Active Ingredient

Pramoxine HCl

Purpose Section

External Analgesic

Active Ingredient

Polymyxin B Sulfate

Purpose

First Aid Antibiotic

Active Ingredient

Bacitracin Zinc

Purpose

First Aid Antibiotic

Neomycin Sulfate

Purpose

First Aid Antibiotic

Uses

First Aid to help prevent infection and for the temporary relief of pain or discomfort in minor cuts, scrapes and burns.

Warnings

For external Use Only

Do Not Use

Do Not use

- In the eyes
- Over large areas of the body
- If you are allergic to any of the ingredients

Ask Doctor before use if you have

- Deep or puncture wounds
- Animal Bites
- Serious Burns

Stop Use and ask a doctor if

- Condition persists or gets worse
- You need to use longer than 1 week
- A rash or other allergic reaction develops
- symptoms persist for more than 1 week or clear up and occur again within a few days

Directions

Adults and children 2 years of age and older:

- Clean the affected area and dry thoroughly.
- Apply a small amount of this product , an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- May be covered with a sterile bandage

Children under 2 years old: ask a doctor

Inactive Ingredients

White Petrolatum

Questions

Call 1-888-296-9067

Storage Information

- Store at 20° to 25°C (68° to 77°F).
- See carton or tube crimp for lot number and expiration date

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center immediately.

Other Information

Distributed By:

Trifecta Pharmaceuticals USA®

101 NE Third Avenue, Suite 1500

Fort Lauderdale, FL. 33301 USA

www.trifecta-pharma.com

OUTSIDE BOX

Compare to the active ingredients in Neosporin® + Burn Relief

Always wash, apply, and cover your wounds.

WASH
Wash the affected area and dry thoroughly.

APPLY
Apply a small amount of Globe Burn Ointment.

COVER
Cover with a sterile bandage.

DRUG FACTS

Active Ingredients	Purpose
Bacitracin Zinc 500 units	First aid antibiotic
Neomycin Sulfate 3.5 mg	First aid antibiotic
Polymyxin B Sulfate 10,000 units	First aid antibiotic
Pramoxine HCl 10 mg	External analgesic

Uses
First aid to help prevent infection and for the temporary relief of pain or discomfort in minor:
• Cuts • Scrapes • Burns

Warnings
For external use only.
Do not use
• In the eyes
• Over large areas of the body
• If you are allergic to any of the ingredients
Ask a doctor before use if you have
• Deep or puncture wounds
• Animal bites
• Serious burns
Stop use and ask a doctor if
• Condition persists or gets worse
• You need to use longer than 1 week
• A rash or other allergic reaction develops
• Symptoms persist for more than 1 week, or clear up and occur again within a few days
Keep out of the reach of children.
If swallowed, get medical help or contact a Poison Control Center immediately.

Directions
Adults and children 2 years of age and older:
• Clean the affected area and dry thoroughly.
• Apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily.
• May be covered with a sterile bandage.
Children under 2 years of age: ask a doctor

Other information
• Store at 20° to 25°C (68° to 77°F).
• See carton or tube crimp for lot number and expiration date.

Inactive ingredient
White petrolatum

Questions?
Call 1-888-296-5067

* This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Neosporin® + Burn Relief.


EXP
LOT



INNER TUBE

NDC: 69396-138-02

Compare to the active ingredients in Neosporin® + Burn Relief



FIRST AID ANTIBIOTIC
+Burn Relief

Maximum Strength Pain Relieving Ointment

Relieves Pain and Prevents Infection in Minor Cuts, Burns and Scrapes

Triple Antibiotic Infection Protection

Bacitracin Zinc
Neomycin Sulfate
Polymyxin B Sulfate
Pramoxine HCl

NET WT. 2 oz. (57g)

DRUG FACTS

Active ingredients	Purpose
Bacitracin Zinc 500 units	First aid antibiotic
Neomycin Sulfate 3.5 mg	First aid antibiotic
Polymyxin B Sulfate 10,000 units	First aid antibiotic
Pramoxine HCl 10 mg	External analgesic

Uses
First aid to help prevent infection and for the temporary relief of pain or discomfort in minor:
• Cuts • Scrapes • Burns

Warnings
For external use only.

Do not use

- In the eyes
- Over large areas of the body
- If you are allergic to any of the ingredients

Ask a doctor before use if you have

- Deep or puncture wounds
- Animal bites
- Serious burns

Stop use and ask a doctor if

- Condition persists or gets worse
- You need to use longer than 1 week
- A rash or other allergic reaction develops
- Symptoms persist for more than 1 week, or clear up and occur again within a few days

Keep out of the reach of children.
If swallowed, get medical help or contact a Poison Control Center immediately.

Directions
Adults and children 2 years of age and older:

- Clean the affected area and dry thoroughly.
- Apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily.
- May be covered with a sterile bandage.

Children under 2 years of age: ask a doctor

Other information

- Store at 20° to 25°C (68° to 77°F).
- See section on tube stamp for lot number and expiration date.

Inactive ingredient
White petrolatum

Questions?
Call 1-888-288-9067

Distributed by:
Trifecta Pharmaceuticals USA®
101 NE Third Avenue, Suite 1500
FL, Lauderdale, FL 33301 USA
www.trifecta-pharma.com

FIRST AID ANTIBIOTIC PLUS BURN RELIEF

bacitracinzinc,neomycinsulfate,polymxinbsulfate,pramoxinehci ointment

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69396-138
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PRAMOXINE HYDROCHLORIDE (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)	PRAMOXINE HYDROCHLORIDE	10 mg in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ.07J96K)	POLYMYXIN B	10000 [USP'U] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g

BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	500 [USP'U] in 1 g
--	------------	-----------------------

Inactive Ingredients

Ingredient Name	Strength
WHITE PETROLATUM (UNII: B6E5W8RQJ4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69396-138-02	1 in 1 BOX	09/04/2023	
1		57 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	09/04/2023	

Labeler - Trifecta Pharmaceuticals USA LLC (079424163)

Revised: 9/2023

Trifecta Pharmaceuticals USA LLC