

ASPIRIN 81 MG- aspirin tablet, delayed release
VALU MERCHANDISERS COMPANY

1080B-AWS-2024-1015

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

* nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for this product

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-844-292-1112

PRINCIPAL DISPLAY PANEL

Always Save®

LOW DOSE

ASPIRIN 81 MG

PAIN RELIEVER (NSAID)

TALK TO YOUR DOCTOR OR OTHER HEALTHCARE PROVIDER BEFORE USING THIS
PRODUCT FOR YOUR HEART

81 MG EACH

ACTUAL SIZE

500 ENTERIC COATED TABLETS



LOW DOSE
ASPIRIN 81 MG
PAIN RELIEVER (NSAID)

TALK TO YOUR DOCTOR OR OTHER HEALTHCARE PROVIDER BEFORE USING THIS PRODUCT FOR YOUR HEART

81 MG EACH

ACTUAL SIZE

500 ENTERIC COATED TABLETS

DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin 81 mg (NSAID)*Pain reliever
* nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for this product.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

PEEL BACK HERE FOR MORE DRUG FACTS



ALWAYS SAVE



66912-0823-AAA

PROUDLY DISTRIBUTED BY:
ASSOCIATED WHOLESOME GROCERS, INC.
KANSAS CITY, KANSAS 66106

Scan here for more product information.
Call 1-844-292-1112 for more product information.

Aspirin is not appropriate for everyone so be sure to talk to your doctor before you begin an aspirin regimen.

L11080804AWS_R0

Adhesive

Drug Facts (continued)

- Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- Have 3 or more alcoholic drinks every day while using this product
- Take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- store between 20°-25°C (68°-77°F) in a dry place

Inactive ingredients anhydrous lactose, carnaua wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, poly sorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-844-292-1112

ASPIRIN 81 MG
aspirin tablet, delayed release

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-080
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	81 mg
Inactive Ingredients			
Ingredient Name			Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)			
CARNAUBA WAX (UNII: R12CBM0EIZ)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			

ALUMINUM OXIDE (UNII: LMI26O6933)	
BROWN IRON OXIDE (UNII: 1N032N7MFO)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	heart
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-080-04	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	11/01/2023	

Labeler - VALU MERCHANDISERS COMPANY (868703513)