

ASPIRIN- aspirin tablet, coated
VALU MERCHANDISERS COMPANY

1018-AWS-2024-1015

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

* nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours

- children under 12 years: consult a doctor

Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Questions or comments?

1-844-292-1112

PRINCIPAL DISPLAY PANEL

Always Save®

REGULAR STRENGTH

ASPIRIN

PAIN RELIEVER/FEVER REDUCER

(NSAID)

COATED TABLETS

ACTUAL SIZE

300 TABLETS

325 mg EACH



Always Save
REGULAR STRENGTH
ASPIRIN
PAIN RELIEVER/FEVER REDUCER (NSAID)
COATED TABLETS
ACTUAL SIZE
300 TABLETS
325 mg EACH

DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts
Active ingredient (in each tablet) Purpose
Aspirin 325 mg (m) (NSAID)*.....Pain reliever/fever reducer
* nonsteroidal anti-inflammatory drug

Uses temporarily relieves:
 ■ headache
 ■ muscle pain
 ■ menstrual pain
 ■ toothache
 ■ pain and fever of colds
 ■ minor pain of arthritis

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
 ■ hives
 ■ facial swelling
 ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

PEEL BACK HERE FOR MORE DRUG FACTS



Scan here for more product information. Call 1-844-292-1112 for more product information.

PROUDLY DISTRIBUTED BY:
 ASSOCIATED WHOLESALE GROCERS, INC.
 KANSAS CITY, KANSAS 66106

ALWAYS SAVE
 5
 7003869913
 0
 66913-0823-AAA




L101819AWS_RO

Adhesive

STOP PEELING

Drug Facts (continued)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed
 Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 ■ you are taking a diuretic
 ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for: ■ diabetes ■ gout ■ arthritis

Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ pain gets worse or lasts for more than 10 days
 ■ fever gets worse or lasts for more than 3 days
 ■ redness or swelling is present ■ new symptoms occur
 ■ ringing in the ears or loss of hearing occurs
 These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 1 to 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
 ■ children under 12 years: consult a doctor

Other information
 ■ store between 20-25°C (68-77°F) in a dry place

Inactive ingredients corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Questions or comments? 1-844-292-1112

ASPIRIN			
aspirin tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-118
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
Inactive Ingredients			
	Ingredient Name		Strength
	CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)		
	HYPROMELLOSES (UNII: 3NXW29V3WO)		
	CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)		
	MINERAL OIL (UNII: T5L8T28FGP)		
	STARCH, CORN (UNII: O8232NY3S)		

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	TCL;011
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-118-19	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/17/2023	06/30/2028

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/17/2023	06/30/2028

Labeler - VALU MERCHANDISERS COMPANY (868703513)

Revised: 12/2025

VALU MERCHANDISERS COMPANY