

**IBUPROFEN- ibuprofen capsule, liquid filled**  
**Best Choice (Valu Merchandisers Company)**

-----

**Drug Facts**

**Active ingredient (in each capsule)**

**Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\***

**(present as the free acid and potassium salt)**

**\*nonsteroidal anti-inflammatory drug**

**Purpose**

**Pain reliever/fever reducer**

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - slurred speech
  - trouble breathing
  - leg swelling
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**

- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

#### **Other information**

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F).
- swallow whole, do not crush, chew, or dissolve

#### **Inactive ingredients**

FD&C green #3, gelatin, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol solution, white ink

#### **Questions or comments?**

**Call 1-877-753-3935 Monday-Friday 9AM-5PM EST**

#### **Principal Display Panel**

COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® LIQUI-GELS®†

**Ibuprofen** Capsules, 200 mg

Pain Reliever/Fever Reducer (**NSAID**)

Softgels\*\*

\*\* LIQUID FILLED CAPSULES

\*\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

PROUDLY DISTRIBUTED BY:

VALU MERCHANDISERS, CO.

5000 KANSAS AVE

KANSAS CITY, KS 66106

#### **Product Label**

Exp. Date: \_\_\_\_\_

Lot No.: \_\_\_\_\_

TAKE PRECAUTION: DO NOT USE IF PRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.  
KEEP OUT OF REACH OF CHILDREN.  
WARNINGS AND PRODUCT INFORMATION.  
FD004794

PROUDLY DISTRIBUTED BY:  
WALL MERCHANTS, CO.  
5000 KANSAS AVE  
KANSAS CITY, KS 66106

Best Choice  
100% Guaranteed  
www.bestchoicebrand.com

0 70038 63444 7

**COMPARE TO THE ACTIVE INGREDIENT IN  
ADVIL® LIQUI-GELS®†**

**Ibuprofen**  
Capsules, 200 mg

**Pain Reliever/  
Fever Reducer (NSAID)**

**20 Softgels\*\***  
\*\* LIQUID FILLED CAPSULES

**Best Choice®**

**BEST CHOICE Ibuprofen 200 mg Softgels**

**Drug Facts (continued)**

■ you have symptoms of heart problems or stroke:  
■ chest pain ■ slurred speech ■ trouble breathing  
■ leg swelling ■ weakness in one part or side of body  
■ pain gets worse or lasts more than 10 days ■ fever  
■ gets worse or lasts more than 3 days ■ redness or  
■ swelling is present in the painful area ■ any new  
■ symptoms appear

**If pregnant or breast-feeding**, ask a health professional  
before use. It is especially important not to use ibuprofen  
during the last 3 months of pregnancy unless definitely  
directed to do so by a doctor because it may cause  
problems in the unborn child or complications during  
delivery. **Keep out of reach of children.** In case of  
overdose, get medical help or contact a Poison Control  
Center right away (1-800-222-1222).

**Directions** ■ do not take more than directed  
■ the smallest effective dose should be used  
■ adults and children 12 years and over: ■ take  
1 capsule every 4 to 6 hours while symptoms persist  
■ if pain or fever does not respond to 1 capsule, 2  
capsules may be used ■ do not exceed 6 capsules in  
24 hours, unless directed by a doctor ■ children under  
12 years: ask a doctor

**Other information** ■ each capsule contains:  
potassium 20 mg ■ read all warnings and directions  
before use. Keep carton.

**Drug Facts (continued)**

product ■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except  
aspirin, increase the risk of heart attack, heart failure, and  
stroke. These can be fatal. The risk is higher if you use  
more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction  
to any other pain reliever/fever reducer ■ right before  
or after heart surgery

**Ask a doctor before use if** ■ the stomach bleeding  
warning applies to you ■ you have problems or serious  
side effects from taking pain relievers or fever reducers  
■ you have a history of stomach problems, such as  
heartburn ■ you have high blood pressure, heart  
disease, liver cirrhosis, kidney disease, asthma, or had a  
stroke ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**  
■ under a doctor's care for any serious condition  
■ taking aspirin for heart attack or stroke, because  
ibuprofen may decrease this benefit of aspirin  
■ taking any other drug

**When using this product**, take with food or milk if  
stomach upset occurs.

**Stop use and ask a doctor if** ■ you experience any of  
the following signs of stomach bleeding: ■ feel faint  
■ vomit blood ■ have bloody or black stools ■ have  
stomach pain that does not get better

**Drug Facts (continued)**

■ store at 20° to 25°C (68° to 77°F)  
■ avoid excessive heat above 40°C (104°F)  
■ swallow whole; do not crush, chew, or dissolve

**Inactive ingredients** FD&C green #3, gelatin,  
medium-chain triglycerides, polyethylene glycol,  
potassium hydroxide, purified water, sorbitol sorbitan  
solution, white ink

**Questions or comments?** Call  
1-877-753-3935 Monday-Friday 9AM-5PM EST

†This product is not manufactured or distributed by  
Pfizer Consumer Healthcare, distributor of Advil®  
Liqui-Gels®.

IBUPROFEN			
ibuprofen capsule, liquid filled			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-745
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

**Inactive Ingredients**

Ingredient Name	Strength
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	

**Product Characteristics**

Color	BLUE	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	PC16
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-745-40	1 in 1 BOX	01/31/2018	
1		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:63941-745-20	1 in 1 BOX	01/31/2018	
2		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206568	01/31/2018	

**Labeler** - Best Choice (Valu Merchandisers Company) (868703513)