

VICKS FORMULA 44DM COUGH- dextromethorphan hbr liquid
The Procter & Gamble Manufacturing Company

VICKS® FORMULA 44™ DM COUGH Liquid

Drug Facts

Active ingredient (in each 15 mL)

Dextromethorphan HBr 15 mg

Purpose

Cough suppressant

Use

temporarily relieves cough due to minor throat and bronchial irritation associated with a cold

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Stop use and ask a doctor if

cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hours

| | |
|---------------------------------------|-----------------------|
| adults and children 12 years and over | 30 mL every 6-8 hours |
| children 6 to under 12 years | 15 mL every 6-8 hours |
| children 4 to under 6 years | ask a doctor |
| children under 4 years | do not use |

Other information

- **each 15 mL contains:**sodium 12 mg
- store at no greater than 25°C
- does not meet USP requirements for light resistant packaging

Inactive ingredients

citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: Do not use if printed shrinkband is broken or missing.

**DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202**

PRINCIPAL DISPLAY PANEL - 354 ml Bottle Label

VICKS®

FORMULA

44™ DM

COUGH

Dextromethorphan HBr -cough suppressant

ATTACKS COUGH AT THE SOURCE

Non-Drowsy

12 FL OZ (354 ml)



Drug Facts (continued)

Other information

- each 15 mL contains: sodium 12 mg
- store at no greater than 25°C
- does not meet USP requirements for light resistant packaging

Inactive ingredients citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions? 1-800-362-1683

Drug Facts (continued)

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Stop use and ask a doctor if cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hours

| | |
|---------------------------------------|-----------------------|
| adults and children 12 years and over | 30 mL every 6-8 hours |
| children 6 to under 12 years | 15 mL every 6-8 hours |
| children 4 to under 6 years | ask a doctor |
| children under 4 years | do not use |



VICKS FORMULA 44DM COUGH

dextromethorphan hbr liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:69423-809 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------------------|-------------------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 15 mg in 15 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| SODIUM CITRATE (UNII: 1Q73Q2JULR) | |
| SORBITOL (UNII: 506T60A25R) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |

XANTHAN GUM (UNII: TTV12P4NEE)

Product Characteristics

| | | | |
|-----------------|-------|---------------------|--|
| Color | red | Score | |
| Shape | | Size | |
| Flavor | BERRY | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:69423-809-12 | 354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 12/21/2020 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 12/21/2020 | |

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 12/2025

The Procter & Gamble Manufacturing Company