VICKS FORMULA 44DM NIGHTTIME COUGH AND COLD- acetaminophen, dextromethorphan hbr, chlorpheniramine maleate liquid Procter & Gamble Manufacturing Company

VICKS [®]
FORMULA 44™ DM
NIGHTTIME COUGH & COLD

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg
Chlorpheniramine maleate 4 mg
Dextromethorphan HBr 30 mg

Purpose

Pain reliever/Fever reducer Antihistamine Cough suppressant

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters

Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 30 mL dose cup contains:sodium 25 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: Do not use if printed shrinkband is broken or missing.

DIST. BY PROCTER & GAMBLE, CINCINNATI OH 45202

PRINCIPAL DISPLAY PANEL - 354 ml Bottle Label

VICKS ®

FORMULA 44™ DM

Nighttime COUGH & COLD

Acetaminophen

Dextromethorphan HBr,

Chlorpheniramine Maleate

Fever, Minor Aches & Pains

Sneezing, Runny Nose 12 FL OZ (354 mL)





Drug Facts (continued)

Warnings
Liver warning: This product contains acetaminophen. Severe liver

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take among the first product of the severe with other drugs containing acetaminophen and other drugs containing acetaminophen and or drugs containing acetaminophen and other drugs drugs drugs and other drugs dr

- o not use

 with any other drug containing acetaminophen (prescription or norprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are not sulleng a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for despression, psychiatric or emotional conditions, or Parkinso's disease), or lot 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

 Adv. a doctor before ask if you have

- Ask a doctor before use if you have

 liver disease glaucoma
 ough that occurs with too much phlegm (mucus)
 a breathing problem or chronic cough that lasts or as occur
 with smoking, asthma, chronic bronchis or emphysema
 trouble urinating due to enlarged prostate gland
- Ask a doctor or pharmacist before use if you are

 taking sedatives or tranquilizers

 taking the blood thinning drug warfarin
- When using this product

 excitability may occur, especially in children

 marked drowsiness may occur
- avoid alcoholic drinks
 be careful when driving a motor vehicle or operating machinery
 • alcohol, sedatives, and tranquilizers may increase

 ◀ drowsiness

Drug Facts (continued)

- Urtig Facts (communed)

 Stop use and ask a doctor if

 pain or cough jets worse or lasts more than 7 days

 feren gets worse or lasts more than 3 days

 redness or swelling is present

 new symptoms occur

 or cough comes back or occurs with rash or headache that lasts
 see could be signs of a serious condition.

I reset vulnu us signis or a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Polson Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
 only use the dose cup provided
 do not exceed 4 doses per 24 hrs
 adults & children 12 yrs & over | 30 mL every 6 hrs

ildren under 4 yrs do not use	or	_

Other information
• each 30 mL contains: sodium 25 mg
• store at no greater than 25°C and do not refrigerate

Inactive ingredients citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralos xanthan gum

Questions? 1-800-362-1683

VICKS FORMULA 44DM NIGHTTIME COUGH AND COLD

acetaminophen, dextromethorphan hbr, chlorpheniramine maleate liquid

Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:69423-811 **Route of Administration ORAL**

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 15 mL	
CHLORPHENIRAMINE MALEATE (UNII: V1Q0090J9Z) (CHLORPHENIRAMINE - UNII: 3U6I01965U)	CHLORPHENIRAMINE MALEATE	4 mg in 15 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 15 mL	

Inactive Ingredients	
Ingredient Name	Strength

XANTHAN GUM (UNII: TTV12P4NEE)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
SORBITOL (UNII: 506T60A25R)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	

Product Characteristics			
Color	red	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

l	P	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
	1	NDC:69423- 811-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/25/2021		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	01/25/2021		

Labeler - Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2023 Procter & Gamble Manufacturing Company