

**DOLEX FLEX- ibuprofen capsule, liquid filled**  
**Pharmadel LLC**

-----  
**DOLEX FLEX (HHH)**

**Drug Facts**

**Active Ingredient & Purposes**

---

**Active ingredient (in each capsule)**

Solubilized ibuprofen equal to  
200 mg ibuprofen (**NSAID\***).....  
(present as the free acid and potassium salt)  
\* nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/ fever reducer

---

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain from arthritis
- and to reduce fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach
- bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- slurred speech
- weakness in one part or side of body
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**

---

**adults and children 12 years and over**

- take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor

**children under 12 years**

- ask a doctor
- 

**Other information**

- each capsule contains: **20mg potassium**
- store at 68-77°F (20-25°C)
- avoid excessive heat above 104°F (40°C)

**Inactive ingredients**

FD&C blue #1, gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitol and sorbitan

**Questions or comments?**

**1-866-359-3478** (M-F) 9 AM - 5 PM EST or [www.pharmadel.com](http://www.pharmadel.com)

**Package Principal Display Panel**

NDC 55758-014-20

Compare to Advil® Liqui-Gels  
active ingredients

# DOLEX<sup>®</sup> FLEX

**IBUPROFEN**  
Capsules, 200 mg

*Fast relief of Muscular Aches*

*Pain Reliever/ Fever Reducer (NSAID)*



**20** Liquid Filled  
Capsules

## DOLEX FLEX

ibuprofen capsule, liquid filled

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55758-014
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SORBITAN</b> (UNII: 6O921CV9RU)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	

**POLYETHYLENE GLYCOL, UNSPECIFIED** (UNII: 3WJQ0SDW1A)


**GELATIN** (UNII: 2G86QN327L)

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	score with uneven pieces
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	IB1
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55758-014-20	1 in 1 CARTON	09/07/2023	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55758-014-99	50 in 1 CARTON	11/01/2023	
2	NDC:55758-014-02	2 in 1 PACKET; Type 0: Not a Combination Product		



### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078682	09/01/2023	

**Labeler** - Pharmadel LLC (030129680)